



COVERS UP TO 5,000 SQUARE FEET
464.5 m²

26-0-12
MONEY BACK GUARANTEE[†]
— Garantía de devolución de dinero —

NET WT 14lb
6.35 kg



PHOSPHORUS FREE Winterizer
Weed & Feed
WEED CONTROL + FERTILIZER



PROOF OF PURCHASE
0 21496 01776 1

PHOSPHORUS FREE Winterizer
Weed & Feed

WEED CONTROL + FERTILIZER

Herbicida y abono con acondicionador invernal sin fósforo

**ENCOURAGES
EARLY SPRING GREEN-UP**

Estimula el reverdecimiento a principios de la primavera

KILLS 250+ LISTED WEEDS

INCLUDING HENBIT, DOLLARWEED, WHITE CLOVER, PLANTAIN & CHICKWEED

Mata más de 250 tipos de malezas, incluidas ortiga mansa, acaricoba, trébol blanco, plántago y pamplina



**FEEDS FOR UP TO
3 MONTHS**

Abona de hasta 3 meses

COVERS UP TO 5,000 SQUARE FEET
464.5 m²

26-0-12
MONEY BACK GUARANTEE[†]
Garantía de devolución de dinero[†]

NET WT 14lb
6.35 kg

KEEP OUT OF REACH OF CHILDREN MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS

CAUTION
PRECAUCIÓN

See back panel for complete Precautionary Statements, including First Aid and Directions For Use, including Storage and Disposal.

Consulte el panel posterior para conocer las declaraciones de precaución, incluidos los primeros auxilios y las instrucciones de uso, almacenamiento y eliminación.

THIS PRODUCT CONTAINS:

14.70 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.735% CAS 1928-43-4.
3.34 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.167% CAS 16484-77-8.
1.42 lb 3,6-dichloro-o-anisic acid equivalent per ton or 0.071% CAS 1918-00-9.
Isomer Specific by AOAC Methods

ACTIVE INGREDIENTS:
2,4-D, 2-ethylhexyl ester . . . 1.108%
Mecoprop-p acid . . . 0.167%
Dicamba acid . . . 0.071%
OTHER INGREDIENTS . . . 98.654%
TOTAL . . . 100.000%



PROOF OF PURCHASE
0 21496 01776 1



COVERS UP TO 5,000 SQUARE FEET
464.5 m²

26-0-12
MONEY BACK GUARANTEE[†]
— Garantía de devolución de dinero —

NET WT 14lb
6.35 kg

PHOSPHORUS FREE Winterizer
Weed & Feed
WEED CONTROL + FERTILIZER

Herbicida y abono con acondicionador invernal sin fósforo



PHOSPHORUS FREE Winterizer
Weed & Feed

WEED CONTROL + FERTILIZER

Herbicida y abono con acondicionador invernal sin fósforo



PROOF OF PURCHASE
0 21496 01776 1

PHOSPHORUS FREE

Winterizer Weed & Feed

WEED CONTROL + FERTILIZER

GUARANTEED ANALYSIS

F525

Total Nitrogen (N) 26.0%
1.2% Ammoniacal Nitrogen
24.8% Urea Nitrogen*
Soluble Potash (K₂O) 12.0%
Sulfur (S) 9.0%
1.4% Combined Sulfur (S)
7.6% Free Sulfur

Derived from: Urea, Muriate of Potash, Polymer-Coated Urea, Sulfur and Ammonium Sulfate.

*4.5% Slow-Release Urea Nitrogen From Polymer-Coated Urea.

FIRST AID

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this product, contact the National Pesticide Information Center, 1-800-858-7378, Monday-Friday, 8:00 AM to 12:00 PM PST. You may also contact the National Poison Control Center, 1-800-222-1222, day or night, for emergency medical treatment information. FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure or accident call CHEMTREC 1-800-424-9300.

ENVIRONMENTAL HAZARDS This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Do not use on dichondra, carpetgrass, 'Floratum' St. Augustinegrass, creeping bentgrass, or turfgrass where desirable clovers are present.
- Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees. Do not exceed specified dosage for any area and do not apply this product within the dripline of, or over surface roots of trees, shrubs, and other ornamentals.
- Do not use this product on or near desirable plants and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of the roots of desirable trees and shrubs. Since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not apply water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

INSTRUCTIONS:

This product is a combination of three proven weed killers--2,4-D, mecoprop-p, and dicamba. This herbicide controls a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, and spurge In addition, a 26-0-12 fertilizer provides a supply of plant nutrients.

USE PRECAUTIONS:

- Do not apply this product to newly seeded grasses until well-established or after the second or third mowing.
- Reseeding or overseeding areas may occur three (3) weeks after the application of this product.

HOW MUCH TO APPLY:

For use on: Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, bahiagrass, zoysiagrass, St. Augustinegrass, centipedegrass, hybrid bermudagrass, and colonial bentgrass: Apply 2.8 pounds of product per 1,000 square feet of turfgrass. A second application may be made in approximately 30 days. The maximum number of broadcast applications per treatment site is 2 per year.

Do not apply more than 4 pounds of product per 1,000 square feet of turfgrass per application. If necessary, a second application may be made no sooner than 30 days after the first. Do not apply more than 8 pounds of product per 1,000 square feet per year.

WHEN TO APPLY:

Apply when broadleaf weeds are actively growing in spring or early fall.

Apply when the air is calm (Choose a calm day) to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery.

Possible retreatment may be necessary if lawn is heavily overrun by weeds, if adverse low moisture conditions prevail, or if weeds are in a state of poor growth.

APPLICATION INSTRUCTIONS:

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).
3. Apply when broadleaf weeds are young and actively growing, preferably in the morning when dew is on the grass. At the time of application, moisture on the weed leaves from dew, rainfall or irrigation/watering aids in control. If grass is not moist at time of application, sprinkle lightly with water before applying product.
4. Avoid applying when temperatures are consistently above 90°F.
5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

SPREADER SETTINGS

BROADCAST SPREADERS

RATE

STA-GREEN® Basic	4 1/2
SCOTTS® SPEEDYGREEN™	4 1/2
SCOTTS® Basic	4 1/2
SCOTTS® Lawn Pro	4 1/2
SCOTTS® DELUXE EDGEGUARD®	4 1/2
SCOTTS® EDGEGUARD® Mini	4 1/2
Precision Rotary	5
EARTHWAY® EV-N-SPRED®	13

DROP SPREADER

RATE

SCOTTS® ACCUGREEN®	6
--------------------	---

HOW TO APPLY

SPREADERS & CALIBRATION

Always apply with a rotary or drop spreader to distribute the granules evenly. Never apply by hand.

COVERAGE: To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to 5,000 square feet.

IMPORTANT: It is important to follow directions as indicated on this quality lawn care product as overapplication may cause injury to your lawn. Spreader settings are approximate.

Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult Owner's Manual. Never apply by hand.

SINGLE PASS DISTRIBUTION:

- For rotary spreaders: Operate at intervals corresponding to the swath width.
- For drop spreaders: Overlap the wheel tracks to ensure complete coverage.
- For best results, apply this product uniformly and avoid streaks, skips or overlaps. Walk at a steady pace to ensure even coverage. Close the spreader opening when turning to change directions or when stopping.

CALIBRATION INSTRUCTIONS AND SPREADER SETTINGS:

Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended rate under actual application conditions. To provide proper distribution, calibrate spreader before application.

• For 2.8 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 357 square feet and adjust to apply at the rate of 1 pound of product per 357 square feet.

BROADCAST SPREADERS: For best results, apply in a circular path starting on the outside and working toward the center overlapping slightly to ensure complete coverage.

DROP SPREADERS: Apply 2 strips along each end. Then apply along length of lawn, overlapping slightly and shutting off the spreader before each turn at edge of border strips.

†MONEY BACK GUARANTEE

If you are not completely satisfied with this product, please mail your name, address and proof of purchase (package UPC) and a brief explanation for seeking a refund.

GARANTÍA DE DEVOLUCIÓN DE SU DINERO:

Si no está completamente satisfecho con este producto, envíenos por correo su nombre, dirección, el comprobante de la compra (UPC del empaque) y una breve explicación del motivo de la solicitud de reembolso.

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

WEEDS CONTROLLED

Amaranth, purple Amaranth, slender Aster, heath Aster, white heath Aster, purple Aster, white prairie Artichoke Austrian fieldcress Bedstraw, Catchweed Bedstraw, smooth Bedstraw, Northern Bedstraw, pineywoods Beggarticks, hairy Betony, Florida Bindweed, Field Bindweed, Hedge Bitter Wintercress Bittercress, hairy Bitterweed Black-eyed Susan Black Medic Blood flower milkweed Blue lettuce Brassbuttons Bristly oxtongue Broomweed Buckhorn Bullnettle Burclover Burdock Burweed, lawn (spurweed) Buttercup, bulbous Buttercup, creeping Buttonweed, common (poorjoe) Buttonweed, Virginia Carolina geranium (wild) Carpetweed Catnip Catsear, spotted (false dandelion) Chamber bitter Chickweed, common Chickweed, mouseear Chickweed, sticky Chicory Cinquefoil Clover, crimson Clover, alcyce Clover, hop Clover, rabbit foot Clover, red	Clover, strawberry Clover, yellow sweet Clover, white sweet Clover, white Cockle Cocklebur Cocklebur, oriental Cocklebur, spiny Common mullein Corn Spurry Creeping Beggarweed Creeping Jenny (creeping charlie) Cupid shaving Brush Cudweed Daisy, English Daisy, oxeye Dandelion, common Dandelion, false Day flower Deadnettle Dichondra Dock, broadleaf Dock, curly Dogbane Dogfennel Dollardweed (Pennywort) Dove weed Elderberry Falseflax False sunflower Fiddleneck Filaree, redstem Filaree, whitestem Fleabane, daisy (annual) Fleabane, rough Galinsoga, hairy Galinsoga, smallflower Goldenrod Ground ivy Groundsel Gumweed Hairy fleabane Hawkweed, meadow Hawkweed, mouseear Hawkweed, orange Hawkweed, yellow Heallall Heartleaf drymary Hemp Henbit Hoary cress	Horsenettle Horseweed Jimsonweed Johnny-Jumpup violet Knawel Knotweed, prostrate Kochia Lambsquarters Lespedeza Mallow, alkali Mallow, bristly Mallow, common Mallow, dwarf Mallow, venice Marcela Matchweed Mexicanweed Moneywort Morningglory, bigroot Morningglory, ivyleaf Morningglory, Mexican Morningglory, red Morningglory, tall Morningglory, woolly Mugwort Mustard, ball Mustard, black Mustard, blue Mustard, hedge Mustard, Indian, Mustard, tansy Mustard, tumble Mustard, white Mustard, wormseed Nettle, stinging (burning) Nettle, tall Nettle, wood Parsley-piert Parsnip Pearwort Pennycress (frenchweed) Pepperweed, perennial Pepperweed, virginia (Peppergrass) Pigweed, prostrate Pigweed, redroot Pigweed, russian Pigweed, smooth Pigweed, tumble Pineapple weed Plains coreopsis (tickseed) Plantain, blackseed	Plantain, bracted Plantain, broadleaf Plantain, buckhorn Plantain, hoary Plantain, narrowleaf Plantain, slender Plantain, woolly Poison ivy Poison oak Pokeweed Prairie Sunflower Prickly lettuce (compass plant) Prickly sida Prostrate knotweed Puncturevine Purslane, common Pusley, brazil Pusley, florida Pusley, large flower Ragweed, bur Ragweed, common Ragweed, lanceleaf Ragweed, western Redstem filaree Rough cinquefoil Scarlet pimpernel Shepherd's purse Smartweed, ladysthumb Smartweed, pale Smartweed, pennsylvania Smooth chaff-flower Smoothdock Sorrel, red (sheep) Sowthistle, annual Sowthistle, spiny Spanish needles Speedwell, bi-lobed Speedwell, common Speedwell, corn Speedwell, ivyleaf Speedwell, persian Speedwell, purslane Speedwell, slender Speedwell, snow Speedwell, thymeleaf Spiny amaranth Spurge, spotted Spurge, prostrate Spurweed Star of Bethlehem Strawberry, India mock	Tansy ragwort Tanweed Thistle, blessed Thistle, bull Thistle, canada Thistle, fiodman (prairie) Thistle, musk Thistle, plate Thistle, plumeless Thistle, russian Thistle, scotch Thistle, wavyleaf Thistle, yellow (horrible) Trailing crownvetch Velvetleaf Vervain, blue Vervain, hoary Vervain, prostrate Vervain, tall Vetch, bird Vetch, hairy Vetch, milk Vetch, narrowleaf Vetch, wild Virginia creeper Western clematis Western salsify Wild aster Wild buckweat Wild carrot Wild four-o'clock Wild garlic Wild lettuce Wild marigold Wild mustard (yellow rocket) Wild onion Wild pansnip Wild radish Wild rape Wild strawberry Wild sweet potato Wild violet Woodsorrel, creeping Woodsorrel, violet Woodsorrel, yellow (Oxalis) Woolly croton Wormseed Yarrow, common Yarrow, western Yarrow, yellow
--	--	---	---	--

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container.

If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Keep this product off sidewalks, patios, driveways, etc. This fertilizer may stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders, which scatter fertilizer beyond lawn edge and onto concrete.

Apply only to actively growing turf. Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands in the driveway, sidewalk, or street, back onto your lawn. Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use.

EPA Reg. No. 2217-884-59144

EPA Est. No. 59144-GA-1 (A), 32802-MO-1 (B), 37351-CO-1 (C), 8378-IN-1 (D), 9404-FL-1 (E), 961-PA-01 (F), 961-IL-01 (G), 35497-OR-1 (H), 49991-TX-2 (Q), 69759-OR-002 (R), 7401-TX-01 (S)

When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials. Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.html>

LIMITED WARRANTY AND DISCLAIMER FOR USE ONLY AS DIRECTED. THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

¡DETÉNGASE! LEA LA ETIQUETA COMPLETA PRIMERO. CUMPLA CON TODAS LAS PRECAUCIONES Y SIGA LAS INSTRUCCIONES DETENIDAMENTE.

DECLARACIONES DE PRECAUCIÓN

RIESGOS PARA SERES HUMANOS Y ANIMALES DOMÉSTICOS

PRECAUCIÓN: Provoca una irritación ocular moderada. Evite el contacto con los ojos o la ropa. Lávese cuidadosamente con jabón y agua después de manipular el producto y antes de comer, beber, masticar chicle, usar tabaco a usar el baño.

CUÁNDO APLICAR

• Aplique cuando las hierbas de hojas anchas estén en pleno crecimiento en primavera o a principios del otoño.

• Aplique cuando no haya brisa para asegurar una cobertura uniforme y evitar que se esparzan los gránulos sobre flores, vegetales y arbustos ornamentales.

• Podría ser necesario un nuevo tratamiento si el césped está excesivamente invadido por malezas; si prevalecen condiciones adversas de humedad baja o si las malezas están en un período de crecimiento pobre.

CUÁNTO APLICAR

Para uso en césped azul de Kentucky, festuca spp., ballico perenne, césped Bermuda común, césped bahía, césped zoysia, césped de San Agustín, césped ciempiés, césped Bermuda híbrido y césped agrostis colonial. Aplique 1,17 kg del producto por cada 92,9 m² de césped. Se puede realizar una segunda aplicación en aproximadamente 30 días. El número máximo de aplicaciones por dispersión por sitio de tratamiento es de 2 por año.

No aplique más de 1,81 kg del producto por cada 92,9 m² de césped cada vez. Si es necesario, se puede aplicar por segunda vez 9 días o menos 30 días después de la primera aplicación. No aplique más de 3,62 kg del producto por cada 92,9 m² al año.

INSTRUCCIONES DE APLICACIÓN

1. Corte el césped a una altura normal 1 a 2 días antes de la aplicación.
2. Riegue bien el césped al menos 1 a 2 días antes de la aplicación para mantener la humedad hasta el siguiente riego (consulte el paso 5).
3. Aplique cuando las malezas de hojas anchas sean jóvenes y estén en pleno crecimiento, de preferencia por la mañana cuando haya ración sobre el césped. En el momento de la aplicación, la humedad en las hojas de las hierbas proveniente del rocío, lluvia o riego/regado ayuda en el control. Si el césped no está húmedo al momento de la aplicación, rocíelo ligeramente con agua antes de aplicar el producto.
4. Evite aplicar el producto cuando las temperaturas sean constantes sobre los 32,2 °C.
5. No moje las hojas de las malezas por 1 a 2 días después de la aplicación. Tras ese período, se debe realizar un riego completo.

SCOTTS®, ACCUGREEN®, and EDGEGUARD® are registered trademarks of OMS Investments, Inc.
SPEEDYGREEN™ is a trademark of Everris International B.V.
EARTHWAY® and EV-N-SPRED® are registered trademarks of Earthway Products, Inc.
STA-GREEN® is a registered trademark of LF, LLC. All rights reserved

Distributed & Guaranteed By:
Gro Tec, Inc.
P.O. Box 290
Madison, GA 30650

COVERS UP TO 5,000 SQ FT
NET WT 14 LB (6,35 KG)

SGF13v14-2/20

PROOF OF PURCHASE

0 21496 01776 1