

63287

AUG 11 2015 **NEW** [Signature]  
DATE REVIEWER

OR 42 750-196-89819



**PROMIS<sup>TM</sup>**  
**MULTI-PURPOSE FUNGICIDE**

**PROTECTS FRUITS AND VEGETABLES**

---

ACTIVE INGREDIENT:  
Myclobutanil.....2.0 %

OTHER INGREDIENTS:.....98.0 %

TOTAL:.....100.0 %

**STOP - Read the entire label before use**

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See back panel and fold out label for additional precautionary statements and direction for use.

**FOR HOME USE ONLY**

EPA Reg. No 42750-196-89819      EPA Est. No 53883 - TX - 002  
Net Contents: 1 Quart (32 fl. oz) (946 ml)

FIRST AID:	
<b>IF IN EYES:</b>	Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a Poison Control Center or doctor for treatment advice.
<b>IF SWALLOWED:</b>	Call a Poison Control Center or doctor immediately for treatment advice. Have person sip glass of water if able to swallow. Do not induce vomiting unless told to by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For transportation or medical emergency information involving this product, please call (800) 424 - 9300.	
<b>PRODUCT FACTS</b>	
<p><b>What it Does:</b> Promis Multi-Purpose Fungicide prevents and controls anthracnose, brown patch, copper spot, dollar spot, leaf spot (including septoria, zonate), melting out, crown rot, leaf smuts, powdery mildew, rust, red thread, Spring dead spot, necrotic ring spot and Summer patch.</p> <p><b>Where to Use:</b> Outdoors on roses, flowers, ornamental shrubs and trees, fruit trees and vegetables.</p> <p><b>When to Use:</b> Start treating when signs of disease are first noticed.</p> <p><b>NOTE:</b> This product is non-staining to most home siding depending on age and cleanliness. However, before using in areas where the spray may contact home siding (vinyl siding in particular), test in an inconspicuous area and recheck in a few hours. Do not use if any staining is observed.</p>	

**STOP. READ ENTIRE LABEL  
BEFORE USE.**

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. For home use only. Shake well before using. Keep pets and people other than applicator off treated areas until spray has dried.



**WHEN TO USE**

This product is highly effective when it is used to prevent diseases or applied at the first sign of disease. Start when signs of diseases are first noticed. For best disease control, treat every 2 weeks.

**RESTRICTIONS**

- Do not apply more than 13 treatments per year. In Nassau and Suffolk Counties in New York State, do not apply more than 3 times per given area per year.
- Do not treat plants during periods of stress, i.e. heat, drought, flooding, insects, excessive fertilizer, herbicides or pesticide use, etc.
- For application to individual plants only use conventional pump up sprayers at labeled rates.
- If it rains before product has dried, reapply.
- If disease is already present, this product will not eliminate it.

If you need more information, contact your local County Extension Service.

## TO PROTECT FRUITS

### APPLE AND MAYHAW

controls powdery mildew, rusts and scab

Mix 1/2 fl oz of Promis Multi-Purpose Fungicide with 1 gallon of water and uniformly apply to all parts of the tree to point of runoff. Reapply every 7 to 10 days so long as disease symptoms persist. For powdery mildew and rusts, apply at pink bud stage when the buds are opening and the pink petals are visible. For scab, apply Promis Multi-Purpose Fungicide when the first green leaf tips are visible.

- Do not apply within 14 days of harvest.
- Do not apply more than 10 times per season.

### STONE FRUITS

For all stone fruits, Promis Multi-Purpose Fungicide may be applied up to the day of harvest.

### APRICOT AND NECTARINE

controls brown rot blossom blight, brown rot, powdery mildew, and shothole

Mix 1/3 fl oz of Promis Multi-Purpose Fungicide with 1 gallon of water and uniformly apply to all parts of the tree to point of runoff. Apply at early red bud stage (the tips of the petals are visible and the flowers are still closed) before infection occurs. Reapply every 7 to 14 days so long as the disease symptoms persist.

- Do not apply more than 7 times per season.

### CHERRY

Controls brown rot blossom blight, brown rot, leaf spot, and powdery mildew

Mix 1/3 fl oz of Promis Multi-Purpose Fungicide with 1 gallon of water and uniformly apply to all parts of the tree to point of runoff. Apply when single flower buds are visible (but still closed), on short stalks and before infection occurs. Reapply every 7 to 14 days so long as disease symptoms persist. For leaf spot, make an additional application of Promis Multi-Purpose Fungicide after harvest.

- Do not apply more than 7 times per season.

## PEACH

controls brown rot blossom blight, brown rot, powdery mildew, and rust

Mix 1/3 fl oz of Promis Multi-Purpose Fungicide with 1 gallon of water and uniformly apply to all parts of the tree to point of runoff. For brown rot blossom, brown rot, and powdery mildew, apply at early pink bud stage (the tips of tile petals are visible and the flowers are still closed) before infection occurs. Reapply every 7 to 14 days so long as disease symptoms persist. For rust, apply Promis Multi-Purpose Fungicide about 8 weeks after flowering if conditions favoring disease development (hot, moist weather) exist. Reapply every 21 days so long as disease symptoms persist.

- Do not apply more than 7 times per season.

## PLUM AND PRUNE

controls brown rot blossom blight, brown rot, and rust

Mix 1/3 fl oz of Promis Multi-Purpose Fungicide with 1 gallon of water and uniformly apply to all parts of the tree to point of runoff. For brown rot blossom blight and brown rot, apply when the first green leaf tips are visible and before infection occurs. Reapply every 7 to 14 days so long as disease symptoms persist. For rust, apply Promis Multi-Purpose Fungicide about 8 weeks after flowering if conditions favoring

disease development (hot, moist weather) exist. Reapply every 21 days so long as disease symptoms persist.

- Do not apply more than 7 times per season.

## STRAWBERRY

controls leaf blight, leaf spot, and powdery mildew

Mix 1 fl oz of Promis Multi-Purpose Fungicide with 1 gallon of water and uniformly apply to all parts of the plant to point of runoff. Promis Multi-Purpose Fungicide may be applied up to the day of harvest. Apply when the disease first appears or when conditions favoring disease development (hot, moist weather) exist. Reapply every 14 to 21 days so long as disease symptoms persist.

- Do not apply more than 6 times per season.
- Do not plant a new crop in the same place as the crop that was treated within 30 days of the last application.

## BERRIES

For all berries, Promis Multi-Purpose Fungicide may be applied up to the day of harvest.

## BLACKBERRY\*, RASPBERRY\*, LOGANBERRY

controls cane and leaf rust, leaf spot, orange rust, powdery mildew, and yellow rust

Mix 1/2 fl oz of Promis Multi-Purpose Fungicide with 1 gallon of water and uniformly apply to all parts of the plant to point of runoff. Apply at budbreak and reapply every 10 to 14 days so long as disease symptoms persist.

- Do not apply more than 4 times per season.

\*Use on the following members of the caneberry group

including varieties and/or hybrids: Bingleberry, black satin berry, boysenberry, cherokee blackberry, Chesterberry, Cheyenne blackberry, coryberry, Darrowberry, dewberry, Oregon evergreen berry, phenomenal berry, rangeberry, ravenberry, rossberry, Shawnee blackberry, youngberry, Himalayaberry, Dirkson Thornless, hullberry, Lavacaberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, nectarberry

## CURRENT

controls powdery mildew

Mix 1 fl oz of Promis Multi-Purpose Fungicide with 1 gallon of water and uniformly apply to all parts of the plant to point of runoff. Apply before bloom, through bloom and up to 2 weeks post-bloom.

- Do not apply more than 8 times per season.

## GOOSEBERRY

controls anthracnose and powdery mildew

Mix 1 fl oz of Promis Multi-Purpose Fungicide with 1 gallon of water and uniformly apply to all parts of the plant to point of runoff. For anthracnose, apply when the first leaf has completely unfolded. If conditions favoring disease development (hot, moist weather) exist, reapply every 10 to 14 days. For powdery mildew, apply Promis Multi-Purpose Fungicide before bloom, through bloom, and up to 2 weeks post-bloom.

- Do not apply more than 8 times per season.

## GRAPES

controls anthracnose, black rot and powdery mildew

Mix 1 fl oz of Promis Multi-Purpose Fungicide with 1 gallon of

water and uniformly apply to all parts of the plant to point of runoff. For anthracnose and black rot, apply when new shoots are 1 to 3 inches in length. Reapply every 14 days so long as disease symptoms persist. For powdery mildew, apply Promis Multi-Purpose Fungicide when shoots are 12 to 18 inches in length (before bloom). Reapply every 21 days so long as disease symptoms persist.

- Do not apply within 14 days of harvest.
- Do not apply more than 5 times per season.

## PEPPERMINT AND SPEARMINT

controls powdery mildew and rust

Mix 1 fl oz of Promis Multi-Purpose Fungicide with 1 gallon of water and uniformly apply to all parts of the plant to point of runoff. Apply in the early spring when the plants have broken dormancy. Reapply every 14 to 21 days so long as disease symptoms persist.

- Do not apply within 30 days of harvest.
- Do not apply more than 3 times per season.

## VEGETABLES

### ASPARAGUS

controls rust

Mix 1 fl oz of Promis Multi-Purpose Fungicide with 1 gallon of water and uniformly apply to all parts of the plant to point of runoff. Do not apply to harvestable spears. In all locations, apply to the developing ferns after harvest. Reapply every 14 days so long as disease symptoms persist.

In all states except California:

- Promis Multi-Purpose Fungicide may be applied within 180 days of the harvest.

- Do not apply more than one time per growing season.
- In California only:
- Promis Multi-Purpose Fungicide may be applied within 30 days of the harvest.
  - Do not apply more than four times per year.

## CUCURBITS

controls powdery mildew

Includes use on the following members of the cucurbits group:

acorn squash, balsam apple, balsam pear, bitter melon, butternut squash, calabaza, cantaloupe, casaba, Chinese cucumber, Chinese waxgourd, citron melon, creshaw melon, crookneck squash, cucumber, edible gourd, gherkin, golden pershaw melon, honey balls, honeydew melon, hubbard squash, mango melon, muskmelon, ornamental gourd, Persian melon, pineapple melon, pumpkin, Santa Claus melon, scallop squash, snake melon, spaghetti squash, straightneck squash, summer squash, vegetable marrow, watermelon, winter squash, zucchini

Mix 1 fl oz of Promis Multi-Purpose Fungicide with 1 gallon of water and uniformly apply to all parts of the plant to point of runoff. Apply Promis Multi-Purpose Fungicide when the disease first appears. Reapply every 7 to 10 days so long as disease symptoms persist.

- Promis Multi-Purpose Fungicide may be applied up to and including the day of harvest.
- Do not apply more than five times per season.
- Do not plant a new crop in the same place as the crop that was treated within thirty days of the last application.

## SNAP BEAN

controls pod tip rot and rust

Mix 1 fl oz of Promis Multi-Purpose Fungicide with 1 gallon of water and uniformly apply to all parts of the plant to point of runoff. For rust, apply when the disease first appears. For pod tip rot, apply when pods begin to develop. For both diseases, if conditions favoring disease development (hot, moist weather) exist, reapply every 7 to 10 days.

- Promis Multi-Purpose Fungicide may be applied up to and including the day of harvest.
- Do not apply more than 4 times per season.
- Do not plant a new crop in the same place as the crop that was treated within 30 days of the last application.

## TOMATO

controls powdery mildew

Mix 3/4 fl oz of Promis Multi-Purpose Fungicide with 1 gallon of water and uniformly apply to all parts of the plant to point of runoff. Apply when the disease first appears or if conditions favoring disease development (hot, moist weather) exist. Reapply every 21 days so long as disease symptoms persist.

- Promis Multi-Purpose Fungicide may be applied up to and including the day of harvest.
- Do not apply more than 4 times per season.
- Do not plant a new crop in the same place as the crop that was treated within 30 days of the last application.

Notice: Buyer assumes all responsibility for safety and use not in accordance with directions.

## ROSES, FLOWERS, ORNAMENTAL SHRUBS & TREES

Promis Multi-Purpose Fungicide prevents and controls Black spot, Blight (blossom, cercospora leaf, ovulinia petal), Leaf spot (including cercospora), Powdery mildew, Rust (including needle, fusiform), Scab (including fusicladium) and other diseases on a wide variety of roses, flowers, trees and ornamental shrubs. See table for a complete list of plants.

## HOW TO TREAT

Use 3/4 fl. oz. / gallon of water. Apply to the point when spray just begins to runoff the leaves. Apply once every 2 weeks throughout the season or until conditions favoring disease development (hot, moist weather) are no longer present. Allow spray to dry before watering.

### USE TIPS

1. Treat roses on a 7 to 10 day schedule. If black spot is not a problem, a 10 to 14 day schedule is recommended.
2. When treating azaleas, mountain laurel or rhododendrons, begin spraying when flowers start to show color.
3. When treating irises, increase use rate to 1 1/2 fl.oz. / gallon of water.
4. For Douglas fir, loblolly pine and slash pine use 2 1/4fl. oz. / gallon of water. Begin treating in early spring and continue treating every 2 to 3 weeks until threat of infection has passed.

PLANT (Number following plant refers to the specific Use Tip above.)	DISEASES
Rose(1)	Black spot, Powdery mildew, Rust
Azalea (2), Mountain laurel (2), Rhododendron (2)	Petal blight ( <i>ovulinia</i> spp.) Powdery mildew
Abelia, Acalypha (Copper-leaf), Catalpa, Flowering cherry	Cercospora leaf spot
Hackberry, Honeysuckle, Privet, Trumpet creeper, Zinnia	Powdery mildew
Ageratum, Anemone, Angelica, Ash, Aster, Birch, China aster, Chokeberry, Chrysanthemum, Columbine, Cornflower, Dianthus, Flowering quince, Flowering maple (Abutilon), Four o'clock, Fuchsia, Iris (3), Marigold, Moonflower, Ninebark, Russian olive, Smoke-tree (Gothus)	Rust
Achillea (Yarrow), Alder, Amelanchier (Juneberry, Shadbush), Barberry, Carnation, Delphinium, Gaillardia, Gardenia, Seranium, Hollyhock, Mock-orange, Pansy, Petunia, Photinia, Salvia, Snap-dragon, Viburnum	Powdery mildew, Rust
African Violet, Begonia, Bittersweet, Buckeye, California poppy, Chestnut (Horse), Cosmos, Crepe myrtle, Dahlia, Elm, Euonymus, Gourd (Ornamental), Grape leaf ivy, Lilac, Locust, Maple, Oak, Poinsettia, Spirea, Sycamore, Walnut, Willow	Powdery mildew

PLANT (Number following plant refers to the specific Use Tip above.)	DISEASES
Flowering almond	Blossom blight (Monilinia spp.)
Amorpha (False indigo), Bellflower, Buttonbush, Fleabane, Phlox, Sunflower	Cercospora leaf spot, Powdery mildew, Rust
Calendula, Hydrangea, Leucothoe	Cercospora leaf spot
Flowering crabapple, Flowering pear, Hawthorn	Powdery mildew, Rust, Scab
Dogwood	Anthracnose, Powdery mildew
Douglas Fir (4)	Needle rust
Loblolly pine, Siasirpine (4)	Fusiform rust
Pyracantha	Fusicladium scab

### STORAGE AND DISPOSAL

**PESTICE STORAGE:** Store in a cool, dry area away from heat and open flame. **PESTICIDE DISPOSAL:** If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain. **CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available.

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear protective eyewear. Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

### USER SAFETY RECOMMENDATIONS

1. User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
2. User should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area.

Manufactured For:  
Oregon Global Distribution, Inc.  
1600 Skypark Dr, Suite 212  
Medford, OR 97504  
1-800-994-4508

031715v7