32802-67-90098_Pennington Ultragreen Crabgrass Plus Fert II 30-0-4_20160830_10_90098_ .pdf

Top Seal Customer 15mm



ENHANCED EFFICIENCY FERTILIZER
Feed More Lawn with Less Fertilizer

PENNINGTON®

ultraGreen

CRABGRASS PREVENTER

PLUS FERTILIZER II

30-0-4

PREVENTS CRABGRASS FOR UP TO 5 MONTHS

PREVENT CRABGRASS AND FERTILIZE YOUR LAWN IN ONE APPLICATION

Boosts Nutrient Uptake for a Lush, Green Lawn

Increases Drought Tolerance & Reduces Watering Needs by Extending Root Absorption Area

COVERS UP TO 5,000 SQ. FT.

Net Weight 12.5 LB | Peso Neto 5.7 kg

ACTIVE INGREDIENT:
Prodiamine (CAS 29091-21-2) 0.58%
OTHER INGREDIENTS: 99.42%
TOTAL: 100.00%

Read the entire "Directions for Use" and "Warranty Statements" before using this product. If terms are not acceptable, return the unopened product container at once to the seller.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for further Precautions and First Aid information.

MANTÉNGALO LEJOS DEL ALCANCE DE LOS NIÑOS

PRECAUCIÓN

Consulte la parte de atras el para conocer los avisos de precauciones e información de primeros auxilios.

Side Seal 8mm

CUT HERE TO OPEN

The Pennington® UltraGreen® Difference

Pennington® UltraGreen® products are more efficient than other fertilizers. Half of the nitrogen in this premium line*** is stabilized**, which allows grass to use more of the nutrients you put down and reduces nutrient loss to the environment. The other half of the nitrogen is slow-release for extended feeding for up to 3 months.

Additionally, these unique blends contain mycorrhizal fungi, natural organisms that have been present in vigorous soils for millions of years. These microscopic fungi increase the surface absorbing area of roots, greatly improving the ability of the lawn to access vital soil resources such as water and nutrients. These tiny fertilizer granules are packed with essential plant nutrients and fungi, so you can feed more lawn with less fertilizer.

***Excluding the Pennington® UltraGreen® Starter Fertilizer.

Benefits of Pennington® UltraGreen® Crabgrass Preventer Plus Fertilizer+ 30-0-4

- Prevent crabgrass and fertilize your lawn in one application
- Up to 5 months of weed grass control
- Quick greening and extended feeding
- Phosphorus Free- Helping Reduce Phosphorus in Water Sources

Pennington® UltraGreen® Crabgrass Preventer Plus Fertilizer+ 30-0-4 WILL NOT BURN when used as directed. We're so confident you'll love the results of using this product that it comes with the Pennington Satisfaction Guarantee.*

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS: To protect the environment, do not allow this product to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather will help to ensure that weed does not blow this product off the treated area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

FIRST AID

If in eyes	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed	• Call a poison control center or physician immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or physician. • Do not give anything by mouth to an unconscious person.
If inhaled	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

Hot Line Number

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call 1-800-265-0761.

USE PRECAUTIONS: The following precautions apply to the use of this product in residential lawns. Application of this product may thin emerged annual ryegrass and newly overseeded grasses. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil). Do not apply to newly sod sod until the following year. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control. Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn and sweep any product that lands on the driveway, sidewalk or street back onto your lawn.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Not for sale or use but being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in established resident turf lawns. It controls susceptible weeds by inhibiting weed seed germination and root development. When used according to directions, it will control: barnyardgrass, bluegrass, annual (Poa annua), carpetweed; chickweed, common, chickweed, mouseear (from seed), crabgrass (large, smooth), crowfootgrass, crupgrass, woolly, foxtails, annual; goosegrass; henbit; lichgrass; johnsongrass (from seed); junglerice, knotweed, kochia, lambsquarter, common, liverpass; panicum (Texas, fall, browntop); pigweed; purslane, common, puskly (Florida, rosacegrass); shepherd's purse; signalgrass, broadleaf, speedwell, Persian, sprangletop; spurge, prostrate, stichgrass; woodsorrel, yellow (from seed).

[1] In many areas where not prohibited by fertilizer use restrictions[2] a single application of 2.5 to 6 pounds/1000 square feet (112 to 255 pounds/acre) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 2.5 to 3.9 pounds/1000 square feet (112 to 173 pounds/acre) to areas not prohibited by fertilizer use restrictions, followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Minimum and Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label).
[2] For this weed, apply in late summer, fall or winter prior to germination.
[3] Suppression only.
[4] Consult your State Cooperative Extension Service for current information on annual application limitations for lawn fertilizers in your region.

WHEN TO APPLY TO ESTABLISHED TURF: This product must be applied prior to germination of weed seed, as it will not control established weeds.

APPLICATION DIRECTIONS: Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 24 days after application.

RATES OF APPLICATION: This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed.

MINIMUM AND MAXIMUM APPLICATION RATES (by Turf Grass Species)[1]

Turf Species	Minimum Rate	Maximum Rate
Bahiagrass[2], Centipedegrass[2], Bermudagrass[2], Zoysiagrass, Seashore Paspalum, St. Augustinegrass, Tall Fescue (including turf-type)	4 pounds per 1000 sq. ft. (175 lbs/acre)	6 pounds/1000 sq. ft. per year (259 lbs/acre per year)
Buffalograss, Kentucky Bluegrass, Perennial Ryegrass	2 pounds per 1000 sq. ft. (87.5 lbs/acre)	4 pounds/1000 sq. ft. per year (173 lbs/acre per year)
Creeping Red Fescue	1.5 pounds per 1000 sq. ft. (65.3 lbs/acre)	2.9 pounds/1000 sq. ft. per year (130 lbs/acre per year)
Creeping Bentgrass	1.5 pounds per 1000 sq. ft. (65.3 lbs/acre)	2.5 pounds/1000 sq. ft. per year (112 lbs/acre per year)

[1] These are maximum rates per calendar year.
[2] In Florida, Do not exceed 2 applications per year to Bahiagrass and Centipedegrass in the north. Do not exceed 3 applications per year to Centipedegrass in the central and south regions. Consult your State Cooperative Extension Service for current information on annual application limitations for lawn fertilizers in your region.
[3] May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 2 lbs/1000 square feet (87.5 pounds/acre). Newly sprigged or plugged Bermudagrass stolons rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and nutsedge due to reduced rates used in sprigging situations.

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

BROADCAST SPREADERS: For best results, apply in a circular path starting on the outside and working toward the center overlapping slightly to ensure complete coverage.
DROP SPREADERS: Apply 2 strips along each end. Then apply along length of lawn, overlapping slightly and shutting off the spreader before each turn at edge of border strips.

SPREADER SETTINGS / AJUSTE DEL ESPARCIDOR	Setting / Ajuste (2.5 pounds per 1000 sq. ft.)
Scotts® Drop Spreaders	2 ¾
Scotts® Broadcast Spreaders	3 ½
Sta-Green® Broadcast Spreader	3 ¼
Pennington® Broadcast Spreader	3

SETTING YOUR LAWN SPREADER

THESE SETTINGS ARE A GUIDE. They were determined by landscape professionals using new spreaders. The condition of your spreader, your walking speed, the roughness of the lawn, and even the humidity can affect how fast the lawn food comes out. Watch flow rate. If fertilizer appears to be coming out either slower or faster than you are covering your lawn, adjust the setting. Close the spreader when lifting, turning or stopping.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.
PESTICIDE STORAGE: Store in original container in a cool, dry, locked storage area out of reach of children and domestic animals.
PESTICIDE DISPOSAL: IF EMPTY: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available.
IF PARTLY FILLED: Call your local solid waste agency for disposal instructions. Never place any unused product down any indoor or outdoor drain or toilet.

WARRANTY

The directions for use of this product are believed to be adequate and must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Plant injury, ineffectiveness, or other unintended consequences may result due to such factors as weather conditions, presence or absence of other materials, or the manner of use or application, all of which are beyond the control of Central Garden & Pet, Garden Division, the manufacturer, or the seller. To the extent consistent with applicable law, the products sold to you are furnished "as is" by Central Garden & Pet, Garden Division. The manufacturer and seller are not liable to the manufacturer's warranties, if any, which appear on the label of the product sold to you. Except as warranted by this label Central Garden & Pet, Garden Division, the manufacturer, or the seller makes no warranties, guarantees, or representations of any kind to the buyer or the user, either express or implied, or by usage of trade, statutory or otherwise, with regard to the product sold or use of the product including, but not limited to, merchantability, fitness for a particular purpose or use, or eligibility of the product for any particular trade usage. To the extent consistent with applicable law, Buyer's or user's exclusive remedy, and Central Garden & Pet, Garden Division, the manufacturer's or the seller's total liability shall be limited to damages not exceeding the cost of the product. No agent or employee of Central Garden & Pet, Garden Division or the seller is authorized to amend the terms of this warranty disclaimer or the product's label or to make a presentation or recommendation different from or inconsistent with the label of this product.
To the extent consistent with applicable law, Central Garden & Pet, Garden Division, the manufacturer, or the seller shall not be liable for consequential, special, or indirect damages resulting from the use, handling, application, storage, or disposal of this product or for damages in the nature of penalties, and the buyer and the user waive any right that they may have to such damages.
***Pennington Satisfaction Guarantee.** If, after following the instructions, you are not satisfied with the product results, send your name, address, and proof of purchase (original dated receipt) to: **Pennington Refund, 1000 Parkwood Circle, Suite 700, Atlanta, GA 30339**

30-0-4

GUARANTEED ANALYSIS
Total Nitrogen (N) 30%
30% Urea Nitrogen**
Soluble Potash (K2O) 4%
Calcium (Ca) 1%
Derived from Urea, Polymer Coated Urea, Muriate of Potash, and Limestone

**Contains 15% Slowly Available urea nitrogen from polymer coated urea.
***Contains 15% nitrogen stabilized with N-(n-butyl) thiophosphoryl triamide
**Stabilized nitrogen minimizes nitrogen losses to the environment and enhances its efficiency.

F002063

ALSO CONTAINS NON PLANT FOOD INGREDIENTS
SOIL AMENDING GUARANTEED ANALYSIS
Active Ingredients:
Mycorrhizal Fungi: ..... 1.12 pieces/gm
Glomus intraradices ..... 0.28 pieces/gm
Glomus mosseae ..... 0.28 pieces/gm
Glomus etunicatum ..... 0.28 pieces/gm
Glomus aggregatum ..... 0.28 pieces/gm
Inert Ingredients (as a soil amendment):
99.03% Fertilizer ingredients
0.58% Herbicide ingredients

F002063

Apply only to actively growing turf. Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn. Check with your local Cooperative Extension Agent to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm

Information concerning the raw materials composing this product can be obtained by writing to the Guarantor listed on this label and referring to the batch number found on this container.

NOTICE - This product contains chemicals known to the State of California to cause cancer and birth defects or other reproductive harm.

PRECAUCIONES

PELIGROS PARA PERSONAS Y ANIMALES DOMÉSTICOS

PRECAUCIÓN: Causa lesiones (irritación) moderada de los ojos. Dañino si se ingiere o absorbe a través de la piel. Evite el contacto con la piel, los ojos o la ropa. Evite respirar el polvo. Lávese con agua y jabón después de manipularlo. El contacto prolongado o repetido de este producto con la piel puede ocasionar reacciones alérgicas en algunas personas.
PELIGROS AMBIENTALES: Para proteger el ambiente, no permita que este producto penetre o sea arrastrado hacia desagües para tormentas, zanjas de drenaje, cunetas o aguas de superficies. Aplique este producto en tiempo de calma para evitar que el viento no lo desplace del área tratada. Barra hacia el área tratada del jardín o césped cualquier residuo de producto que caiga en la entrada del garaje, la acera o la calle para evitar que sea arrastrado hacia los cuerpos de agua o sistemas de drenaje.

Primeros auxilios

En caso de contacto con los ojos	• Mantenga el ojo abierto y enjuague lenta y suavemente con agua durante 15-20 minutos. • Si usa lentes de contacto, retírelos después de los primeros 5 minutos y continúe enjuagando posteriormente. • Llame a un centro de control de envenenamiento o un médico para recibir consejos sobre el tratamiento.
En caso de contacto con la piel o la ropa	• Quítese la ropa contaminada. • Enjuague la piel inmediatamente con agua abundante durante 15-20 minutos. • Llame a un centro de control de envenenamiento o un médico para recibir consejos sobre el tratamiento.
En caso de ingestión	• Llame a un centro de control de envenenamiento o un médico para recibir consejos sobre el tratamiento. • Haga que la persona beba un vaso de agua si es capaz de tragar. • No induzca el vómito a menos que se lo indique el médico o el centro de control de envenenamiento. • No le de nada por la boca a una persona en estado inconsciente.
En caso de inhalación	• Traslade a la persona al aire fresco. • Si la persona no puede respirar, llame al 911 o a un servicio de ambulancia y luego dele respiración artificial, preferiblemente boca a boca si es posible. • Llame a un centro de control de envenenamiento o un médico para recibir consejos adicionales sobre el tratamiento.

Teléfono de asistencia de emergencia Tenga a mano el recipiente o la etiqueta del producto cuando llame al centro de control de envenenamiento o el médico, o cuando acuda para recibir tratamiento. Para recibir más información sobre este pesticida (incluyendo preocupaciones sobre la salud, emergencias médicas o incidentes con el pesticida) llame al 1-800-265-0761.

PRECAUCIONES DE USO: Las siguientes precauciones se aplican al uso de este producto en céspedes residenciales. La aplicación de este producto puede afectar la densidad de los céspedes de poa anual recién sembrados y los céspedes recién resembrados. No lo aplique en lapes resembrados durante los 60 días posteriores a la siembra de semillas o hasta después de la segunda poda, lo que suceda después. Pueden producirse daños a las plantas deseables si se aplica el producto antes de que las raíces secundarias de las plantas alcancen la segunda pulgada de profundidad del suelo (no de paja con suelo). No lo aplique al lape recién colocado hasta el año siguiente. La aplicación a un lape afectado por la sequía, la poca fertilidad o la acción de las plagas puede ocasionar daños al lape. La alteración de la barrera de herbicida con prácticas de cultivo (como la aireación) puede conducir a un control reducido de la maleza. No lo aplique cerca del agua, desagües para tormentas o zanjas de drenaje ni tampoco si se espera lluvia fuerte. Aplique solo el césped y barra hacia el césped cualquier producto que caiga en las aceras, entradas de carros o calles.

INDICACIONES DE USO

El uso de este producto de manera diferente a la indicada en su etiqueta es una violación de las leyes federales.
Este producto no está comercializado para usar en céspedes cultivados para la venta o cualquier uso comercial como es el caso de lapes, o para la producción comercial de semillas, ni para fines investigativos.
Este producto es un herbicida selectivo de preemergencia que brinda un control residual de muchos variedades de hierbas y malezas de hojas anchas en céspedes. El mismo controla las malezas susceptibles al inhibir la germinación de la semilla y el desarrollo de las raíces de la maleza. Cuando se usa según las indicaciones, el producto controla: equinochloa o pata de gallo, pata (Poa annua), verdolaga afelonza, altiere, verdolaga (de semillas), digitaria (grande, lisa), hierba pata de gallo, woolly crupgrass, cola de zorro anual, grama de caballo; ortiga menor, lichgrass, sarga de abrojo (a partir de semillas), pasto del arroz, centinodia, kochia, cenizo común, zacate llanero, paniceo (Texas, rapidente, browntop); maleza colorada, verdolaga común, trébol de Florida, cenadillo; berza de pastor; bracaná de hojas anchas, verónica persa, zacate gigante, galinchina, ortiga menor, ace amarillo (a partir de semilla).
En muchas regiones donde no está prohibido por las restricciones de uso de fertilizantes[1], una sola aplicación de 2.5 a 6 libras/1000 pies cuadrados (de 112 a 255 libras/acre) es suficiente para controlar la grama de caballo. Sin embargo, en condiciones de mucha presión de hierba grama de caballo y/o una temporada de crecimiento extendida, se puede obtener un control más efectivo con una aplicación inicial de 2.5 a 3.9 libras/1000 pies cuadrados (de 112 a 173 libras/acre) en las regiones donde no le prohiben las restricciones del uso de fertilizantes, seguida de una segunda aplicación en dosis que no exceda la tasa anual máxima (vea la tabla "Dosis de aplicación mínima y máxima según las especies de hierbas" en la sección "Dosis de aplicación" de esta etiqueta).
[1] Para esta maleza, aplíquelo a fines de verano, en otoño o invierno antes de la germinación.
[2] Supresión solamente.
[3] Consulte a su Servicio de Extensión Cooperativa de Estado para obtener información actualizada sobre las limitaciones de aplicación anuales de fertilizantes del césped en su región.

CUÁNDO APLICARLO AL CÉSPED ASENTADO: Este producto se debe aplicar antes de que germinen las semillas de la maleza, ya que no sirve para controlar la maleza asentada.
INDICACIONES DE USO: Aplique este producto uniformemente, usando un esparcidor adecuado que esté calibrado correctamente. El control más efectivo de la maleza se obtiene cuando este producto es activado por una capa mínima de 0.5 pulgada de lluvia o irrigación, o con la incorporación poco profunda (1-2 pulgadas) antes de que germinen las semillas de la maleza y dentro de las 24 días posteriores a la aplicación.
DOSIS DE APLICACIÓN: Se recomienda usar este producto en las especies de hierba mencionadas en la tabla siguiente. La cantidad máxima del producto por año varía según la especie de hierba. No exceda la tasa de aplicación anual que se indica.

DOSIS DE APLICACIÓN MÁXIMOS Y MÍNIMOS (según la especie de hierba)[1]

Especie de hierba	Dosis mínima	Dosis máxima
Pasto bahía[2], Gramparis[2], hierba Bermuda[2], grama japonesa zoysia, chepica blanca, hierba de San Agustín, festuca alta (incluyendo las de lepa)	4 libras por 1000 pies cuadrados (175 lbs/acre)	6 libras/1000 pies cuadrados por año (259 lbs/acre por año)
Zacate búfalo, poa común, césped perenne	2 libras por 1000 pies cuadrados (87.5 lbs/acre)	4 libras/1000 pies cuadrados por año (173 lbs/acre por año)
Festuca roja	1.5 libras por 1000 pies cuadrados (65.3 lbs/acre)	2.9 libras/1000 pies cuadrados por año (130 lbs/acre por año)
Nombre de dios	1.5 libras por 1000 pies cuadrados (65.3 lbs/acre)	2.5 libras/1000 pies cuadrados por año (112 lbs/acre por año)

[1] Estas son las dosis máximas por año-calendario.
[2] En Florida: No exceda las 2 aplicaciones por año para hierba bahía y gramparis en la región norte. No exceda las 3 aplicaciones por año para hierba gramparis en las regiones central y sur. Consulte en el Servicio de Extensión Cooperativa Estatal para conocer la información actual sobre las limitaciones anuales en cuanto a la aplicación de fertilizantes de césped en su región.
[3] Se puede usar en hierba Bermuda recién sembrada por ramas o postures o una vez que ha crecido las 2 lbs/1000 pies cuadrados (87.5 libras/acre). La formación de estolones en la hierba Bermuda recién sembrada por ramas o postures puede retardarse temporalmente. La supresión solo de la cola de zorro, grama de caballo y cocalote debido a las dosis reducidas que se usan en situaciones de siembra de ramas.

Este producto se puede aplicar como una aplicación simple o aplicaciones secuenciales para controlar la germinación de las malezas durante todo el año. Se pueden realizar aplicaciones secuenciales siempre y cuando la cantidad del producto aplicado no exceda la tasa de aplicación anual máxima recomendada para cada variedad de hierba. El período de tiempo del control residual de maleza que ofrece este producto está relacionado con la tasa aplicada. Para obtener un mayor período de control de la maleza, use las dosis superiores recomendadas.

ESPARCIDORES POR DISPERSIÓN. Para lograr mejores resultados, aplíquelo en una trayectoria circular comenzando por la parte exterior y trabajando hacia el centro, solapando ligeramente sobre el área recién aplicada para garantizar una cobertura completa.
ESPARCIDORES POR GOTEO. Aplíquelo en 2 franjas a lo largo de cada extremo y aplíquelo luego a lo largo del césped, solapando ligeramente sobre el área recién aplicada y apagando el esparcidor antes de cada giro al final de cada trayectoria por los bordes.

ALMACENAJE Y DESECHO

No contamine el agua, la comida o el alimento al almacenarlo o desecharlo.
ALMACENAJE DEL PESTICIDA: Almacénelo en el recipiente original en una área de almacenaje fresca y seca tapa-llave que se encuentre lejos del alcance de los niños y las mascotas domésticas. **PARA DESECHAR EL PESTICIDA: SI ESTÁ VACÍO.** Este es un recipiente no rellenable. No rellene o reuse este recipiente. Recíclelo, si es posible. **SI ESTÁ PARCIALMENTE LLENO:** llame a la agencia de desechos sólidos de su localidad para recibir instrucciones sobre cómo desecharlo. Nunca deseche un producto que no se vaya a usar en un desagüe o vertedero para exteriores o interiores.

AJUSTE DEL ESPARCIDOR PARA CÉSPED

ESTOS AJUSTES SON SOLO UNA GUÍA. Y han sido determinados por jardineros profesionales usando esparcidores nuevos. Las condiciones de su esparcidor, la velocidad con la que usted camina, la aspereza de su césped e incluso la humedad pueden afectar la velocidad con la que sale el nutriente de su esparcidor. Puede alterar el flujo si el fertilizante sale más rápido o más despacio de lo que usted cabe el césped, modifique el ajuste del esparcidor. Cierre siempre el esparcidor cuando lo rellene, cuando gire o cuando se detenga.

GARANTÍA

Las indicaciones para el uso de este producto se consideran adecuadas y se deben seguir minuciosamente. Es imposible eliminar todos los riesgos asociados inherentemente con el uso de este producto. Los daños a plantas, la ineficacia o cualquier otra consecuencia no deseada puede deberse de diversos factores como las condiciones del clima, la presencia o ausencia de otros materiales, o la manera de usarlo o aplicarlo, todos y cada uno de los cuales están más allá del control de Central Garden & Pet, la División de Jardinería, el fabricante o el vendedor. En consonancia con lo estipulado por la ley correspondiente, los productos vendidos se ofrecen tal y como viene el producto por parte de Central Garden & Pet y la División de Jardinería. El fabricante y el vendedor están sujetos solamente a las garantías del fabricante, si existeran, que se mencionan en la etiqueta del producto vendido a usted. Excepto cuando sea garantizado por la referida en la etiqueta, la división de jardinería de Central Garden & Pet, el fabricante o el vendedor no emiten garantías ni representaciones de ninguna índole al vendedor o usuario, ya sean explícitas e implícitas, de uso, reglamentarias o de ninguna índole, con referencia al producto vendido a al uso del producto, incluyendo, pero sin limitarse a, la comercialidad, adecuación para un propósito o uso particular, o elegibilidad del producto para cualquier uso comercial exclusivo. En consonancia con lo estipulado por la ley correspondiente, la única solución para el comprador o usuario, y la única responsabilidad total de Central Garden & Pet, la División de Jardinería, el fabricante o el vendedor quedará limitada a los daños que no excedan el costo del producto. Ningún agente o empleado de Central Garden & Pet, la División de Jardinería o el vendedor está autorizado a enmendar los términos de esta declaración de garantía o de la etiqueta del producto, o a efectuar presentaciones o recomendaciones diferentes o que difieren con lo mencionado en la etiqueta de este producto.
En consonancia con lo estipulado por la ley correspondiente, Central Garden & Pet, la División de Jardinería, el fabricante o el vendedor no asumen responsabilidad alguna por los daños consecuentes, especiales e indirectos que se deriven del uso, la manipulación, la aplicación, el almacenaje o el desecho de este producto o por los daños relacionados con las penalizaciones, y el comprador y usuario renuncian a cualquier derecho que puedan tener en relación con dichos daños.
***Garantía de satisfacción de Pennington.** Si, después de usarlo siguiendo las instrucciones, usted no está satisfecho con los resultados, envíe su nombre, dirección y prueba de la compra (recibo original con fecha) a: **Pennington Refund, 1000 Parkwood Circle, Suite 700, Atlanta, GA 30339**

30-0-4

ANÁLISIS GARANTIZADO
Total de nitrógeno (N) 30%
30% de nitrógeno ureico**
Potasa soluble (K2O) 4%
Calcio (Ca) 1%
Derivado de la urea, urea con recubrimiento de polímero, muriato de potasa y cal.

**Contiene un 15% de nitrógeno ureico de liberación lenta a partir de la urea con recubrimiento de polímero.
***Contiene un 15% de nitrógeno estabilizado con triamida N-(n-butil) tiofosfórica.
** El nitrógeno estabilizado minimiza la pérdida de nitrógeno hacia el ambiente y mejora su eficiencia.

CONTIENE TAMBIÉN INGREDIENTES QUE NO SON NUTRIENTES DE PLANTAS
ANÁLISIS GARANTIZADO PARA ENMENDAR EL SUELO
Ingredientes activos:
Hongos micorrizos ..... 1.12 piezas/gm
Glomus intraradices ..... 0.28 piezas/gm
Glomus mosseae ..... 0.28 piezas/gm
Glomus etunicatum ..... 0.28 piezas/gm
Glomus aggregatum ..... 0.28 piezas/gm
Ingredientes inertes (como acondicionadores del suelo):
99.03% de ingredientes fertilizantes
0.58% de ingredientes herbicidas

F002063

Aplíquelo solo a céspedes en crecimiento. No lo aplique en la proximidad de agua, drenajes o zanjas de drenaje. No lo aplique si se espera mucha lluvia. Aplique este producto solamente a huertas o jardines y barra los residuos que caigan en el área de porgueo, la acera o la calle hacia el césped. Verifique con la Agencia de Extensión Cooperativa de su comunidad para obtener información específica acerca de las mejores prácticas del cuidado del césped. Verifique con el gobierno de su condado o ciudad para determinar si hay restricciones locales sobre el uso o la aplicación de fertilizantes.
Para ver la información disponible acerca del contenido y los niveles de metales en este producto, visite la página de Internet:
http //www.aapfco.org/metals.htm

La información acerca de las materias primas que componen este producto se puede obtener escribiendo al avalador mencionado en esta etiqueta y refiriendo el número de lote que se encuentra en este recipiente.

AVISO - Este producto contiene sustancias químicas reconocidas en el estado de California como causantes de cáncer y defectos congénitos u otros daños al sistema reproductivo.

Expiration Date

Distributed & Guaranteed by
Central Garden & Pet
Garden Division
1000 Parkwood Circle, Suite 700
Atlanta, GA 30339

gardendivision@central.com
www.penningtonfertilizer.com
1-800-265-0761

EPA Reg. No. 32802-67-90098
EPA Est. No. 59144-CA-1, 35497-OR-1, 32802-MO-1, 37351-CO-1
Subscript used is the First letter of Run Code on Container

© 2014, 2015 Central Garden & Pet Company
300520898 - 8/16 • UG123 v.3

CENTRAL Garden & Pet

Pennington and Pennington with design are registered trademarks of Pennington Seed, Inc. Central Garden & Pet, Central Garden & Pet with design and UltraGreen are trademarks of Central Garden & Pet Company.
Sta-Green is a registered trademark of LF LLC.
Scotts is a registered trademark of OMS Investments, Inc.

0 21496 01329 9