

REF **MSC351430A**

Manufactured for Medline Industries Inc., Three Lakes Drive, Northfield, IL 60093. 1-800-MEDLINE. Made in the USA. RK1919ATH

For product emergency, contact: Poison Control Center, 1-800-222-1222

For Technical Information call: 1-800-MEDLINE

For an SDS, call 1-800-MEDLINE and request document MSC351430A

Exp

100% of the respondents (n = 100) were female, and 100% (n = 100) were white. The majority of the respondents (n = 80) were aged 18–24 years, and 100% (n = 100) were currently attending a postsecondary institution. The majority of the respondents (n = 80) were currently employed, and 100% (n = 100) were currently attending a postsecondary institution.

***SPECIAL INSTRUCTIONS FOR CLEANING PRIOR TO DISINFECTING AGAINST *Candida auris*.

- **PERSONAL PROTECTION:** Wear appropriate barrier protection such as gloves, gowns, masks, or eye covering.
- **CLEANING PROCEDURE:** Fecal matter/waste must be thoroughly cleaned from surfaces/objects before disinfection by application with a clean cloth, mop, and/or sponge saturated with the product. Pre-cleaning is to include vigorous wiping and/or scrubbing and all visible soil is removed. Surfaces in patient rooms are to be cleaned in an appropriate manner, such as from right to left or left to right, on horizontal surfaces, and top to bottom, on vertical surfaces, to minimize spreading the organism. Restrooms are to be cleaned last. Do not reuse soiled cloths.
- **INFECTIOUS MATERIAL DISPOSAL:** Materials used in the cleaning process that may contain feces/wastes are to be disposed of immediately in accordance with local regulations for infectious materials disposal.

*SPECIAL INSTRUCTIONS FOR CLEANING PRIOR TO DISINFECTING AGAINST *Clostridium difficile* SPORES.

- **PERSONAL PROTECTION:** Wear appropriate barrier protection such as gloves, gown, mask, or eye covering.
- **CLEANING PROCEDURE:** Fecal matter /waste must be thoroughly cleaned from surfaces /objects before disinfection by application with Micro-Kill Bleach Wipes. Cleaning is to include vigorous wiping and/or scrubbing, until all visible soil is removed. Special attention is needed for high-touch surfaces. Surfaces in patient rooms are to be cleaned in an appropriate manner, such as from right to left or left to right, on horizontal surfaces, and top to bottom, on vertical surfaces, to minimize spreading of the spores. Restrooms are to be cleaned last. Do not reuse soiled wipes.
- **INFECTIOUS MATERIAL DISPOSAL:** Materials used in the cleaning process that may contain feces /wastes are to be disposed of immediately in accordance with local regulations for infectious materials disposal.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barriers or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

*Follow the Special Instructions for Cleaning prior to disinfection for *Clostridium difficile* spores*.

+Follow the Special Instructions for Cleaning prior to disinfection for bloodborne pathogens.

^A pre-cleaning step is required for efficacy against +HIV, +HBV, +HCV, *Candida auris****, and *Clostridium difficile* spores*.

†This product is for the exterior surfaces of Blood Glucose Meters and is not for use on the needle and/or the monitors.

FIRST AID
If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. Have the product container with you

when calling a poison control center or doctor, or going for treatment.

PRECAUTIONARY STATEMENTS
Hazards to humans and domestic animals.

CAUTION: Causes moderate eye irritation. Avoid contact with eyes and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PHYSICAL AND CHEMICAL HAZARDS
This product contains bleach. Do not use this product with other chemicals such as ammonia, toilet bowl cleaners, rust removers, or acid, as this releases hazardous gases.

STORAGE AND DISPOSAL
Do not contaminate water, food or feed by storage or disposal

Pesticide Storage:
Keep this product in a tightly closed canister, when not in use. Store canister in cool, dry, and well-ventilated area away from heat, open flame, direct sunlight and inaccessible to children.

Pesticide Disposal:
Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling and Disposal:
Nonrefillable rigid canister. Do not re-use or refill this canister. Triple rinse (or equivalent), promptly after emptying. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

REF MSC351440AN



(01) 10888277652538



Manufactured for Medline Industries Inc.,
Three Lakes Drive, Northfield, IL 60093.
Micro-Kill is a trademark of Medline Industries, Inc.
1-800-MEDLINE. Made in the USA. RK19MGW

For product emergency, contact:
Poison Control Center, 1-800-222-1222
For Technical Information call: 1-800-MEDLINE
For an SDS, call 1-800-MEDLINE
and request document MSC351440AN

MEDLINE

MICRO-KILL™ Bleach

GERMICIDAL BLEACH WIPES



***SPORICIDAL** 3 minute contact time
Clostridium difficile spores*

****VIRUCIDAL** 30 second contact time
Adenovirus Type 5
Avian Influenza A Virus (H3N2)
Avian Influenza A Virus (H5N1)
Canine Parvovirus
Bovine Viral Diarrhea Virus (BVDV) (HCV)
Duck Hepatitis B Virus (DHBV) (HBV)
Enterovirus D68
Feline Calicivirus (ATCC VR-782)
Feline Panleukopenia Virus
Hepatitis A Virus
+Hepatitis B Virus
+Hepatitis C Virus
Herpes Simplex Virus Type 2
Human Coronavirus
+Human Immunodeficiency Virus Type 1
Influenza A Virus (Novel H1N1)
Influenza B Virus (Strain B/Hong Kong/5/72)
Norovirus (Feline Calicivirus)
Polio Virus Type 1
Respiratory Syncytial Virus
Rhinovirus Type 37
Rotavirus

FUNGICIDAL
1 minute contact time
Candida albicans
Trichophyton interdigitale
2 minute contact time
*Candida auris****

BACTERICIDAL 30 second contact time
Acinetobacter baumannii-MDR
Bordetella pertussis
Campylobacter jejuni
CA-Methicillin Resistant *Staphylococcus aureus*
Enterobacter aerogenes
Enterobacter cloacae
Enterococcus faecalis-VRE
Enterococcus faecium-MDR
Escherichia coli
Escherichia coli O157:H7
Escherichia coli-ESBL
Escherichia coli-NDM-1



Klebsiella pneumoniae
Klebsiella pneumoniae-CRKP (CRE)
Klebsiella pneumoniae-ESBL
Klebsiella pneumoniae-NDM-1
Legionella pneumophila
Linezolid Resistant *Staphylococcus aureus*
Listeria monocytogenes
Methicillin Resistant *Staphylococcus aureus* (MRSA)
Proteus mirabilis
Pseudomonas aeruginosa
Salmonella enterica
Serratia marcescens
Shigella dysenteriae
Staphylococcus aureus
Staphylococcus aureus-VISA
Staphylococcus aureus-VISA
Streptococcus pneumoniae
Streptococcus pyogenes

Klebsiella pneumoniae is a member of the family *Enterobacteriaceae*.

These wipes are sporicidal*, virucidal**, bactericidal and fungicidal*** and can be used to clean, deodorize and disinfect hard, non-porous healthcare and environmental surfaces, patient care equipment and point-of-care equipment in one-step^.

Equivalent to 1:10 dilution of 6.5% bleach

Active Ingredient: Sodium hypochlorite 0.65%
Other Ingredients~: 99.35%
Total 100.00%

~Other ingredients do not include the weight of wipes.

KEEP OUT OF REACH OF CHILDREN CAUTION

SUGGESTED AREAS OF USAGE:

Medical, Dental and Laboratory Counters, Exam Tables, Carts, Patient Care Equipment, Point-of-Care Equipment, Telephones, Sink Tops, Toilet Seats, Digital Screens, on Plastics, Glazed Porcelain, Stainless Steel, Formica, Vinyl Surfaces and Blood Glucose Meters†.

See side panels for directions for use, first aid statement and additional precautionary statements.

EPA Reg. No. 37549-1 EPA Est. No. 10088-WI-2
73027-WI-01

NET CONTENTS:

55

PRE-SATURATED WIPES
12 in x 12 in (30.5 cm x 30.5 cm)
Net Wt: 2 lbs 1.58 oz (952 g)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

HOSPITAL DISINFECTION

1. Always use personal protective equipment.*
2. Open Micro-Kill Bleach Germicidal Bleach Wipes canister.
3. Remove pre-saturated 12 in x 12 in wipe.
4. Apply pre-saturated towelette and wipe desired surface to be disinfected.
5. Gross soil must be removed prior to disinfecting. A 30 second contact time is required to kill the bacteria and viruses** on the label except 1 minute contact time is required to kill *Candida albicans* and *Trichophyton interdigitale*, a 2 minute contact time is required to kill *Candida auris****, and a 3 minutes contact time is required to kill *Clostridium difficile* spores*. Reapply as necessary to ensure that the surface remains visibly wet for the entire contact time.
6. Allow surface to air dry and discard used wipe and empty canister (see **STORAGE AND DISPOSAL**).

TO OPEN CANISTER:

1. Always use personal protective equipment.*
2. Remove lid from canister.
3. Remove seal from mouth of canister.
4. Lift flap on canister lid.
5. Pull up corner of center wipe and snap off and discard. Take the next wipe, twist to a point and thread through dispenser hole in lid.
6. DO NOT PUSH FINGER THROUGH OPENING.
7. Replace lid. Pull out exposed wipe and snap off. The next wipe pops up automatically.
8. Snap center flap down when finished to retain moisture.
9. Do not flush wipe down the toilet.

***KILLS HIV-1, HBV AND HCV ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids and in which the surfaces/objects likely to be soiled with blood/body fluids can be associated with the potential for transmission of HIV-1 (associated with AIDS), HBV and HCV.

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV, AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

- **PERSONAL PROTECTION:** When handling items soiled with blood or body fluids, use disposable gloves, gown, masks and eye protection.
- **CLEANING PROCEDURE:** Blood/body fluids must be thoroughly cleaned from surfaces/objects before application of Micro-Kill Bleach Germicidal Bleach Wipes.
- **CONTACT TIME:** Allow surface(s) to remain visibly wet for 30 seconds to kill the bacteria and viruses** on the label except a 1 minute contact time is required to kill *Candida albicans* and *Trichophyton interdigitale*, a 2 minute contact time is required to kill *Candida auris****, and a 3 minute contact time is required to kill *Clostridium difficile* spores*.
- **DISPOSAL OF INFECTIOUS MATERIAL:** Blood/body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

FRONT

MEDLINE

MICRO-KILL™ Bleach

GERMICIDAL BLEACH WIPES

**3 MINUTE
KILL TIME**

<p>BACTERICIDAL</p> <ul style="list-style-type: none"> 30 second contact time <i>Acinetobacter baumannii</i>-MDR <i>Bordetella pertussis</i> <i>Campylobacter jejuni</i> CA-Methicillin Resistant <i>Staphylococcus aureus</i> <i>Enterobacter aerogenes</i> <i>Enterobacter cloacae</i> <i>Enterococcus faecalis</i>-VRE <i>Enterococcus faecium</i>-MDR <i>Escherichia coli</i> 	<ul style="list-style-type: none"> <i>Escherichia coli</i> O157:H7 <i>Escherichia coli</i>-ESBL <i>Escherichia coli</i>-NDM-1 <i>Klebsiella pneumoniae</i> <i>Klebsiella pneumoniae</i>-CRKP (CRE) <i>Klebsiella pneumoniae</i>-ESBL <i>Klebsiella pneumoniae</i>-NDM-1 <i>Legionella pneumophila</i> Linezolid Resistant <i>Staphylococcus aureus</i> <i>Listeria monocytogenes</i> Methicillin Resistant 	<ul style="list-style-type: none"> <i>Staphylococcus aureus</i> (MRSA) <i>Proteus mirabilis</i> <i>Pseudomonas aeruginosa</i> <i>Salmonella enterica</i> <i>Serratia marcescens</i> <i>Shigella dysenteriae</i> <i>Staphylococcus aureus</i> <i>Staphylococcus aureus</i>-VISA <i>Staphylococcus aureus</i>-VRSA <i>Streptococcus pneumoniae</i> <i>Streptococcus pyogenes</i>
---	--	---

***SPORICIDAL**

- 3 minute contact time
- Clostridium difficile* spores*

****VIRUCIDAL**

- 30 second contact time
- Adenovirus Type 5
- Avian Influenza A Virus (H3N2)
- Avian Influenza A Virus (H5N1)
- Avian Influenza A Virus (H7N9)
- Bovine Viral Diarrhea Virus (BVD) (HCV)
- Canine Parvovirus
- Cytomegalovirus (Strain AD-169)

- Duck Hepatitis B Virus (DHBV) (HBV)
- Enterovirus D68
- Feline Calicivirus (ATCC VR-782)
- Feline Panleukopenia Virus
- Hepatitis A Virus
- +Hepatitis B Virus
- +Hepatitis C Virus
- Herpes Simplex Virus Type 2
- Human Coronavirus
- +Human Immunodeficiency Virus Type 1
- Influenza A Virus (Novel H1N1)
- Influenza A Virus (Hong Kong)

- Influenza B Virus (Strain B/Hong Kong/5/72)
- Norovirus (Feline Calicivirus)
- Polio Virus Type 1
- Respiratory Syncytial Virus
- Rhinovirus Type 37
- Rotavirus

FUNGICIDAL

- 1 minute contact time
- Candida albicans*
- Trichophyton interdigitale*
- 2 minute contact time
- Candida auris****

These wipes are sporidical*, virucidal**, bactericidal and fungicidal*** and can be used to clean, deodorize and disinfect hard, non-porous healthcare and environmental surfaces, patient care equipment and point-of-care equipment in one-step*.

Equivalent to 1:10 dilution of 6.5% bleach

Active Ingredient: Sodium hypochlorite	0.65%
Other Ingredients*:	99.35%
Total	100.00%

* Other ingredients do not include the weight of wipes.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SUGGESTED AREAS OF USAGE:

Medical, Dental and Laboratory Counters, Exam Tables, Carts, Patient Care Equipment, Point-of-Care Equipment, Telephones, Sink Tops, Toilet Seats, Digital Screens, on Plastics, Glazed Porcelain, Stainless Steel, Formica, Vinyl Surfaces and Blood Glucose Meters†.

See back panel for directions for use, first aid statement and additional precautionary statements.

EPA Reg. No. 37549-1
EPA Est. No. 10088-WI-2

73027-WI-01

[illegible]

MEDLINE

MICRO-KILL™ Bleach

GERMICIDAL BLEACH WIPES & BUCKET



BACTERICIDAL

30 second contact time
Acinetobacter baumannii-MDR
Bordetella pertussis
Campylobacter jejuni
CA-Methicillin Resistant
Staphylococcus aureus
Enterobacter aerogenes
Enterobacter cloacae
Enterococcus faecalis-VRE
Enterococcus faecium-MDR
Escherichia coli
Escherichia coli O157:H7
Escherichia coli-ESBL
Escherichia coli-NDM-1
Klebsiella pneumoniae

Klebsiella pneumoniae-CRKP (CRE)

Klebsiella pneumoniae is a member of the family *Enterobacteriaceae*.
Klebsiella pneumoniae-ESBL
Klebsiella pneumoniae-NDM-1
Legionella pneumophila
Linezolid Resistant
Staphylococcus aureus
Listeria monocytogenes
Methicillin Resistant
Staphylococcus aureus (MRSA)
Proteus mirabilis
Pseudomonas aeruginosa
Salmonella enterica

Serratia marcescens
Shigella dysenteriae
Staphylococcus aureus

Staphylococcus aureus-VISA
Staphylococcus aureus-VRSA
Streptococcus pneumoniae
Streptococcus pyogenes

*SPORICIDAL

3 minute contact time
Clostridium difficile spores*

**VIRUCIDAL

30 second contact time
Adenovirus Type 5
Avian Influenza A Virus (H3N2)
Avian Influenza A Virus (H5N1)

Avian Influenza A Virus (H7N9)

Bovine Viral Diarrhea Virus (BVDV) (HCV)
Canine Parvovirus
Cytomegalovirus (Strain AD-169)
Duck Hepatitis B Virus (DHBV) (HBV)
Enterovirus D68
Feline Calicivirus (ATCC VR-782)
Feline Panleukopenia Virus
Hepatitis B Virus
+Hepatitis C Virus
Herpes Simplex Virus Type 2
Human Coronavirus
+Human Immunodeficiency Virus Type 1

Influenza A Virus (Novel H1N1)

Influenza A Virus (Hong Kong)
Influenza B Virus (Strain B/Hong Kong/5/72)
Norovirus (Feline Calicivirus)
Polo Virus Type 1
Respiratory Syncytial Virus
Rhinovirus Type 37
Rotavirus

FUNGICIDAL

1 minute contact time
Candida albicans
Trichophyton interdigitale
2 minute contact time
*Candida auris****

Klebsiella pneumoniae is a member of the family *Enterobacteriaceae*.



These wipes are sporidical*, virucidal**, bactericidal and fungicidal*** and can be used to clean, deodorize and disinfect hard, non-porous healthcare and environmental surfaces, patient care equipment and point-of-care equipment in one-step*.

Active Ingredient: Sodium hypochlorite	0.65%
Other Ingredients**	99.35%
Total	100.00%

*Other ingredients do not include the weight of wipes.

KEEP OUT OF REACH OF CHILDREN CAUTION

SUGGESTED AREAS OF USAGE:

Medical, Dental and Laboratory Counters, Exam Tables, Garts, Patient Care Equipment, Point-of-Care Equipment, Telephones, Sink Tops, Toilet Seats, Digital Screens, on Plastics, Glazed Porcelain, Stainless Steel, Formica, Vinyl Surfaces and Blood Glucose Meters.†
See back panel for directions for use, first aid statement and additional precautionary statements.

EPA Reg. No. 37549-1 EPA Est. No. 10088-WI-2, 73027-WI-01

NET CONTENTS: **110** PRE-SATURATED WIPES
12 in x 12 in (30.5 cm x 30.5 cm)
Net Wt: 4 lbs 5.96 oz (2085g)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

TO FILL OR REFILL BUCKETS

1. Always use personal protective equipment.
2. To fill or refill the bucket with towels, place the sealed pouch containing the roll of towels in the bucket and then cut open the pouch.
3. Pull up corner of center towel and snap off and discard. Take the next towel and twist to a point and thread through dispenser hole in lid.
4. DO NOT PUSH FINGER THROUGH OPENING.
5. Replace lid. Pull out exposed towel and snap off. The next towel pops up automatically.
6. Snap center flap down when finished to retain moisture.
7. Do not flush towel down the toilet.
8. Refer to label on the bucket for DIRECTIONS FOR USE.

HOSPITAL DISINFECTION

1. Always use personal protective equipment.*
2. Open Micro-Kill Bleach Germicidal Bleach Wipes pouch.
3. Remove pre-saturated 12 in x 12 in wipe.
4. Apply pre-saturated towelette and wipe desired surface to be disinfected.
5. Gross soil must be removed prior to disinfecting. A 30 second contact time is required to kill the bacteria and viruses** on the label except 1 minute contact time is required to kill *Candida albicans* and *Trichophyton interdigitale*, a 2 minute contact time is required to kill *Clostridium difficile* spores*. Resupply as necessary to ensure that the surface remains visibly wet for the entire contact time.
6. Allow surface to air dry and discard used wipe and empty bucket (see STORAGE AND DISPOSAL).

*KILLS HIV-1, HBV AND HCV ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids and in which the surfaces/objects likely to be soiled with blood/body fluids can be associated with the potential for transmission of HIV-1 (associated with AIDS), HBV and HCV.

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION

AGAINST HIV-1, HBV, AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

- **PERSONAL PROTECTION:** When handling items soiled with blood or body fluids, use disposable gloves, gown, masks and eye protection.
- **CLEANING PROCEDURE:** Blood/body fluids must be thoroughly cleaned from surfaces/objects before application of Micro-Kill Bleach Germicidal Bleach Wipes.
- **CONTACT TIME:** Allow surface(s) to remain visibly wet for 30 seconds to kill the bacteria and viruses** on the label except a 1 minute contact time is required to kill *Candida albicans* and *Trichophyton interdigitale*, a 2 minute contact time is required to kill *Candida auris****, and a 3 minute contact time is required to kill *Clostridium difficile* spores*.
- **DISPOSAL OF INFECTIOUS MATERIALS:** Blood/body fluids must be autotested and disposed of according to local regulations for infectious waste disposal.

*SPECIAL INSTRUCTIONS FOR CLEANING PRIOR TO DISINFECT-ING AGAINST Clostridium difficile SPORES.

- **PERSONAL PROTECTION:** Wear appropriate barrier protection such as gloves, gown, mask, or eye covering.
- **CLEANING PROCEDURE:** Fecal matter /waste must be thoroughly cleaned from surfaces/objects before disinfection by application with Micro-Kill Bleach Wipes. Cleaning is to include vigorous wiping and /or scrubbing, until all visible soil is removed. Special attention is needed for high-touch surfaces. Surfaces in patient rooms are to be cleaned in an appropriate manner, such as from right to left or left to right, on horizontal surfaces, and top to bottom, on vertical surfaces, to minimize spreading of the spores. Restrooms are to be cleaned last. Do not reuse soiled wipes.

- **INFECTIOUS MATERIALS DISPOSAL:** Materials used in the cleaning process that may contain feces /wastes are to be disposed of immediately in accordance with local regulations for infectious materials disposal.

***SPECIAL INSTRUCTIONS FOR CLEANING PRIOR TO DISINFECTING AGAINST Candida auris:

- **PERSONAL PROTECTION:** Wear appropriate barrier protection such as gloves, gown, masks, or eye covering.

- **CLEANING PROCEDURE:** Fecal matter/waste must be thoroughly cleaned from surfaces/objects before disinfection by application with a clean cloth, mop, and/or sponge saturated with the product. Pre-cleaning is to include vigorous wiping and/or scrubbing and all visible soil is removed. Surfaces in patient rooms are to be cleaned in an appropriate manner, such as from right to left or left to right, on horizontal surfaces, and top to bottom, on vertical surfaces, to minimize spreading of the organisms. Restrooms are to be cleaned last. Do not reuse soiled cloths.

- **INFECTIOUS MATERIALS DISPOSAL:** Materials used in the cleaning process that may contain feces/wastes are to be disposed of immediately in accordance with local regulations for infectious materials disposal.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barriers or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

† This product is for the exterior surfaces of Blood Glucose Meters and is not for use on the needle and/or the monitors.

*Follow the Special Instructions for Cleaning prior to disinfection for *Clostridium difficile* spores*.

+Follow the Special Instructions for Cleaning prior to disinfection for bloodborne pathogens.

*A pre-cleaning step is required for efficacy against +HIV, +HBV, +HCV, *Candida auris****, and *Clostridium difficile* spores*.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. Have the product container with you when calling a poison control center or doctor, or going for treatment.

PRECAUTIONARY STATEMENTS

Hazards to humans and domestic animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PHYSICAL AND CHEMICAL HAZARDS

This product contains bleach. Do not use this product with other chemicals such as ammonia, toilet bowl cleaners, rust removers, or acid, as this releases hazardous gases.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

Pesticide Storage:

Keep this product in a tightly closed bucket, when not in use. Store bucket in cool, dry, and well-ventilated area away from heat, open flame, direct sunlight and inaccessible to children.

Pesticide Disposal:

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling and Disposal:

Refillable bucket. Refill this bucket with Micro-Kill Bleach Germicidal Bleach Wipes only. Do not reuse this bucket for any other purpose. To clean the bucket before final disposal, triple rinse (or equivalent), promptly after emptying. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.



Manufactured for Medline Industries Inc.,
Three Lakes Drive, Northfield, IL 60093.
Medline Micro-Kill is a trademark of Medline Industries, Inc. 1-800-MEDLINE Made in the USA RK19MGW

REF MSC351450AN



For product emergency, contact:
Poison Control Center, 1-800-222-1222

For Technical Information call: 1-800-MEDLINE

For an SDS, call 1-800-MEDLINE

and request document MSC351450AN

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barriers or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

†This product is for the exterior surfaces of Blood Glucose Meters and is not for use on the needle and/or the monitors.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. Have the product container with you when calling a poison control center or doctor, or going for treatment.

PRECAUTIONARY STATEMENTS Hazards to humans and domestic animals.

CAUTION: Causes moderate eye irritation. Avoid contact with eyes and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PHYSICAL AND CHEMICAL HAZARDS

This product contains bleach. Do not use this product with other chemicals such as ammonia, toilet bowl cleaners, rust removers, or acid, as this releases hazardous gases.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Keep this product in a tightly closed canister, when not in use. Store canister in cool, dry, and well-ventilated area away from heat, open flame, direct sunlight and inaccessible to children.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling and Disposal: Nonrefillable rigid canister. Do not re-use or refill this canister. Triple rinse (or equivalent), promptly after emptying. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Follow the Special Instructions for Cleaning prior to disinfection for Clostridium difficile spores.

+Follow the Special Instructions for Cleaning prior to disinfection for bloodborne pathogens.

^A pre-cleaning step is required for efficacy against +HIV, +HBV, +HCV, Candida auris**, and Clostridium difficile spores*.

REF MSC351410AN



(01)10888277652521



Manufactured for Medline Industries Inc.,
Three Lakes Drive, Northfield, IL 60093.
Micro-Kill is a trademark of Medline Industries,
Inc. 1-800-MEDLINE. Made in USA. RL19ATH

For product emergency, contact: Poison Control Center, 1-800-222-1222

For Technical Information call: 1-800-MEDLINE. For a SDS
call 1-800-MEDLINE and request document MSC351410AN

MICRO-KILL™ Bleach
GERMICIDAL BLEACH WIPES



*SPORICIDAL 3 minute contact time
Clostridium difficile spores*

FUNGICIDAL 1 minute contact time
Candida albicans
Trichophyton interdigitale

***FUNGICIDAL 2 minute contact time
Candida auris



Equivalent to 1:10 dilution of 6.5% bleach.

Active Ingredient: Sodium hypochlorite.....	0.65%
Other Ingredients**.....	99.35%
Total	100.00%

~Other ingredients do not include the weight of wipe.

BACTERICIDAL 30 second contact time
Acinetobacter baumannii-MDR
Bordetella pertussis
Campylobacter jejuni
CA-Methicillin Resistant Staphylococcus aureus
Enterobacter aerogenes
Enterobacter cloacae
Enterococcus faecalis-VRE
Enterococcus faecium-MDR
Escherichia coli

These wipes are sporicidal*, virucidal**, bactericidal and fungicidal*** and can be used to clean, deodorize and disinfect hard, non-porous healthcare and environmental surfaces, patient care equipment and point-of-care equipment in one-step^.

Escherichia coli O157:H7
Escherichia coli-ESBL
Escherichia coli-NDM-1
Klebsiella pneumoniae
Klebsiella pneumoniae-CRKP (CRE)
Klebsiella pneumoniae-ESBL
Klebsiella pneumoniae-NDM-1
Legionella pneumophila
Linezolid Resistant Staphylococcus aureus
Listeria monocytogenes
Methicillin Resistant Staphylococcus aureus (MRSA)
Proteus mirabilis
Pseudomonas aeruginosa
Salmonella enterica
Serratia marcescens
Shigella dysenteriae
Staphylococcus aureus
Staphylococcus aureus-VISA
Staphylococcus aureus-VRSA
Streptococcus pneumoniae
Streptococcus pyogenes
Klebsiella pneumoniae is a member of the family Enterobacteriaceae.

KEEP OUT OF REACH OF CHILDREN
CAUTION

SUGGESTED AREAS OF USAGE:

Medical, Dental and Laboratory Counters, Exam Tables, Carts, Patient Care Equipment, Point-of-Care Equipment, Telephones, Sink Tops, Toilet Seats, Digital Screens, on Plastics, Glazed Porcelain, Stainless Steel, Formica, Vinyl Surfaces and Blood Glucose Meters†

See side panels for directions for use, first aid statement and additional precautionary statements.

EPA Reg. No. 37549-1 EPA Est. No. 10088-WI-2
73027-WI-01

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

HOSPITAL DISINFECTION

1. Always use personal protective equipment.*
2. Open Micro-Kill Bleach Germicidal Bleach Wipes canister.
3. Remove premoistened 6 in x 5 in wipe.
4. Apply pre-saturated towelette and wipe desired surface to be disinfected.
5. Gross soil must be removed prior to disinfecting. A 30 second contact time is required to kill the bacteria and viruses** on the label except 1 minute contact time is required to kill Candida albicans and Trichophyton interdigitale, a 2 minute contact time is required to kill Candida auris***, and a 3 minutes contact time is required to kill Clostridium difficile spores*. Reapply as necessary to ensure that the surface remains visibly wet for the entire contact time.
6. Allow surface to air dry and discard used wipe and empty canister (see STORAGE AND DISPOSAL).

TO OPEN CANISTER:

1. Always use personal protective equipment.*
2. Remove lid from canister.
3. Remove seal from mouth of canister.
4. Lift flap on canister lid.
5. Pull up corner of center wipe and snap off and discard. Take the next wipe, twist to a point and thread through dispenser hole in lid.
6. DO NOT PUSH FINGER THROUGH OPENING.
7. Replace lid. Pull out exposed wipe and snap off. The next wipe pops up automatically.
8. Snap center flap down when finished to retain moisture.
9. Do not flush wipe down the toilet.

*KILLS HIV-1, HBV AND HCV ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS

in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids and in which the surfaces/objects likely to be soiled with blood/body fluids can be associated with the potential for transmission of HIV-1 (associated with AIDS), HBV and HCV.

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV, AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

- PERSONAL PROTECTION: When handling items soiled with blood or body fluids, use disposable gloves, gown, masks and eye protection.
- CLEANING PROCEDURE: Blood/body fluids must be thoroughly cleaned from surfaces/objects before application of Micro-Kill Bleach Germicidal Bleach Wipes.
- CONTACT TIME: Allow surface(s) to remain visibly wet for 30 seconds to kill the bacteria and viruses** on the label except a 1 minute contact time is required to kill Candida

albicans and Trichophyton interdigitale, a 2 minute contact time is required to kill Candida auris***, and a 3 minute contact time is required to kill Clostridium difficile spores*.

• DISPOSAL OF INFECTIOUS MATERIAL: Blood/body fluids must be autoclaved and disposed of according local regulations for infectious waste disposal.

*SPECIAL INSTRUCTIONS FOR CLEANING PRIOR TO DISINFECTING AGAINST Clostridium difficile SPORES.

- PERSONAL PROTECTION: Wear appropriate barrier protection such as gloves, gown, mask, or eye covering.
- CLEANING PROCEDURE: Fecal matter/waste must be thoroughly cleaned from surfaces/objects before disinfection by application with Micro-Kill Bleach Germicidal Bleach Wipes. Cleaning is to include vigorous wiping and/or scrubbing, until all visible soil is removed. Special attention is needed for high-touch surfaces. Surfaces in patient rooms are to be cleaned in an appropriate manner, such as from right to left or left to right, on horizontal surfaces, and top to bottom, on vertical surfaces, to minimize spreading of the spores. Restrooms are to be cleaned last. Do not reuse soiled wipes.

• INFECTIOUS MATERIAL DISPOSAL: Materials used in the cleaning process that may contain feces/ wastes are to be disposed of immediately in accordance with local regulations for infectious materials disposal.

***SPECIAL INSTRUCTIONS FOR CLEANING PRIOR TO DISINFECTING AGAINST Candida auris:

- PERSONAL PROTECTION: Wear appropriate barrier protection such as gloves, gowns, masks, or eye covering.
- CLEANING PROCEDURE: Fecal matter/waste must be thoroughly cleaned from surfaces/objects before disinfection by application with a clean cloth, mop, and/or sponge saturated with the product. Pre-cleaning is to include vigorous wiping and/or scrubbing and all visible soil is removed. Surfaces in patient rooms are to be cleaned in an appropriate manner, such as from right to left or left to right, on horizontal surfaces, and top to bottom, on vertical surfaces, to minimize spreading the organism. Restrooms are to be cleaned last. Do not reuse soiled cloths.
- INFECTIOUS MATERIAL DISPOSAL: Materials used in the cleaning process that may contain feces/wastes are to be disposed of immediately in accordance with local regulations for infectious materials disposal.