

ALL-IN-ONE LAWN WEED KILLER*

READY-TO-SPRAY



DANDELION



NUTSEDGE



CLOVER

WEEDS



*Controls Listed Annual Grasses, Broadleaf Weeds and Sedges **When used as directed

COVERS 6,000 FT.

CPABGRASE

Active Ingredients:	By Wt.
Sulfentrazone	1.6%
Quinclorac	5.0%
Other Ingredients	93.4%
	100.0%

NET CONTENTS 24 FL OZ (710 mL)



READY-TO-SPRAY

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CALITION

Harmful if svallowed or absorbed through skin. Causes moderate ever initiation. Avoid contact with eyes, skin, or dothing. Wear long-sleeed shirt, long pants, shoes, socks and welending, gloves. Wear throughly with soap and water after handling and before eating, drinking, thewing gurn, using tobacco, or using the toilet. Remove contaminated dothing and wesh hefore reise.

See inside booklet for additional Precautionary Statements and complete Directions

,		
	FIRST AID	
If swallowed	Call poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.	
lf on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.	
If in eyes	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continus rinsing eye. Call a poson control center or doctor for treatment advice. 	
HOTH INF MUMBER		

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-265-0761 for emergency medical treatment information.

STORAGE AND DISPOSAL Pesticide Storage

Keep from freezing. Do not store below 32°F. Keep out of reach of children and animals. Store in original container only. Store in a cool, dry place, preferably in a locked storage area and avoid excess heat.

in case of a spill, avoid contact, footbe area and keep out animals and unprotected person confine spills. Call Central Garden & Pet, Garden Division: 1-800-265-016. <u>Ioconfine spill</u>: If liquid, like surrounding area or absorb with sand, call titer or commercial day. If dry material, cover to prevent dispersal. Place damaged package in a holding contains: identify contents.

Pesticide Disposal

Pedicide westes are acutely hazardous. Improper disposal of excess pedicide, spray minture, or misate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pedicide of Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for quidants.

Container Handling

Non-refillable container. Do not reuse or refill this container. If empty: Offer for recycling, if available, or dispose of in trash.

If partially filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Manufactured for: Central Garden & Pet, Garden Division 1000 Parkwood Circle, Suite 700 Atlanta. GA 30339

1-800-265-0761 gardendivision@central.com EPA Reg No. 279-3463-90098 EPA Est. No. 59144-GA-1_a, 67572-GA-1_b, 67572-GA-2_c, 74750-GA-1_b, 70815-GA-1_t 5905-GA-1_t

Subscript used is first letter of Run Code on container.



© 2015-2023 Central Garden & Pet Company 300534282 - 01/23 | IM197 V5 KILLS WEEDS FAST



ALL-IN-ONE LAWN WEED KILLER

READY-TO-SPRAY

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Observe all cautions and limitations on this label. The use of this product not consistent with this label may result in plant injury. Keep container closed to avoid solils and contamination.

APPLICATION INSTRUCTIONS AND PRECAUTIONS

п					
ı	W	н	Ε	R	Ε
ı	TC)	U	s	E

Well established turfgrass

Cool Season Grasses: Fine & Tall Fescue, Perennial Ryegrass
Kentucky & Rough Bluegrass.

Warm Season Grasses: Common & Hybrid Bermudagrass, Buffalograss, Centipedegrass, Seashore Paspalum, Zoysiagra

KILLS

Crabgrass, Foxtail, Ground Iwy, Wild Violet, Plantain, Dandelion, Clover, Dollarweed, Spurge, Woodsorrel, Knotweed, Chickweed, Curly Dock, Wild Onion, Wild Garlic, and other listed weeds.

WHEN TO USE HOW MUCH TO USE

4fl. oz. per 1 gallon of water per 1,000 ft². Spot spray weed directly until thoroughly covered. Some weeds may require repeat application. (See detailed

COVERAGE

24 fl.oz. covers up to 6.000 sa.f

See inside booklet for additional Precautionary Information, First Aid and complete Directions For Use.

Manufactured for: Central Garden & Pet, Garden Division 1000 Parkwood Circle, Suite 700 Atlanta, GA 30339

1-800-265-0761 gardendivision@central.com EPA Reg No. 279-3463-90098 EPA Est. No. 59144-GA-1₈, 67572-GA-1₈, 67572-GA-2_c, 74750-GA-1₈, 70815-GA-1_e, 5905-GA-1_e subscript used is first letter of Run Code on container.

Image, Central Garden & Pet, and Central Garden & Pet with design are trademarks of Central Garden & Pet Company.

CENTRAL
GARDEN & PET

300534282 - I/25 | IM197 V5

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed or absorbed through skin, Causes moderate eve irritation. Avoid contact with eyes, skin, or clothing. Wear long-sleeved shirt, long pants. shoes, socks and waterproof gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to marine/estuarine invertebrates. To protect the environment. do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems.

This product may impact surface water quality due to runoff of rain water or spray drift. Under certain conditions this product may have a high potential for runoff into surface water for several months after application. This is especially true for poorly draining soils and soils with shallow ground water.

The active ingredients in this product are known to leach through soil into groundwater under certain conditions as a result of labeled use. Use of this product in areas where soils are permeable, particularly where the water table is shallow. may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Observe all cautions and limitations on this label. The use of this product not consistent with this label may result in plant injury. Keep container closed to avoid spills and contamination.

PRODUCT INFORMATION

This product selectively controls tough weeds like crabgrass, chickweed, dandelion. dover, ground ivy, purplet and yellow nutsedge (nutgrass), Kyllinga, wild onion and garlic, broadleaf plantain, purslane, wild violet, spurge, and other listed annual. biennial, and broadleaf weeds in a variety of lawns, including:

Cool Season Grasses

Tall & Fine Fescue, Kentucky Bluegrass, Rough Bluegrass, Ryegrass (perennial)

Warm Season Grasses

Common & Hybrid Bermudagrass, Zoysiagrass, Centipedegrass, Buffalograss, Seashore Paspalum

Not for use in California

This product has demonstrated tolerance on both cool and warm season turfgrasses. However, not all varieties have been evaluated. Some turf varieties are more susceptible to temporary turf discoloration than others. It is recommended to treat a small area prior to treatment of larger areas.

This unique formula causes visual symptoms to develop quickly within 3 to 7 days after application. This product involves uptake by both weed roots and shoots. Complete kill should occur in 2-3 weeks. This product also controls yellow nutsedge tubers (nutlets) in the soil over time.

A postemergence application of this product is improved when adequate soil moisture is present at application.

HOW TO APPLY:

Ready to Spray Hose-End Sprayer

1. Measure the entire area to be sprayed. For rectangular areas, multiply the length by width to obtain total square feet

2 See below for directions for the ready-to-use hose-end sprayer.

3 Snray evenly over the entire measured area

5. Spray evenily over the challe measured area.	1 1
Hose End Sprayer Connection	2
1. Shake well.	4
2. Make sure control lever is in the OFF position.	6
Attach to hose.	8
3. Turn on water.	24

Product of Area Size: fl. 07. 250 sa.ft. 7 fl. oz. 500 sq.ft. 4 fl. oz. 1000 sq.ft. 1500 sq.ft. 5 fl 07 B fl. 07. 2000 sa.ft. 4 fl. oz. 6000 sa.ft.

Amount of | Provides Coverage

Spray

1. Aim towards direction of application and away from you.

2. Turn control lever to ON to begin spraying.

Spray evenly and as directed for specified type of treatment area.

Ston

1. When finished turn control lever to OFF.

Turn faucet OFF. Discharge residual water pressure by turning control lever ON and OFF again

Disconnect sprayer and store / discard as instructed.

COVERAGE: 4 fl.oz. per 1000 sq. ft.

APPLICATION INSTRUCTIONS:

 Best weed control results will be obtained when no rainfall or irrigation occurs within 24 hours after application.

· Listed sedges and grassy weeds: Spray directly on emerged weeds. This product is effective against newly emerged and established sedges and grassy weeds.

· Listed broadleaf weeds: Spray directly on newly emerged weeds. For best results, apply to weeds when less than 3 inches high.

Annly in calm weather

 Begin spraying at the farthest point of the treatment area and work backward to avoid contact with wet treated surfaces.

Good spray coverage is needed for optimum control.

Avoid overlap spraving.

Avoid side to side or swinging arm motion, as it can result in uneven coverage.

 If listed weeds are not controlled by the initial application, apply a second application 4 to 6 weeks after the initial treatment.

 Water lawn with at least 0.5 inches within 7 days (but not sooner than 1 day after application) if no rainfall or irrigation occurs within 7 days after application.

APPLICATION RESTRICTIONS

Keep children and pets out of the treated area until sprays have dried. Do not spray over the top of flowers, ornamentals (including herbaceous) or garden vegetables.

Do not use on turforasses other than those listed on the label. Do not use in greenhouses.

Do not graze or feed animals with treated grass clippings. Do not apply to newly planted, plugged, or sodded turfgrass.

APPLICATION PRECAUTIONS

Use only on well established non-stressed turfgrass. Applications to stressed plants (turfgrass or target weeds) may result in temporary discoloration or poor weed control. Allow turfgrass to develop good root system before applying.

If discoloration occurs, treated turfgrass will recover with new growth and discolored leaf tissue will be removed with mowing. To reduce potential for discoloration, do not apply on turfgrass that is weakened by weather, mechanical, chemical, disease or other related stress. Maintain proper cultural practices such as adequate moisture and fertility levels to promote healthy turfgrass growth.

Avoid applying when windy to prevent product drifting to other plants. Avoid reseeding, winter overseeding, or plugging lawns for at least 4 weeks after treatment

Grass and sedge weed species controlled or suppressed by this product:

,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,			
Common Name	Scientific Name	Common Name	Scientific Name
Barnyardgrass	(Echinochloa crus-galli)	Kyllinga, fragrant	(Kyllinga sesguiflorus)
Broadleaf Signalgrass	(Brachiaria platyphylla)	Kyllinga, tufted	(Kyllinga pumila)
Crabgrass, Large			(Microstegium vimineum)
Crabgrass, Smooth	(Digitaria ischaemum)	Junglerice	(Echinochloa colona)
Deergrass	(Panicum dadestinum)	Nutsedge, purple'	(Cyperus rotundus)
Goosegrass	(Eleusine indica)	Nutsedge, yellow	(Cyperus esculentus)
Foxtail, giant	(Setaria faberi)	Sedge, annual	(Cyperus compressus)
Foxtail, green	(Setaria viridis)	Sedge, globe	(Cyperus globulosus)
Foxtail, yellow		Sedge, cylindrical	(Cyperus retrorsus)
Kyllinga, cocks-comb		Sedge, Surinam	(Cyperus surinamensis)
Kyllinga, green		Sedge, Texas	(Cyperus polystachyos)
Kyllinga, false green	(Kyllinga gracillima)		

For optimal control of Goosegrass, Junglerice, or Purple Nutsedget, repeat treatment every 3-4 weeks

†Not for use in California

Broadleaf weeds controlled or suppressed by this product:

suppressed by this product.	
Common Name	Scientific Name
Bedstraw, catchweed	(Galium aparine)
Beggarticks†	(Bidens frondosa)
Beggarweed, Florida	(Desmodium tortuosum)
Bindweed, field	(Convolvulus arvensis)
Bittercress	(Cardamine spp.)
Black medic	(Medicago lupuline)
Buttercups	(Ranunculus spp.)
Buttonweed, common (poorjoe)	(Diodia teres)
Carolina geranium	(Geranium carolinianum)
Carpetweed	(Mollugo verticillata)
Chickweed, common	(Stellaria media)
Chickweed, mouseear	(Cerastium vulgatum)
Cinquefoil	(Potentilla spp.)
Clover	(Trifolium spp.)
Clover, alyce	(Alysicarpus ovalifolius)
Clover, hop	(Trifolium campestre)
Clover, red	(Trifolium pretense)
Clover, white	(Trifolium repens)
Cudweed	(Gnaphalium spp.)
Daisy, English	(Bellis perennis)
Dandelion	(Taraxacum officianale)
Dandelion, false	(Microseris cuspidate)
Dayflower, common	(Commelina diffusa)
Dock, Curly	(Rumex crispus)
Dollarweed (pennywort)	(Hydrocotyle spp.)
Edipta	(Eclipta prostrata)
Evening primrose	(Oenother biennis)
Fiddleneck	(Amsinckia spp.)
Filaree	(Erodium spp.)
Filaree, redstem	(Erodium cicutarium)
Galinsoga	(Galinsoga ciliate)
Goldenrod	(Solidago spp.)
Ground ivv	(Glechmea hederacea)
Groundsel, common	(Senecio vulgaris)
Henbit	(Lamium amplexicaule)
Knawel	(Scleranthus annuus)
Knotweed, prostrate	(Polygonum aviculare)
Kochia	(Kockia scoparia)
Lambsquarters, common	(Chenopodium album)
Lawn burweed (Spurweed)	(Soliva pterosperma)
Lespedeza, common	(Lespedeza striafa)
London Rocket	(Sisymbrium irio)
Mallow, common	(Malva neglecta)
Morning glory	(Ipomea spp.)
Morning glory, entireleaf	(Ipomoea hederacea var. integriuscula)
Morning glory, ivyleaf	(Ipomoea hederacea)
Morning glory, palmleaf	(Ipomoea wrightii)
Morning glory, pitted	(Ipomoea lacunose)
Morning glory, tall	(Ipomoea purpurea)
Parsley piert	(Alchemilla arvensis)
Pigweed, Redroot	(Amaranthus retroflexus)
Pigweed Smooth	(Amaranthus hybridus)
Pigweed, Smooth Pigweed, Tumble	(Amaranthus albus)
Pineapple weed	(Matricaria matricariodes)
Plantain, Broadleaf	(Plantago comanalata)
Plantain, Buckhorn	(Plantago lanceolata)
Plantain, Narrowleaf	(Plantago narowiforma)
Puncturevine	(Tribulus terrestris)
Purslane, common	(Portulaca oleracea)
Pusley, Florida	(Richardia scabra)
Redweed	(Melochia corchorifolia)
Shepherd's-Purse	(Capsella bursa-pastoris)
Smartweed, Pennsylvania	(Polygonum pensylvanicum)
Sorrel, Red	(Rumex acetosella)
Speedwell	(Veronica spp.)
Speedwell, slender	(Veronica filiformis)
Speedwell, thymeleaf	(Veronica serpyllifolia)
Spurge, (annuals)	(Euphorbia spp.)
Spurge, prostrate	(Euphorbia humistrata)
Spurge, spotted	(Euphorbia maculate)
-p-3-, spoucu	(Euprioridia Hacaiate)

Broadleaf weeds controlled or suppressed by this product (cont'd):

Common Name	Scientific Name	
Star-of-Bethlehem Velvetleaf Violet, wild Violet, Johnny-Jump-Up Wild garlic Wild onion Woodsorrel, yellow	(Omithogalum umbellatum) (Abutilon theophrasti) (Viola pratiriona) (Viola raleinesquii) (Allium vineale) (Allium anaderse) (Daalis comiculata) (Okalis stricta)	

*Not for use in California

Certain listed weeds may require more than one application to achieve season-long control

STORAGE AND DISPOSAL

Pesticide Storage

Keep from freezing. Do not store below 32°F. Keep out of reach of children and animals. Store in original container only. Store in a cool, dry place, preferably in a locked storage area and avoid excess heat.

In case of a spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call Central Garden & Pet, Garden Division: 1-800-265-0761.

<u>To confine spill</u>: If liquid, dike surrounding area or absorb with sand, cat litter or commercial day. If dry material, cover to prevent dispersal. Place damaged package in a holding container, Identify contents.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes candid be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Recional Office for outdance.

Container Handling

Non-refillable container. Do not reuse or refill this container. <u>If empty</u>: Offer for recycling, if available, or dispose of in trash.

If partially filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: To the extent consistent with applicable law, buyer assumes all risks of use, storage or handling of this product not in accordance with label directions.