

FIRST AID	
If in eyes	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Hotline Number
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

In the event of a medical or chemical emergency, contact 
North America 1-800-255-3924 or Worldwide Int'l.
+01-813-248-0585

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, and using the toilet.

ENVIRONMENTAL HAZARDS
To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

DIRECTIONS FOR USE
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

HOW TO APPLY: Scatter the slug and snail bait granules on the soil around or near the plants to be protected. For broadcast application, use standard broadcast spreaders. For row application, use standard granular spreaders. Apply bait evenly at the rate of 0.5-1 lb. per 1000 square feet, or for smaller areas, 0.15 oz., or 1 level teaspoon, per square yard, in a 6 inch circular band around the base of the stem. Apply at 0.15 oz., or 1 level teaspoon, per square yard, in a 6 inch circular band around the base of the plants to be protected. For older trees, spread the bait around the base of the tree to intercept slugs and snails traveling to the trunk. Apply the bait at the rate of 0.5-1 lb. per 1000 square feet for orchards using standard fertilizer granular spreaders.

Reapply as the bait is consumed or at least every two weeks. Apply at higher rate if the infestation is severe, if the area is heavily watered or after long periods of heavy rain. See specific directions for different plant types and for inside residential greenhouses.

Manufactured under a license of W. Neudorff GmbH KG



Manufactured For:
VPG
Valley Packaging
Group, Inc.
230 FM 87 • Bonham, TX 75418
EPA Reg. No. 67702-3-7401
EPA Est. No. 7401-TX-01
Visit Us At: www.fertilome.com
Product Questions? 855-270-4776



WHEN TO APPLY: Apply the bait during the evening, as slugs and snails travel and feed mostly by night or early morning.

WHERE TO APPLY: Treat all likely areas of infestation, especially around the perimeter of garden plots because these pests travel into plant areas from daytime refuges. They favor damp places around vegetable plants such as beans, tomatoes, lettuce, cabbage, celery and squash. Other favorite areas are flower gardens, rockeries, hedges, dichondra lawns, citrus groves, ivy patches, and other ground cover where they obtain shelter by day.

Vegetables

Use the bait to protect any vegetables from slug and snail damage, including (but not limited to): artichokes, asparagus, beans, beets, blackeyed peas, broccoli, Brussels sprouts, cabbage, cantaloupe, carrots, cauliflower, corn, cucumbers, eggplants, garlic, lettuce, onions, peas, peppers, potatoes, radishes, rutabagas, spinach, squash, Swiss chard, tomatoes and turnips. Scatter the bait around the perimeter of the vegetable plot at the rate of 0.5-1 lb. per 1000 square feet, or for smaller areas, 0.15 oz., or 1 level teaspoon, per square yard, to provide a protective "barrier" for slugs and snails entering the garden plot. If slugs or snails are inside the rows, then scatter the bait on the soil around the base of the plants and between the rows.

Fruits Including Citrus

Use the bait to protect fruits from slugs and snails, including (but not limited to): apples, avocados, apricots, cherries, grapes, melons, peaches, plums, citrus, and pears. For seedlings spread the bait around the base of the stem. Apply at 0.15 oz., or 1 level teaspoon, per square yard, in a 6 inch circular band around the base of the plants to be protected. For older trees, spread the bait around the base of the tree to intercept slugs and snails traveling to the trunk. Apply the bait at the rate of 0.5-1 lb. per 1000 square feet for orchards using standard fertilizer granular spreaders.



Slug & Snail Bait

- Kills Snails And Slugs
- Use Around Flowers, Gardens, Vegetables And Fruit Trees
- Remains Effective After Rain Or Sprinkling

ACTIVE INGREDIENT:

By Weight

Iron Phosphate 1.0%

OTHER INGREDIENTS:99.0%

TOTAL: 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

NET WEIGHT 1.5 lbs (0.68 KG)

OPEN
HERE

Berries

Use the bait to protect berries from slugs and snails, including (but not limited to): strawberries, blackberries, blueberries, boysenberries, loganberries, raspberries. Spread the bait around the perimeter of the plot to intercept slugs and snails migrating toward the berries. Use a rate of 0.5-1 lb. per 1000 square feet, or for smaller areas, 0.15 oz., or 1 level teaspoon, per square yard, and scatter by hand or with granular spreaders. If slugs and snails are already in the plots, then carefully spread bait between the furrows near the base of the plants. For small plots, treat around the base of the plants to be protected. Do not spread over the entire area but apply selectively.

Outdoor Ornamentals

Scatter bait in a 6 inch circular band around the base of the plants to be protected at 0.15 oz., or 1 level teaspoon, per square yard. If plants are next to a grassy area, spread the bait between the ornamentals and the grass. Slugs and snails traveling to the plants will encounter the bait before reaching the plant. In these situations, scatter the bait around the perimeter of the plot at the rate of 1 lb. per 1000 square feet to intercept snails and slugs traveling to the plot.

Greenhouses

Where slugs or snails are a problem in greenhouses, scatter the bait in the plant pots of plants being damaged or around pots on greenhouse benches. Apply $\frac{1}{2}$ teaspoon per 9 inch pot.

Lawns

Use the bait on lawns as a buffer zone around gardens. When slugs or snails are detected, scatter the bait at a rate of 0.15 oz., or 1 level teaspoon, per square yard. Scatter by hand or with a granular spreader where the slugs or snails are observed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store this product in its original container and keep in a secure storage area out of reach of children and domestic animals.

CONTAINER DISPOSAL: Nonrefillable container. **If empty:** Do not reuse or refill this container. Offer for recycling, if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY

To the extent consistent with applicable law, the seller warrants that this product conforms to the chemical description on this label and is reasonably fit for purposes stated on this label only when used in accordance with directions for use. To the extent consistent with applicable law this warranty does not extend to use of this product contrary to label directions, or under abnormal use conditions, or under conditions not reasonably foreseeable to seller. To the extent consistent with applicable law, seller makes no other warranties, either expressed or implied."

Slug & Snail Bait

ACTIVE INGREDIENT:	By Weight
Iron Phosphate	1.0%
OTHER INGREDIENTS:.....	99.0%
TOTAL:	100.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION