

FIRST AID

HAVE LABEL WHEN OBTAINING TREATMENT ADVICE

If Swallowed:	Call a poison control center, or doctor, or 1-800-498-5743 immediately for treatment advice.
	Have person sip a glass of water if able to swallow.
	Do not induce vomiting unless told to do so by the poison control center or doctor.
If on Skin or on Clothing:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a price property of the state of the s
	 Call a poison control center, or doctor, or 1-800-498-5743 immediately for treatment advice.
If in Eyes:	Hold eye open and rinse slowly and gently with water for 15-20 minutes.
	Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
	Call a poison control center, or doctor, or 1-800-498-5743 immediately for treatment advice.

NOTE TO PHYSICIAN

If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K₁, intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

TREATMENT FOR PET POISONING

If animal eats bait, call veterinarian or 1-800-498-5743

NOTE TO VETERINARIAN

Anticoagulant Diphacinone: For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K.

For 24-hour emergency information on this product, call 1-800-498-5743 (US & Canada) or 1-651-523-0318 (all other areas).





PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Keep away from children, domestic animals and pets. Wear waterproof gloves made of barrier laminate, polyethylene, butyl rubber (≥14 mils), nitrile rubber (≥14 mils), neoprene rubber (≥14 mils), natural rubber (≥14 mils), polyvinyl chloride (≥14 mils), or Viton (≥14 mils). Any person who retrieves carcasses or unused bait following bait application must wear waterproof gloves.

Wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco, or using the toilet and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to mammals, birds and other wildlife. Dogs, cats and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not allow bait to be exposed on soil surface. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment wash water or rinsate.


DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions, use restrictions, and use precautions.

USE RESTRICTIONS:

- This product may only be applied at the use sites and by the application methods indicated on this label.
- This product may only be used to control moles or pocket gophers in manual, below-ground applications. There are Statespecific limitations on use of this product to control moles.
- Bait must be applied directly into moles' and pocket gophers'
 hurrow systems
- Any person who retrieves carcasses or unused bait following application of this product must wear waterproof gloves.
- Do not apply this product in gardens where food or feed may become contaminated.
- . Do not apply this product by means of an artificial burrow builder.



Fish Flavored, Weather-Resistant Rodenticide

For Control of Pocket Gophers and Moles* In Manual Below-Ground Applications Only.

Intended for Residential Use

Do Not Sell In Individual Containers Containing More Than 1 Pound of Product.

*Not Approved for This Use Pattern in the following states:
California, Hawaii, New Hampshire, and Indiana.

For use only against the eastern mole (Scalopus aquaticus) and the hairy-tailed mole (Parascalops breweri) in North Carolina. Not for use against the star-nosed mole (Condylura cristata parva) in North Carolina.

ACTIVE INGREDIENT:

Diphacinone

(2-Diphenylacetyl-1,3-Indandione)	0.005%
OTHER INGREDIENTS:	<u>99.995%</u>
TOTAL:	100.000%

EPA Reg. No. 61282-65

EPA Est. No. 61282-WI-01

KEEP OUT OF REACH OF CHILDREN

CAUTION





NET CONTENTS: 1 pound (454 grams) NET WEIGHT: 1 pound (454 grams)

MOLES

Not approved for the control of moles in the following States: California, Hawaii, Indiana, and New Hampshire

For use only against the eastern mole (Scalopus aquaticus) and the Hairy-tailed mole (Parascalops breweri) in North Carolina

Not for use against the star-nosed mole (Condylura cristata parva) in North Carolina

Lawns: The presence of moles may be indicated by a network of surface ridges in the turf, by a series of conical mounds of earth pushed up from deep burrows, or by damage to flower hulbs

Surface Ridges: Prior to treatment, determine which tunnels are active by pressing down (flattening) on a small section of the tunnel or by removing a one-inch section of the tunnel's roof. Mark these disturbed sections and recheck them in 24 hours. The tunnel is considered to be active if the flattened runway has been raised up or its roof has been repaired. Only treat active tunnels. Using a long-handled spoon, carefully punch a hole in the top of each active tunnel, insert a teaspoon of bait product, and carefully reclose the hole with sod or a stone. Do not collapse the tunnel again or allow loose soil to cover the bait. Using this method, treat all active tunnels at intervals of 10 to 15 feet. After a few days, check to see if the tunnels still are active by using one of the two methods described above. Retreat the tunnels that remain active.

Conical Mounds: Using a long-handled spoon, carefully remove the center plug from each conical mound. Through the hole just created, drop a teaspoon of bait into the underground tunnel. Close the opening with a rock or sod. Do not allow loose dirt to cover bait. Treat all conical mounds in this manner. If mole activity persists several days after one treatment, additional treatments may be made at intervals of 3 to 4 days.

Flower Bulb Protection (Fall Planted): Pack about onequarter inch of soil against each flower bulb. Place several pellets at sides of and around each bulb. Then, cover the bulbs with soil.

POCKET GOPHERS

Lawns: Burrowing pocket gophers (*Thomomys spp.* and *Geomys spp.*) throw out low, fan-shaped mounds on either side of their underground tunnels. These lateral tunnels coming to the surface are on the flat side of the fan and are plugged with soil. With a long-handled spoon, carefully remove the plug. Insert 4 ounces of this product as far down the lateral tunnel as possible. Reclose the opening, being careful not to cover bait with soil.

FOLLOW-UP (Moles and Pocket Gophers): Wearing waterproof gloves, collect and dispose of all dead, exposed animals and dispose of leftover bait in accordance with the disposal directions on this label. Use detergent and hot water to wash spoons. Do not use spoon for mixing, holding, or transferring food or feed. If reinfestation occurs, repeat treatment.



STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

Pesticide Disposal: Nonrefillable container. Do not refill this container.

If Empty: Place in trash or offer for recycling if available.

If Partially Filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

DISCLAIMER

DISCLAIMER: To the extent permitted by applicable law, seller makes no representation or warranty, either expressed or implied, for results due to misuse, improper handling or improper storage of this material. Nor does Seller assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or improper storage of this material.

For non-emergency (e.g., current product information), Call: 1-800-621-8829

*Not Approved for This Use Pattern in the following states: California, Hawaii, New Hampshire, and Indiana. For use only against the eastern mole (Scalopus aquaticus) and the hairy-tailed mole (Parascalops breweri) in North Carolina. Not for use against the star-nosed mole (Condylura cristata parva) in North Carolina.

Manufactured By: HACCO, Inc. 110 Hopkins Drive Randolph, WI 53956

Ramik® is a Registered Trademark of Neogen Corporation. HACCO, Inc is a wholly subsidiary of the Neogen Corporation.

Code (NE.RA.) NT.RA.