17.5 lb treats up to 6,000 sq ft



GRUB CONTROL + FERTILIZER

APPLY anytime from SPRING through late SUMMER

One application stops/kills grubs all season long

- Kills grubs, European crane fly larvae and mole crickets, controls cutworms & Chinch bugs
- For use on lawns, ground covers, flower beds, and around ornamental trees & shrubs



ACTIVE INGREDIENT:

KEEP OUT OF REACH OF CHILDREN

FOR OUTDOOR RESIDENTIAL USE ON LAWNS AND LANDSCAPED AREAS. For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300

For Medical Emergencies Only, Call (877) 325-1840 NET WEIGHT: 17.5 LBS. (7.93 Kg)





NET WEIGHT: 17.5 LBS. (7.93 Kg)



NET WEIGHT: 17.5 lb (7.93 kg)





LANDSCAPERS GRUB CONTROL + FERTILIZER

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through skin. Causes minimal eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

FIRST AID		
IF ON SKIN OR CLOTHING	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.	
IF IN EYES	 Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a polson control center or doctor for treatment advice. 	
IF SWALLOWED	 Call a polson control center or doctor immediately for treatment advice. Have person sip a glass of water If able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. 	

HOT LINE NUMBER Have the product container or label with you when calling a polson control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information. NOTETO PHYSICIAN No specific antidote is available. Treat the patient symptomatically

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For use on lawns, landscaped areas and plantings around ornamental trees, shrubs, flower and ornamental beds.

Restrictions

DO NOT apply near fish pools, ponds, streams or lakes.

DO NOT use on sod farms.

DO NOT use more than 200 pounds of this product (0.4 lbs a.i.) per acre per year [4.5 pounds of this product (0.09 lbs a.i.) per 1,000 square feet per year].

DO NOT apply this product to potential treated areas that are waterlogged or saturated with water. This product may not work effectively in these times of conditions.

DO NOT harvest or consume fruit or nuts from trees that have been treated within 1 year (365 days).

Keep children and pets off treated area until it has dried.

DO NOT apply this product, by any application method, to linden, basswood or other Tilla species.

Product Information

This product can be used for control of the following soil pests: mole crickets and white grubs, including the larvae of Japanese beetle, Orlental beetle, Chafers, Asian garden beetle, May beetle, June beetle and black turforass ataenlus. This product can also be used to control billbugs, cutworms, Chinch bugs and European crane fly larvae

Turf Grasses

Treat when Insects first appear. Apply this product evenly over the turfgrass area at the use rate indicated in the table below, using a properly calibrated drop or rotary spreader. A more uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. If rainfall has not occurred within 24 hours after application, irrigate treated turfgrass area to move the active ingredient through the thatch and down into the soll. Do not mow lawn until after irrigation or rainfall has occurred so that uniformity of the application will not be affected.

PEST	USE RATE PER 1,000 SQ. FT.	
White grubs, including the larvae of Japanese beetle, Oriental beetle, Chafers, Asian garden beetle, May beetle, June beetle and Black turfgrass ataenlus billbugs, Cutworms, Chinch bugs and European crane fly larvae.	2.87 pounds	
Mole crickets	3.80 pounds	

Flower Beds, Ground Covers, Ornamental Trees and Shrubs

Application Instructions

Apply this product at the rate of 2 tablespoons per 10 square feet of planting bed (1-1/4 cups per 100 square feet of planting bed). Evenly spread granules to the soil over flower beds, ground cover plantings, shrub beds and under ornamental trees. The use of a hand-held spreader is recommended. Rake granules Into the soil around plants. Water thoroughly after application. This product works best Wilen applied before mole crickets and grubs hatch. If necessary, consult your local State extension agent for the ideal times to apply this product in your area.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep out of reach of children and animals. Store in original container only. Store in a cool, dry place

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. DO NOT reuse or refill this container. If empty: Place In trash or offer for recycling If available. If partly filled: Call your local solid waste agency for disposal Instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS

Information regarding the contents and level of metals in this product is DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

GUARANTEED ANALYSIS

20-0-4 20.0%

TOTAL NITROGEN (N) 20.0% Urea Nitrogen

SOLUBLE POTASH (K2O)...

DERIVED FROM: Polymer Sulfur-Coated Urea, Urea, Muriate of Potash. CHLORINE (CI) MAXIMUM

*7.2% Coated Slow Release Nitrogen from Polymer Sulfur-Coated Urea

..... 3.0% F1612

Environmental Hazards

This product is highly toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off Into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems

Spreader Settings

Broadcast Spreaders: For best results, apply in a circular path starting on the outside and working toward the center. A little overlap of product will ensure complete coverage. Drop Spreaders: First, apply 2 border strips along the edge of each end of the lawn. Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips.

Applications Instructions

See Spreader Settings chart below. Adjust spreader to the appropriate setting. Make sure that the spreader works properly and spreads granules uniformly. Apply to the border and then fill in the center to ensure the entire area is covered completely. To make sure that the product is evenly distributed, walk at a steady pace making parallel passes. DO NOT use a spreader that applies granules in narrow, concentrated or uneven bands. Water thoroughly after application

Important: It is important to follow the directions as indicated on this quality lawn care product as over-application may cause injury to the lawn. Measure the size of the area to be treated and do not over or under apply. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. Calibrate spreader prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing uniformly. Close spreader when filling, stopping or turning. DO NOT

Broadcast/Rotary Spreaders	2.87 lbs./ 1,000 sq. ft.	3.8 lbs./1,000 sq. ft.
Earthway® Rotary	14	15
Scotts® EdgeGuard®	4	5
Scotts® SpeedyGreen®	4	5
Lowes 32 lb Broadcast	4	5

Typically, a lawn utilizes 1 to 4 lbs of nitrogen per 1,000 sq ft per growing season. When applied as directed, this product provides 0.57 to 0.76 lb of nitrogen per 1,000 sq ft. Contact your local Extension Service for information specific to your geographic region.

PESTS CONTROLLED	WHEN TO APPLY	INDICATION OF DAMAGE	USE RATE PER 1,000 SQ. FT
Grubs	Apply once anytime from May through mid-August damage to lawns by feeding grass roots, causing it to turn then brown and die.		2.87 pounds
European crane fly larvae	Apply once, when these insects are present in large numbers.	European crane fly larvae cause damage by feeding on grass roots and disturbing the soil surface,	2.87 pounds
Mole Crickets	Apply once anytime from mid-May through mid-June. Florida residents - Apply once anytime between early April through early May.	Mole crickets cause widespread damage by tunneling through the soil, while some species also feed on roots. Tunneling through the soil uproots the grass causing it to dry out, become thin and weedy, turn yellow and eventually die.	

Application Instructions

- . Apply early before grubs, mole crickets and European crane fly larvae hatch.
- Irrigate thoroughly within 24 hours of application to form a protective zone in the soil. This product can be applied over soil or mulch. Remove black plastic or other weed barrier material prior to application.
- Application to water saturated soils may decrease product effectiveness.
- Restrictions:
- Keep children and pets off treated area until it has dried. DO NOT apply to water saturated areas.
- DO NOT move turf or lawn area until after sufficient irrigation or rainfall has occurred so that uniformly of application will not be affected.

 DO NOT use more than 200 pounds of this product (0.4 lbs a.i.) per acre per year [4.5 pounds of this product (0.091bs a.i.) per 1,000 square feet per year].

available on the internet at http://www.aapfco.org/metals.htm.

EPA REG. NO. 228-510-82757, EPA EST. No. 82757-FL-001 (S), 82757-MA-002(H), 82757-0H-001 (MF)

First letter of lot code on end of container indicates establishment.

Manufactured and Guaranteed by:

Turf Care Supply Corp., 50 Pearl Road, Suite 200, Brunswick, OH 44212 877-220-1306

MADE IN U.S.A.

(RV103114)



NET WEIGHT: 17.5 LBS. (7.93 Kg)

GRUB CONTROL + FERTILIZER

RS

NDSCAPE

NET WEIGHT: 49 lb (22.22 kg)



GRUB CONTROL + FERTILIZER

APPLY anytime from SPRING through late SUMMER

One application stops/kills grubs all season long

- Kills grubs, European crane fly larvae and mole crickets, controls cutworms & Chinch bugs
- For use on lawns, ground covers, flower beds, and around ornamental trees & shrubs

20 0 4



ACTIVE INGREDIENT:

Imidacloprid
OTHER INGREDIEN

99.8%

0.2% KEEP OUT OF REACH OF CHILDREN

100.0% CAUTION SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

FOR OUTDOOR RESIDENTIAL USE ON LAWNS AND LANDSCAPED AREAS.

RESTRICTION: Not for sale, use, or distribution in or into Nassau, Suffolk, Kings or Queens Counties, New York. For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300 For Medical Emergencies Only, Call (877) 325-1840

NET WEIGHT: 49 LBS. (22.22 Kg)

49 lb treats up to 17,000 sq ft



GRUB CONTROL + FERTILIZER NET WEIGHT: 49 LBS. (22.22 Kg)







LANDSCAPERS GRUB CONTROL + FERTILIZER

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through skin. Causes minimal eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

*7.2% Coated Slow Release Nitrogen from Polymer Sulfur-Coated Urea

DERIVED FROM: Polymer Sulfur-Coated Urea, Urea, Muriate of Potash.

FIF	STAID	
	IF ON SKIN OR CLOTHING	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
	IF IN EYES	Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, If present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
	IF SWALLOWED	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment You may also contact 1-877-325-1840 for emergency medical treatment information. NOTE TO PHYSICIAN No specific antidote is available. Treat the patient symptomatically

Environmental Hazards

CHLORINE (CI) MAXIMUM ..

TOTAL NITROGEN (N) SOLUBLE POTASH (K20)

GUARANTEED ANALYSIS

This product is highly toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

20-0-4 20.0%

4.0%

3.0%

F1612

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
For use on lawns, landscaped areas and plantings around ornamental trees, shrubs, flower and ornamental beds.

DO NOT apply near fish pools, ponds, streams or lakes.

DO NOT use on sod farms.

DO NOT use more than 200 pounds of this product (0.4 lbs a.i.) per acre per year [4.5 pounds of this product (0.09 lbs a.i.) per 1,000 square feet per year].

DO NOT apply this product to potential treated areas that are waterlogged or saturated with water. This product may not work effectively in these times of conditions.

DO NOT harvest or consume fruit or nuts from trees that have been treated within 1 year (365 days).

Keep children and pets off treated area until it has dried. DO NOT apply this product, by any application method, to linden, basswood or other Tilia species.

Product Information

This product can be used for control of the following soil pests: mole crickets and white grubs, including the larvae of Japanese beetle, Oriental beetle, Chafers, Asian garden beetle, May beetle, June beetle and black turfgrass ataenius. This product can also be used to control billbugs, cutworms, Chinch bugs and European crane fly larvae.

Turf Grasses

Treat when insects first appear. Apply this product evenly over the turfgrass area at the use rate indicated in the table below, using a properly calibrated drop or rotary spreader. A more uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. If rainfall has not occurred within 24 hours after application, irrigate treated turfgrass area to move the active ingredient through the thatch and down into the soil. Do not mow lawn until after imgation or rainfall has occurred so that uniformity of the application will not be affected.

PEST	USE RATE PER 1,000 SQ. FT.	
White grubs, including the larvae of Japanese beetle, Oriental beetle, Chafers, Asian garden beetle, May beetle, June beetle and Black turfgrass ataenius billbugs, Cutworms, Chinch bugs and European crane fly larvae.	2.87 pounds	
Mole crickets	3.80 pounds	

Flower Beds, Ground Covers, Ornamental Trees and Shrubs

Apply this product at the rate of 2 tablespoons per 10 square feet of planting bed (1-1/4 cups per 100 square feet of planting bed). Evenly spread granules to the soil over flower beds, ground cover plantings, shrub beds and unde ornamental trees. The use of a hand-held spreader is recommended. Rake granules into the soil around plants. Water thoroughly after application. This product works best Wilen applied before mole crickets and grubs hatch. If necessary consult your local State extension agent for the ideal times to apply this product in your area.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal. **PESTICIDE STORAGE:** Keep out of reach of children and animals. Store in original container only. Store in a cool, dry place

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. DO NOT reuse or refill this container If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER, THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER. MANUFACTURER AND SELLER. AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER INCLUDING USE, APPLICATION, HANDLING, AND DISPOSALTO I HEAZIENT CONSISTENT WITH APPLICABLE LAW, MANDFALTOREN
OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY,
OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR RIS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

Spreader Settings

Broadcast Spreaders: For best results, apply in a circular path starting on the outside and working toward the center. A little overlap of product will ensure complete coverage. Drop Spreaders: First, apply 2 border strips along the edge of each end of the lawn. Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips.

Applications Instructions

See Spreader Settings chart below. Adjust spreader to the appropriate setting. Make sure that the spreader works properly and spreads granules uniformly. Apply to the border and then fill in the center to ensure the entire area is covered completely. To make sure that the product is evenly distributed, walk at a steady pace making parallel passes. DO NOT use a spreader that applies granules in narrow, concentrated or uneven bands. Water thoroughly

Important: It is important to follow the directions as indicated on this quality lawn care product as over-application may cause injury to the lawn. Measure the size of the area to be treated and do not over or under apply. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. Calibrate spreader prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing uniformly. Close spreader when filling, stopping or turning. DO NOT hand spread

Broadcast/Rotary Spreaders	2.87 lbs./ 1,000 sq. ft.	3.8 lbs./1,000 sq. ft.
Earthway® Rotary	14	15
Scotts® EdgeGuard®	4	5
Scotts® SpeedyGreen®	4	5
Lowes 32 lb Broadcast	4	5

Typically, a lawn utilizes 1 to 4 lbs of nitrogen per 1,000 sq ft per growing season. When applied as directed, this product provides 0.57 to 0.76 lb of nitrogen per 1,000 sq ft. Contact your local Extension Service for information

specific to your geographic region.			
PESTS CONTROLLED	WHEN TO APPLY	INDICATION OF DAMAGE	USE RATE PER 1,000 SQ. FT.
Grubs	Apply once anytime from May through mid-August	White grubs cause widespread damage to lawns by feeding on grass roots, causing it to turn yellow, then brown and die.	2.87 pounds
European crane fly larvae	Apply once, when these insects are present in large numbers.	European crane fly larvae cause damage by feeding on grass roots and disturbing the soil surface.	2.87 pounds
Mole Crickets	Apply once anytime from mid-May through mid-June. Florida residents - Apply once anytime between early April through early May.	Mole crickets cause widespread damage by tunneling through the soil, while some species also feed on roots. Tunneling through the soil uproots the grass causing it to dry out, become thin and weedy, turn yellow and eventually results.	3.80 pounds

phication Instructions

- Apply early before grubs, mole crickets and European crane fly larvae hatch.
- Imigate thoroughly within 24 hours of application to form a protective zone in the soil This product can be applied over soil or mulch. Remove black plastic or other weed barrier material prior to application
- Application to water saturated soils may decrease product effectiveness. Restrictions:
- Keep children and pets off treated area until it has dried.

 DO NOT apply to water saturated areas.
- DO NOT mow turf or lawn area until after sufficient irrigation or rainfall has occurred so that uniformity of application will not be affected.
- DO NOT use more than 200 pounds of this product (0.4 lbs a i.) per acre per year [4.5 pounds of this product (0.091bs a.i.)

per 1,000 square feet per year].

Information regarding the contents and level of metals in this product is available on the internet at http://www.aapfco.org/metals.htm EPA REG. NO. 228-510-82757, EPA EST. No. 82757-FL-001 (S),

82757-MA-002(H), 82757-0H-001 (MF) First letter of lot code on end of container indicates establishment.

Manufactured and Guaranteed by: Turf Care Supply Corp., 50 Pearl Road, Suite 200, Brunswick, OH 44212

MADE IN U.S.A.

(RV103114)

