MICRODOT

Covers up to 15,000 sq ft

LANDSCAPERS™ SELECT



FERTILIZER WITH CRAB-BUSTER® CRABGRASS PREVENTER

For control of crabgrass and control or suppression of many other listed annual grass and broadleaf weeds in established lawns and ornamental turf.

1st FIRST APPLICATION ~ Apply Early Spring ~

LANDSCAPERS™ SELECT

FERTILIZER WITH CR

NET WT 40.5 lb (18.37 kg)

RABGRASS PREVENTER

ZER WITH CRAB-BUSTER® CRABGRASS PREVENTER

.37 kg)

CRABGRASS PREVENTER

For control of crabgrass and control or suppression of many other listed annual grass and broadleaf weeds in established lawns and ornamental turf.

- **Contains Crab-Buster® Herbicide**
  - A highly effective herbicide for crabgrass control!
- **Premium Granular Fertilizer**
  - Providing the nutrients needed for a beautiful healthy lawn.

23	0	4

Phosphorus - Free

PREMIUM GRANULAR FERTILIZER

1st

FIRST APPLICATION

~ Apply Early Spring ~

1

FIRST A

~ Apply

FERTILIZER WITH CRAB-BUSTER® CRABGRASS PR

NET WT 40.5 lb (18.37 kg)

APERS

PREVENTER

1st FIRST APPLICATION ~ Apply Early Spring ~



ACTIVE INGREDIENT:

Dithiopyr: 3,5- pyridinedicarbothioic acid, 2-(difluoromethyl)- 4-(2-methylpropyl)-6-(trifluoromethyl)-S,S-dimethyl ester 0.15%

OTHER INGREDIENTS: 99.85%

TOTAL: 100.00%

This product contains 0.075 pound of the active ingredient dithiopyr per 50 pound bag.

KEEP OUT OF REACH OF CHILDREN

CAUTION REFER TO BACK PANEL OF BAG FOR DIRECTIONS FOR USE INCLUDING STORAGE AND DISPOSAL.

NET WT 40.5 lb (18.37 kg)

FERTILIZE

NET WT 40.5 lb (18.37 kg)

LANDSCAPERS SELECT

MICRODOT

Covers up to 15,000 sq ft

LANDSCAPERS SELECT

FERTILIZER WITH CRAB-BUSTER CRABGRASS PREVENTER

1st FIRST APPLICATION ~ Apply Early Spring ~

NET WT 40.5 lb (18.37 kg)

LANDSCAPERS SELECT

FERTILIZER WITH CRAB-BUSTER® CRABGRASS PREVENTER

- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

NOTICE: Read the entire label. Use only according to label directions.
Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feed, drugs or clothing.

GUARANTEED ANALYSIS 23-0-4

TOTAL NITROGEN (N)	23.0%
23.0% Urea Nitrogen*	
SOLUBLE POTASH (K_2O)	4.0%
DERIVED FROM: Polymer Sulfur-Coated Urea, Urea, Muriate of Potash.	
CHLORINE (CI) MAX	3.0%

*7.2% Slowly Available Nitrogen from Polymer Sulfur-Coated Urea

F1612

PRECAUTIONARY STATEMENTS

First Aid	
IF IN EYES	• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF INHALED	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

KEEP OUT OF REACH OF CHILDREN

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUSES EYE IRRITATION - HARMFUL IF INHALED

Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

Environmental Hazards

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn will help to prevent runoff to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

In the state of New York, do not exceed 3.8 lb per 1000 sq ft per year of this product (equivalent to 0.25 lb of active ingredient per acre). In the state of New York, this product may be applied at a minimum of 2.0 lb per 1000 sq ft per year (equivalent to 0.128 lb active ingredient per acre).

In the state of Florida, do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn.

Use Directions For Turf

This turf fertilizer product, containing Crab-Buster® herbicide, can be applied up to 4 weeks later than other crabgrass preventers and provides up to 4 months of crabgrass control.

Landscapers Select™ Fertilizer with Crab-Buster® Crabgrass Preventer provides preemergence control of crabgrass and control or suppression of other listed weeds when applied prior to their germination. In addition, this product provides postemergence control of seedling crabgrass after germination and emergence. Postemergence control is limited to crabgrass only, but is effective up to 4 weeks after crabgrass has germinated. This generally corresponds to the time when crabgrass seedlings first become visible in the established turfgrass. Do not apply this product later than 4 weeks after crabgrass germination.

Florida Applications: Do not exceed 2 applications per year on centipedegrass in the north. Do not apply more than 7.7 lb of this product per 1,000 sq ft per application and no more than 23.1 lb of this product/1000 sq ft per year (equivalent to 0.5 and 1.5 lb of active ingredient per acre, respectively).

Use Precautions For Turf

This product should be applied only to lawns and ornamental turfgrasses that are well established. For newly established lawns, initial application of this product should be made only

Apply this product only to turfgrass species listed on this label. The following turfgrass species are tolerant to this product when applied as directed:	
Cool-Season Grasses	**Warm-Season Grasses**
	Bahiagrass

Weeds Controlled Or Suppressed

Landscapers Select™ Fertilizer with Crab-Buster® Crabgrass Preventer will provide control of crabgrass and control or suppression of other listed grass and broadleaf weeds when applied prior to their germination.

Grasses		Broadleaf Weeds			
barley	goosegrass	bittercress†	common†	parsley-piert†	spurge, prostrate
barnyardgrass	foxtail, green	carpetweed	marestail	purslane, common	spurge, spotted
bluegrass, annual	foxtail, yellow	chickweed†	medic, black	rocket, London	woodsorrel, creeping
brome	kikuyugrass†	geranium, Carolina†	mustard	shepherdspurse	woodsorrel, yellow
crabgrass, large	oats, wild	henbit	oxalis, buttercup	speedwell, corn†	
crabgrass, smooth	ryegrass	knotweed, prostrate	pineappleweed	spurge, garden	
crabgrass, Southern	(annual & perennial)	lespedeza,	pigweed, redroot		
crowfootgrass†	sandbur				
dallisgrass	smutgrass				

† Suppression only.

Weed Resistance Management

Dithiopyr, the active ingredient in this product, is a Group 3 herbicide based on the mode of action classification system of the Weed Science Society of America. Any weed population may contain or develop plants resistant to Group 3 herbicides. Resistant weeds may dominate the weed population if these herbicides are used repeatedly in the same field. Such resistant weed plants may not be effectively managed using Group 3 herbicides but may be effectively managed utilizing other herbicides alone or in mixtures from a different herbicide Groups that are labeled for control of these weeds and/or by using cultural or mechanical practices. However, a herbicide mode of action classification by itself may not adequately address specific weeds that are resistant to specific herbicides. Consult your local company representative, state cooperative extension service, professional consultants or other qualified authorities to determine appropriate actions for treating specific resistant weeds

after crabgrass has germinated. This generally corresponds to the time when crabgrass seedlings first become visible in the established turfgrass. Do not apply this product later than 4 weeks after crabgrass germination.

Florida Applications: Do not exceed 2 applications per year on centipedegrass in the north. Do not apply more than 7.7 lb of this product per 1,000 sq ft per application and no more than 23.1 lb of this product/1000 sq ft per year (equivalent to 0.5 and 1.5 lb of active ingredient per acre, respectively).

Use Precautions For Turf

This product should be applied only to lawns and ornamental turfgrasses that are well established. For newly established lawns, initial applicatio n of this product should be made only after turfgrasses have developed a vigorous root system, uniform stand, and received at least two mowings following seeding, sodding, or sprigging. **Exception**: In newly established lawns of Kentucky bluegrass, perennial ryegrass and/or tall fescue, this product may be applied two weeks after the first sign of germination.

Apply this product only to turfgrass species listed on this label. The following turfgrass species are tolerant to this product when applied as directed:

Cool-Season Grasses	Warm-Season Grasses
Bentgrass, creeping Bluegrass, Kentucky Fescue, fine Fescue, tall Ryegrass, perennial	Bahiagrass Bermudagrass Buffalograss Carpetgrass Centipedegrass St. Augustinegrass Zoysiagrass

- Use of this product on lawns and ornamental turfgrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperatures, or lack of moisture may result in turfgrass injury.
- For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, should be done before application of this product. **Do not** apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.
- **Reseeding, Overseeding, or Sprigging:** Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application program totaling 7.7 lb/ 1,000 sq ft (0.5 lb active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses. When reseeding, overseeding follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.

Use Precautions

- **Do not** use clippings from treated turf for mulching around vegetables or fruit trees.
- **Do not** allow people or pets to enter treated area until dust has settled.

Application Timing

Spring Application

Control of Crabgrass: **Landscapers Select™ Fertilizer with Crab-Buster® Crabgrass Preventer** provides preemergence control of crabgrass and control or suppression of other listed weeds when applied prior to their germination. In addition, this product provides postemergence control of seedling crabgrass after germination and emergence. Postemergence control is limited to crabgrass only and is effective up to 4 weeks after germination of crabgrass. This generally corresponds to the time when crabgrass seedlings are at the 3-leaf stage of growth and first become visible in the established turf. This product will not control crabgrass at later growth stages when crabgrass plants begin to tiller (spread). Properly timed applications will control existing crabgrass and prevent further emergence of crabgrass seedlings. Do not apply this product later than 4 weeks after crabgrass germination.

Control or Suppression of Other Grass and Broadleaf Weeds: For control or suppression of other grass and broadleaf weeds, this product must be applied as a **preventative** treatment prior to their germination and emergence. Only crabgrass will be controlled with applications made up to 4 weeks after weed germination.

Fall Application

This product can also be applied from late summer through early fall (late August through November). When applied in the fall, a sequential or second application of this product must be made the following spring to provide season-long control of crabgrass. If this product is applied in the fall, **do not** reseed, overseed or sprig turfgrasses within 12 weeks after application.

Tips for Improved Weed Control

- For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.
- The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by 1/2 inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.
Pesticide Storage: Store this product only in its original container in a dry, cool, secured storage area.
Container Handling: Nonrefillable container. Do not reuse or refill this container.
If empty: Place in trash or offer for recycling if available.
If partially filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions.
Never place unused product down any indoor or outdoor drain.

dallisgrass	smutgrass

† Suppression only.

Weed Resistance Management

Dithiopyr, the active ingredient in this product, is a Group 3 herbicide based on the mode of action classification system of the Weed Science Society of America. Any weed population may contain or develop plants resistant to Group 3 herbicides. Resistant weeds may dominate the weed population if these herbicides are used repeatedly in the same field. Such resistant weed plants may not be effectively managed using Group 3 herbicides but may be effectively managed utilizing other herbicides alone or in mixtures from a different herbicide Groups that are labeled for control of these weeds and/or by using cultural or mechanical practices. However, a herbicide mode of action classification by itself may not adequately address specific weeds that are resistant to specific herbicides. Consult your local company representative, state cooperative extension service, professional consultants or other qualified authorities to determine appropriate actions for treating specific resistant weeds.


Best Management Practices

Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is recommended. A diversified weed management program may include the use of multiple herbicides with different modes of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistant weeds. Scouting after a herbicide application is important because it can facilitate the early identification of weed shifts and/or weed resistance and thus provide direction on future weed management practices. One of the best ways to contain resistant weed populations is to implement measures to avoid allowing weeds to reproduce by seed or to proliferate vegetatively. Cleaning equipment between sites and avoiding movement of plant material between sites will greatly aid in reducing the spread of resistant weed seed.

Length Of Control

Landscapers Select™ Fertilizer with Crab-Buster® Crabgrass Preventer will provide up to 4 months of crabgrass control when applied according to label directions. The actual length of control will vary depending on use rate, weather conditions, condition of the turfgrass, intensity of weed pressure, and the location or "Region" within the United States where this product is being applied. **Refer to the map below to determine the region of the U.S. that includes your location.** Use the table below to determine the amount of **Landscapers Select™ Fertilizer with Crab-Buster® Crabgrass Preventer** to apply in your location based on the desired length of weed control or need for postemergence crabgrass control.

Use "Regions" within the United States:



Use Rates:

Amount of Landscapers Select™ Fertilizer with Crab-Buster® Crabgrass Preventer to Apply (Pounds of product to apply per 1,000 square feet of turfgrass †)

Region of U.S.	Length of Crabgrass Control		
	2-3 months	3-4 months	Early Postemergence Control
North	2.0 lb/1000 sq ft (0.128 lb ai/acre)	2.7 lb/1000 sq ft (0.174 lb ai/acre)	
South	2.9 lb/1000 sq ft (0.192 lb ai/acre)	3.4 - 3.9 lb/1000 sq ft †† (0.22 - 0.256 lb ai/acre)	3.4 - 7.7 lb/1000 sq ft †† (0.22 - 0.5 lb ai/acre)

† 1,000 square feet equals an area 100 feet x 10 feet or square area of about 32 feet on a side.

†† Use the higher rate in rate range in areas of high weed pressure or for increased length of control.

Maximum Application Rates:

- **Do not** apply more than 7.7 lb of this product per 1,000 sq ft per application. Do not apply more than 23.1 lb of this product/1000 sq ft per year (equivalent to 0.5 and 1.5 lb of active ingredient per acre, respectively).
- In the state of New York, do not exceed 3.8 lbs per 1,000 sq ft per year of this product (equivalent to 0.25 lb of active ingredient per acre).

Application Instructions and Equipment

This product may be applied with drop or rotary-type spreaders designed to apply granular herbicides. For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps. Avoid the use of spreaders that tend to apply granules in narrow rows or concentrated bands. Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the application rate under actual application conditions. The desired calibration setting may be marked or recorded for future reference. Apply this product uniformly over the treatment area. More uniformity of application can usually be achieved by applying one-half of the required amount of product over the treatment area and then applying the remaining one-half in a different direction (e.g., at a right angle to the previous direction). Avoid streaking, skips, or overlaps during application. Check equipment frequently to verify calibration and proper functioning.

Spreader	Calibration Settings at Specified Rate (lb/1000 sq ft)		
Brand and Model of Spreader	**2.7**	**3.4**	**3.9**
Cyclone Rotary	4	4-1/2	5
Earthway® Rotary	14	15	16
STA-Green® Rotary (1-10), Vigoro® Rotary (1-10)	3-1/2	4	4-1/2
Scotts® EasyGreen Rotary	21	23	25
Scotts® SpeedyGreen®, EdgeGuard®, EdgeGuard Mini®, EdgeGuard Deluxe®, Rotary	4	4-1/2	5

Application Timing

Spring Application

Control of Crabgrass: **Landscapers Select™ Fertilizer with Crab-Buster® Crabgrass Preventer** provides preemergence control of crabgrass and control or suppression of other listed weeds when applied prior to their germination. In addition, this product provides postemergence control of seedling crabgrass after germination and emergence. Postemergence control is limited to crabgrass only and is effective up to 4 weeks after germination of crabgrass. This generally corresponds to the time when crabgrass seedlings are at the 3-leaf stage of growth and first become visible in the established turf. This product will not control crabgrass at later growth stages when crabgrass plants begin to tiller (spread). Properly timed applications will control existing crabgrass and prevent further emergence of crabgrass seedlings. Do not apply this product later than 4 weeks after crabgrass germination.

Control or Suppression of Other Grass and Broadleaf Weeds: For control or suppression of other grass and broad-leaf weeds, this product must be applied as a **preventative** treatment prior to their germination and emergence. Only crabgrass will be controlled with applications made up to 4 weeks after weed germination.

Fall Application


This product can also be applied from late summer through early fall (late August through November). When applied in the fall, a sequential or second application of this product must be made the following spring to provide season-long control of crabgrass. If this product is applied in the fall, **do not** reseed, overseed or sprig turfgrasses within 12 weeks after application.

Tips for Improved Weed Control

- For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.
- The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by 1/2 inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.
Pesticide Storage: Store this product only in its original container in a dry, cool, secured storage area.
Container Handling: Nonrefillable container. Do not reuse or refill this container.
If empty: Place in trash or offer for recycling if available.
If partially filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions.
Never place unused product down any indoor or outdoor drain.



	2-3 months	3-4 months	Early Postemergence Control
North	2.0 lb/1000 sq ft (0.128 lb ai/acre)	2.7 lb/1000 sq ft (0.174 lb ai/acre)	
South	2.9 lb/1000 sq ft (0.192 lb ai/acre)	3.4 - 3.9 lb/1000 sq ft †† (0.22 - 0.256 lb ai/acre)	3.4 - 7.7 lb/1000 sq ft †† (0.22 - 0.5 lb ai/acre)

† 1,000 square feet equals an area 100 feet x 10 feet or square area of about 32 feet on a side.

†† Use the higher rate in rate range in areas of high weed pressure or for increased length of control.

Maximum Application Rates:

- **Do not** apply more than 7.7 lb of this product per 1,000 sq ft per application. Do not apply more than 23.1 lb of this product/1000 sq ft per year (equivalent to 0.5 and 1.5 lb of active ingredient per acre, respectively).
- In the state of New York, do not exceed 3.8 lbs per 1,000 sq ft per year of this product (equivalent to 0.25 lb of active ingredient per acre).

Application Instructions and Equipment

This product may be applied with drop or rotary-type spreaders designed to apply granular herbicides. For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps. Avoid the use of spreaders that tend to apply granules in narrow rows or concentrated bands. Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the application rate under actual application conditions. The desired calibration setting may be marked or recorded for future reference. Apply this product uniformly over the treatment area. More uniformity of application can usually be achieved by applying one-half of the required amount of product over the treatment area and then applying the remaining one-half in a different direction (e.g., at a right angle to the previous direction). Avoid streaking, skips, or overlaps during application. Check equipment frequently to verify calibration and proper functioning.

Spreader	Calibration Settings at Specified Rate (lb/1000 sq ft)		
Brand and Model of Spreader	**2.7**	**3.4**	**3.9**
Cyclone Rotary	4	4-1/2	5
Earthway® Rotary	14	15	16
STA-Green® Rotary (1-10), Vigoro® Rotary (1-10)	3-1/2	4	4-1/2
Scotts® EasyGreen Rotary	21	23	25
Scotts® SpeedyGreen®, EdgeGuard®, EdgeGuard Mini®, EdgeGuard Deluxe®, Rotary	4	4-1/2	5

Terms and Conditions of Use - If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer - Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use - It is impossible to eliminate all risks associated with the use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent permitted by law, all such risk shall be assumed by buyer.

Limitation of Remedies - To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at the Seller's elections, one of the following:

(1) Refund of purchase price paid by the buyer or user for product bought, or (2) Replacement of amount of product used.

To the extent permitted by law, Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

If you have questions or comments regarding the quality of this product, please write Turf Care Supply Corp. at the address below, or call Customer Service at 877-220-1306 between the hours of 9 A.M. And 5 P.M., Eastern Time, Monday through Friday. Please refer to the item number found on this bag when writing or calling.

Information regarding the contents and levels of metals in this product is available at the following website:
http://www.aapfco.org/metals.htm

EPA REG. NO. 62719-504-82757
EPA EST. NO. 82757-FL-001 (S), 82757-MA-002 (H), 82757-OH-001 (MF)
First letter of lot code on end of container indicates establishment.

Manufactured and Guaranteed by:
Turf Care Supply Corp., 50 Pearl Road, Suite 200, Brunswick, OH 44212

Produced for:
Dow AgroSciences LLC, 9330 Zionsville Road, Indianapolis, IN 46268

Made in U.S.A.

Crab-Buster® is a trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow.

Item # 7615073

8 50526 00276 3

P7615073

NET WT 40.5 lb (18.37 kg)