



PEEL HERE





SHAKE WELL BEFORE USE. READ ENTIRE LABEL BEFORE EACH USE

Γ		FIRST AID			
I	F ON SKIN OR LOTHING:	Take off contaminated clothi Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center of the state			

- doctor for treatment advice.
- Immediately call a poison control center or doctor.
 Do not induce vomiting unless told to do so by a poison contro center or doctor.

 Do not give any liquid to the person.

 Do not give anything by mouth to an unconscious person.
- Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

 For information regarding medical emergencies For information regarding medical emergencie or pesticide incidents, call 1-888-740-8712.
- EPA Reg. No. 1021-1758-89609 EPA Est. No. 33595-MO-2 (Y), 33595-MO-3 (Q), 33595-MO-4 (N), 11623-GA-1 (A), 11623-GA-2 (B)

- Kills gnats, fleas, deer ticks, brown dog ticks and other listed crawling insects
- Kills listed bugs on lawns, trees, shrubs, roses and flowers.

ADVECTA™ YARD for Flens Ticks & Mosquitoes Yard ADVECTA**—INAD TRUSS, IRAS & MOSQUINESS trut Spray Concentrate contains Permethrin, a long lasting insecticide that provides contact and residual control of many hard-to-kill listed sucking and chewing insects for up to 4 weeks.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

For consumer questions call: 1-888-846-4231

Distributed by: TruRx LLC. 500 E Shore Drive, Suite 130, Eagle, ID 83616



PESTS

Ants, Armyworms, Brown dog ticks, Chinch bugs, Crickets, Cutworms, Deer ticks, Fleas, Grasshoppers, Japanese beetle grubs, Mole crickets and Sod webworms.

HOW TO USE

Arizona Cypress, Azalea, Birch, Non-Bearing Citrus Conifers, Elm, English Iv Mock-Orange, Oak, Palm

Kentucky bluegrass, Bent, Bermuda, Blueg Dichondra, Fescue, Irish moss, Merion and St. Augustine

Aphids, Bagworms, Cicadas, Exposed thrips, Fall eafminers, Mealybugs, pider mites, Whiteflies Fall webworms on Cherry and Han weavourins on Conifer, Elm leaf beetles, Elm spanworms and Fall webworms on Elms, Tussock moths on Douglas and Grand Firs, Citrus black flies on Non-Bearing Citrus, Oakworm: on Oaks, Pine beetles, Pine moths, Pine needleminers, and Needle scales on Pines.

Apply as a thorough spray, wetting leaves and branches but do not let runoff occur. Try to penetrate dense foliage. Spray in the late ofternoon or evening, when the termount of evening when the termore armore from 50 to 75°F and when there is little or no wind. Spray at the first sign of insects.

Repeat as necessary; use intervals of 4-8 days.

HOW TO USE

Thoroughly wet down grass a few hours before applying. Home lawns should be no taller than 3 inches at time of application. Slightly more water may be used as long as the amount of product per area is as listed below. For heavy infestations, repeat application after 2 weeks. For listed insects 32 fl. oz. treats 5,300 sq. ft.

PESTS HOW TO USE

Aaeratum, Aster, Azalea, Begonia, Coleus, Common Ninebark and Snowberry, Exacum, Gladiolus, Gold Bells Hypoestes, Ivy, Lilac, Marigold, Mock-Orange, Nannyberry, Orchid, Pansy, Pea Shrub, Petunia, Poinsettia, Rhododendron, Rose, Snapdraaon, Zinnio

Aphids, Bagworms, Cicadas, Exposed thrips, Fall cankerworms, Fall webworms, Inchworms, Leafminers, Leafrollers, Mealybugs, Spider mites and Whiteflies.

Apply as a thorough spray, wetning the plants thoroughly, but do not let runoff occur. Try to hit underside of leaves and penetrate dense foliage. Spray in the later afternoon or evening, when the temperature ranges from 50 to 75°F and when there is little or no wind. Spray at the first sign of insects. To maintain effectiveness; spray as reinfestation occurs but use intervals of 4-8 days.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soop and water after handling and before eating, drinking, chewing gum, or using tobacco.
Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic organisms, including fish and invertebrates.

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging in the treatment area.

PHYSICAL OR CHEMICAL HAZARDS Do not apply this product around electrical equipment due to possible shock hazard.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE
Store in a cool, dry area, preferably a locked
storage area. Store above freezing. Do not
store diluted spray. Store with container
tightly closed. Always store pesticides in the
original container, do not transfer to food or

drink container. Avoid contamination of food, feed, and domestic water supplies. Store away from food and pet food.

PESTICIDE DISPOSAL AND CONTAINER HANDLIN Nonrefillable container. Do not reuse or refi

this container.
If empty: Place in trash or offer for recycling if available.
If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Distributed by: TruRx LLC. 500 E Shore Drive, Suite 130, Eagle, ID 83616

EPA Reg. No. 1021-1758-89609 EPA Est. No. 33595-MO-2 (Y), 33595-MO-3 (Q), 33595-MO-4 (N), 11623-GA-1 (A), 11623-GA-2 (B)

PRODUCT FACTS		
KILLS INSECTS	Lawn pests: ants, armyworms, brown dog ticks, chinch bugs, crickets, cutworms, deer ticks, fleas, grasshoppers, Japanese beetle grubs, mole crickets and sod webworms.	
	Tree, shrub, rose & flower pests: aphids, bagworms, cicadas, exposed thrips, fall cankerworms, inchworms, leafminers and other listed pests.	
	Nuisance pests (outdoors): ants, boxelder bugs, clover mites, cluster flies, crickets, earwigs, fleas, mosquitoes and other listed pests.	
WHERE TO USE	On lawns, trees and shrubs, roses and flowers and outside surfaces of buildings.	

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with

USE PRECAUTIONS AND RESTRICTIONS

SHAKE WELL BEFORE USE. READ ENTIRE LABEL BEFORE EACH USE.

APPLICATION RESTRICTIONS:

Not for broadcast use on indoor residential surfaces.

Do not apply this product in a way that Do not apply this product in a way that will contact adults, children, or pets, either directly or through drift. Remove pets, birds, and cover fish aquariums before spraying. Not for use in outdoor residential misting systems.

Do not breathe vapors or spray mist. Do not apply when food is present. Do not enter or allow others to enter treated area until sprays have dried.

All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

- (1) Treatment to soil or vegetation around structures;
 (2) Applications to lawns, turf, and
- other vegetation;
 (3) Applications to building foundations,
- up to a maximum height of 3 feet

Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks. driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications, only.

Do not make applications during rain Do not water the treated area to the point of runoff.

Do not apply directly to sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur, except as directed by this label.

ENTRY RESTRICTIONS:

Do not allow adults children or nets to enter the treated area until sprays have dried. Do not enter or allow others to enter until vapors, mists, and aerosols have dispersed, and the treated area has been thoroughly ventilated.

This product helps reduce annoying and troublesome listed outdoor insects. Do not spray animals. People and pets should not be allowed in treated areas until spray has dried.

• Apply insecticide when insects first appear and repeat as necessary but not more frequently than every 3 days. This product kills most troublesome listed outdoor insects, which destroy lawns, gardens, and are an annoyance on patios and at picnics.

HOSE END SPRAYER: Easy to Use. For Application to Lawns, Outdoor Ornamentals, Outside Surfaces of Buildings, and Other Outdoor Residential Areas. Snap off tab on top of sprayer before first use.

- 1. Make sure knob Control lever is in the OFF position. Shake well and attach to hose.
- 2. Turn water on at faucet, aim nozzle toward lawn and turn knob Control
- lever to ON to begin spraying.

 3. When finished, turn Control lever to OFF position. Turn faucet to OFF position. Discharge residual water pressure by turning ON and OFF again. Disconnect sprayer from hose. Store out of reach of children

Barrier Spray For Nuisance Pests: Such as Ants, Boxelder Bugs, Clover mites, Cluster flies, Crickets, Earwigs, Fleas, Flying moths, Gnats, House flies, Millipedes, Mosquitoes, Sowbugs (pillbugs), Spiders, Tent caterpillars, Ticks (Brown dog, Deer, Lone Star).

Use Directions:

Use 32 fl. oz. to treat 1,900 linear feet with a 6-inch spray band. For use on lawns: 32 fl. oz. treats up to 5,300 square feet.

			clove	nce pests (outdoors): ants, boxelder bugs, r mites, cluster flies, crickets, earwigs, fleas, uitoes and other listed pests.
WHERE TO USE			On lawns, trees and shrubs, roses and flowers and outside surfaces of buildings.	
PESTS				HOW TO USE
	Outside surfaces of buildings	Ants, Boxelder Bugs, Clover mites, Crickets, Earwigs, Millipedes, Sowbugs (pillbugs)		To help prevent infestation of buildings, treat the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Also, apply as a residual spary to outside surfaces of buildings, including porches, screens, window

PESTS		HOW TO USE
Outside surfaces of buildings	Ants, Boxelder Bugs, Clover mites, Crickets, Earwigs, Millipedes, Sowbugs (pillbugs)	To help prevent infestation of buildings, treat the building foundation to a height of 2 to 3 feet, where pests are active and may find entronce. Alo, apply as a residual spray to outside surfaces of buildings, including porches, screens, window frames, eaves, patios, picnic areas, garages, retuse piles, and other areas where these pests congregate or have been seen. Hold sprayer from (2) to three (3) feet from surface to be treated. Spray to slightly moisten the surface. To maintain effectiveness, reapply as reinfestation occurs.
Other outside area	Ants, Crickets, Earwigs, Fleas, Spiders, Ticks, (Brown dog, Deer, Lone star)	For treatment of localized infestations of these insects in areas where there are weeds, shrubs or busty non-rop areas, spray infested areas thoroughly. For ants, thoroughly wet hills and runways. Repeat application as reinfestations occur.
	Boxelder Bugs, Cluster flies, Flying moths, Gnats, House flies, Mosquitoes	For use only as an aid in reducing annoyance from these insects. Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porthe, under potio furniture, in garages and other places where these insects alight or congregate. Repeat as necessary.
	Tent caterpillars	Application should be made when caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening, when caterpillars have returned to their tents. Wet tents with spray on all sides. A few caterpillars may be seen escaping from the tent, however, they should die within several hours.