World rights reserved. Marysville, OH 43040 © 2016 Tomcat Brands. Tomcat Brands

vailable Exclusively From: www.tomcat.com **S** C911 1-811-332-0155

Questions or Comments?

continually monitored and retreated it activity returns. fields, streams/lakes, etc. are especially prone to this. These areas will have to be NOTE: All properties are susceptible to recolonization. However, properties near woods,

 $^{**}$  Deep tunnel/mound systems require a slightly different procedure. See bottom panel was used to verify active runways. Return 48–72 hours later and retreat all active runs. Five to seven days after application, check for mole activity using the same procedure that



stlueeA gnitieB gninimneteO &

feet of active runway. application every 5-10 rock. Repeat bait possible, cover with a small hole closed. If this is not Carefully reseal by pinching



grubs should be exposed. типмау. Ио рагго от тпе grubs are completely in the opening, ensuring that the placement) into the Drop two grubs (one



diameter of the bait. larger than the yhtdgils bor a dtiw







baited. (See Step 2) holes resealed within \$\text{N}\$ hours should be 48–72 hours later. Runways that have had Mark opened runways and revisit them



not crush runway. subsurface $^{**}$  runways. NOTE: Be careful to to got ant ni alon a arutanug ot bor wornen Use your finger, small wooden dowel or a



syswnuA evitoA yineV

## APPLICATION DIRECTIONS

## HOW TO APPLY TOMCAT® MOLE KILLER GRUBS





Other mole baits have been developed on the premise that moles are similar to rats and mice, en in reality moles are insectivores (insect eaters). Moles readily consume earthworr when available, grubs. In fact, more than 90% of a mole's average daily dietary requirements are from earthworms and grubs.

Thousands of hours of groundbreaking laboratory and field research have gone into the product you are holding. Tomcat Mole Killer Grubs has been carefully designed to mimic a mole's most common natural food source — earthworms and grubs. Moles may consume a lethal dose in a single feeding, but it may take two or more days from the time of bait consumption for moles





Mimics a Mole's\* Natural Food Source, The Grub

Results Guaranteed+



KEEP OUT OF REACH OF CHILDREN **CAUTION** 

See bottom and side panels for First Aid, additional precautionary statements, and Directions for Use.

8 Grubs 2 Grubs Per Placement NET WT. **0.7 OZ** (20 g)

PIXELAUTIONARY SIAILMENTS
Hazards to Humans and Domestic Animals
CAUTION: Harmful if swallowed or absorbed through the skin.
Keep away from children, domestic animals and pets. Do not get in eyes, on skin, or on clothing.
Anyone handling baits must wear gloves. Wash thoroughly with soop and water after handling and before eating, diriking, chewing gum, or using tobacco.

Prolonged or frequent repeated skin contact may cause allergic

Ready To Use

species

Scientifically designed and tested to

mimic a mole's natural food source

Effective against most common mole\*

 Unit of pulsari country certain the pulsaria from the pulsari Wash with plenty of soap and water.

\*Not for use against the star-nosed mole in North Carolina

FIRST AID

HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE

USE RESTRICTIONS: This product may only be applied at the use sites and by the application methods indicated on this label. This product may only be used in lawns and around homes to control eastern moles (Scalpus aquaficus), stannoed moles\* (Condylura aristate parva) or Scanapus spp.

Boit must be applied directly into moles burnow systems. This be that the public directly into the runnway, using the rod to push any exposed bait completely into the runnway. Carefully reseal the runnway with a model signal and unaways or in subsurface feeding tunnels. This boit may not be used above ground. Wear gloves when boit is handled.

\*\*Not for use against the stannoed mole in North Carolina and unaways or in subsurface places and the public directly into the stannoed mole of cardinary using the same procedure that was used to establish initial activity. Return 48 to 72 hours later and retent all active runs.

\*\*SELECTION OF TREATMENT AREAS: The presence of moles may be indicated by a network of subsurface runways in the turn or by a series of conical mounds of earth pushed up from deep burnows. Subsurface runways are indicated by praised of substanding the subsurface runways are indicated by raised of substanding the subsurface runways are indicated by raised of substanding the substanding the public of the publ

boil. Do not apply directly to water.

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

APPLICATION DIRECTIONS:
Remove and place grub shaped boil according to use directions below and ADDITIONAL GRAPHICS AND TEXT.

To locate deep tunnels, push a probe, such as a broom handle, into the soil between mounds. When the probe enters a tunnel, the resistance on the probe will decrease applied, Mark opened tunnels and revisit them 48 to 72 hours later.

112415/07-15



TREATMENT FOR PET POISONING
If an animal eats bait, call veterinarian or 1-877-332-0755

Bromethalin (CAS #63333-35-7):

OTHER INGREDIENTS:

NOTE TO PHYSICIAN OR VETERINARIAN

EPA REG. NO. 90780-2 EPA EST. NO 12455-WI-1

\*Not for use against the star-nosed male in North Carolina

NOI I OF PATISION OR VELEXINAXION.
Contains the nerve poisons Bonembralia. This is not an anticoagulant type rodenticide. For humans or animals ingesting bait and/or showing poisoning signs (such as muscle tremos, loss of hind limb use, or seizures for animals), limit absorption by either emesis or gastric largue. Sublethal symptoms, if present, would be the result of creebral define and should be treated accordingly through administration of an asmotic diuretic and corticosteroid.

Environmental Hazards
This product is extremely toxic to mammals and birds. Dogs and other predatory and sorvenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water.

below and ADDITIONAL GRAPHICS AND TEXT.

SUBSURFACE RUNWAYS: Tomcar Mole Killer Grubs has been great funnels and revisit them 48 to 72 hours later. Runways that how had holes resealed within 72 hours should be baited.

SUBSURFACE RUNWAYS: Tomcar Mole Killer Grubs has been specially designed to mimic the natural food sources of the placed in areas used habitually by moles for feeding.

STABLISHING INITIAL ACTIVITY:

Be your finger, small wooden dowel or a narrow rod to of soul, and in the tome of subsurface runways. Be careful on the turnel. Cover the hole with a piece of son, a small rock or a piece of nevespoper to allow for visual of son of subsurface runways. Be careful on the turnel. Cover the hole with a piece of son, a small rock or a piece of nevespoper to allow for visual of son of subsurface runways. Be careful on the turnel. Cover the hole with a piece of son, a small rock or a piece of nevespoper to allow for visual on the turnel. Cover the hole with a piece of son, a small rock or a piece of nevespoper to allow for visual on the turnel. Cover the hole with a piece of son, a small rock or a piece of nevespoper to allow for visual on the turnel. Cover the hole with a piece of son, a small rock or a piece of nevespoper to allow for visual of son of subsurface runways. Be careful and the turnel cover the hole with a piece of son, a small rock or a piece of never piece of never piece. The substitute of t