

## PGRAGE MARCH

**Controls over 80 listed insect pests** 

Use on roses, flowers, fruits, & vegetables

For use on dogs and listed domestic animals

Kills and repels for up to 4 weeks

ACTYCHIGHEDISAT:
Permistrati (CAS 116, 52845-48-1). 0,285OTHER INCREDISATE: 58.765TOTAL. 700,00%

EPA Ed. No. 4-MY-1 EPA Rep. No. 4-480-61801

Net Weight 4 lbs (1.8 kg.)

Keep Out of Reach of Children
(See back panel for additional)
(Precautional) Statements and Fors Auth CAUTION

path to other the bound area with the dusts have reliest. Our stoppy as a breaking treatment to recover our say as more, they can contain acrossly treatment and regards before.

ovitata application med he limed to gold of price ped covera humanism only account for the libburing possible year.
Technol 19 ped to septiate formed to pediate.
Libburing to pediate to the septiate of the

Tien application to building Augusticay, of extinor applications to empression such as selectable, differency, pulses, porches and one surfaces (Soft as Windows, change and nevers are builded to personal most earlier applications only.)

And apply when over any score in the area. On you apply find is supposed within 24 hours. On not apply 4 wind causes shift, Do not water the elections on the policy of part of the applications being not

plication a prohibited durity into prevent or chains, or to any over time a puter where durings to severn, other during, water bodge, no aquate Aftel can occur. Do no along the product or error any draw during or other application.

For Dusting Plants: Be sure to lightly cover all plant surfaces

For Dusting Non-plant Surfaces: Using this container or other suitable mechanical duster, such as a bulbous duster, make a light but uniform application lobowing the specific directions for each use area listed on the label. Apply only as a spot treatment indoors. Do not broadcast, Exposed tools should be covered or removed to below application.

NUISANCE PESTS IN OUTDOOR AREAS

NUISANCE PESIS IN OUIDOUR AREAS

OUTSIDE SURFACES OF BUILDING

Use on Ants, Clover imites, Crickets, Earwigs, Millipedes, Spliders, Cockroaches, Sowbuss, (Pillbugs) to help prevent infestation of buildings tend the building foundation to a height of 2 to 3 few, where pests are active and may find entirance. Also, to outside surfaces of building, including porches, decks, window tranes, caves, patios, garages, relute dumps and other areas where these pests congregate or have been seen. Repeat irratment are needed to maintain effectiveness, but no note than one overy? 10 to 40 gar.

Iteatiment as needed to maintain effectiveness, but no more than once every 7 to 10 days.

To temporarily control House Files, Mosquittoes, Chister Files, Grats, Fyring Moths: Oust Peats on outside surfaces of screens, doors, window frames or wherever these viscots may enter the room. Also treat surfaces around light findness on porches, in garages, and other places where these insects may alight or congregate. Repeat treatment as needed to maintain effectiveness, but no more than once every 7 to 10 days.

OTHER OUTDOOR AREAS

Use on Ants, Crickets, Earwigs, Fleas, Lone Star Ticks for treatment of localized infestations of these insects in areas where there are weeds or bushy non-crop areas, cover infested areas with a light coaling of clus, For Ants finebuding Fire Ants), thoroughly treat hills and runways. Repeal treatment as needed to maintain effectiveness, but no more than once every 7 to 10 days.

NUISANCE PESTS INDOORS Application Within Residential Buildings Including Homes and Apartment Buildings

For Spot Treatment Only: Apply as a light but uniform dust to localized areas where cockroaches (including strains resistant to other insecticides), ants, other miles, crickets, silverfish and spiders are found or normally occur including dark corners of rooms and closets, cracks and crivices in walls, silverg and behind baseboards; beneath and behind sirks, slover, ordispersion and calcinets, around plumbing and other utility installations. For Anta: Apply to an trains, also around doors and wedows and wherever their these peats may find enhance.

Corpenier Ants: Apply dust incured doors and windows and other places where anis enter premises and where they crawl or hide. Also force dust into cracks and crevious or through openings or small newly drilled holes into wall voids where anis anction nests are present. Use a sufficient amount to fighty over are sufficient.

to lightly cover area surfaces.

For Control of Cappet Beelles: Apply dust as a spot treatment along baseboards and edges of carpeting, under carpeting, rugs, and furniture, in closes and on shelving, and wherever these insects are seen or suspected.

For Control of Ticks and Fleas: Apply dust as a spot treatment to infested areas, nearby cracks and crevices, along and behind baseboards, window and door learnes on locatived areas of floor and floor covering where these pests may be prosent. For best results, old get bedding should be cleaned and replaced. To control the source of fless affestations, pets whatling the treated premises should be treated as directed below.

And door large and locatized areas of floor and floor covering where triese peace may be a made and a man a man and a man a man and home a man and home a man a

DIRECTIONS FOR APPLYING ROSES & FLOWERS				
Sites	Pests	How to Use		
Marigold, Mock-grange, Nannyberry, Orchid, Palm, Pansy, Pea Shrub, Peperomia, Petunia, Philodendron, Piggy-Back Plant,	Worms, Cicadas, Comworms, Cutworms, Exposed Thrips, Fall Cankerworms, Fall	Apply at the first sign of insects. Lighty cover all plant surfaces. Repeat as necessary, use intervals of 4-8 days. Application can be made up to the day of collection.		

Use with care on ferris, paties and fuschia. As with all plants listed above, especially carnations, chrysanthemums, and roces, there exists a wice variety of species and cultivars, and theeters, a wide variety of succeptibility to injury. To ascertain possible phytotoxicity, cust a few plants as described above, wail 24 hours, and check for possible injury.

Repeat applications may be necessary to control adult Japanese beetles on roses.

On Tent Categoliars: Application should be made when categoliars are young and lenhs are first noticed. For best results, apply in the late afternoon or evening, when categoliars have returned to their tents. A few categoliars may be seen escaping from the tent; however, they should die within several hours.

VEGETABLES
For Home Vegetable Gardens: Dust upper and lower leaf surfaces. Use sufficient amount to obtain full coverage of all foliage. Apply no more than

Sites	Pests	How to Use
Asparagus (pre-harvest)	Asparagus Beelles, Culworms	Apply as needed to ensure uniform coverage. Do not apply more than 4 applications per crop. Can be applied up to 3 days prior to harvest.
	Larval slages of Asparagus Beelles; Tarnished Plant Bugs; Lygus Bugs and Adull Japanese Beelles.	Treat ferns or bush growth after spear harvest when insects are present.
Celery-See Leafy Vegetables		

Sites	Pests	How to Use
Cole Crops: Broccoli, Brussels Sprouls, Cabbage, Cauliflower	Beel Armyworms, Cabbage Aphids, Cabbage Loopers, Diamond Back Moths, Imported Cabbageworms	Apply every 5 days or as needed. Do not make more than 6 applications on Cauliflower, Brussels Sprouts and Broccoli and 10 applications on Cabbage. Can be
Corn-See Sweel Corn Courblist Baham Pear (Biller Melon), Chinese Waxgourd (Chinese Preserving Meton), Cilron Melon, Courumber, Gherkin, Edible Gourds; Melons including hybrids such as Canladupe Casaba, Crenshaw, Honeydew, Honey Balls, Mango Melon, Muskmeton, Perisan Melon, Pumpkin, Summer & Winter Squask, Naturmeton	Aphids, Cabbage Loopers, Cucumber Beelles (Adults), Cutworms, Leafhoppers, Leafminers, Melonworms, Pickieworms, Plant Bugs, Squash Bugs,	Apply as needed to ensure uniform coverage, Do not make more than 8 applications per crop, Can be applied up to the day of harvest
Eggplant	Colorado Potato Beetles, Vegetable Leafminers	Apply as needed to ensure uniform coverage. Do not exceed 16 applications. Can be applied up to 3 days prior to baryes!
Horseradish	Imported Cruciler Weevils	Apply as needed. Do not exceed 3 foliar applications. Can be applied up to 22 days prior to harvest.
Leafy Vegelables: Amaranih, Arugula, Calery, Celluce, Chervil, Corn Salad, Chrysanlhemum (Edible-Leaved and Garland), Cress (Garden and Upland), Dandelion, Dock, Endive, Fennel, Lelluce (Head and Leaf), Orach, Parsley, Purslane (Garden and Winter), Probatts, Someth, Sarss Charle	Alfalla Loopers, Aphiks, Beel Armyworms, Cabbage Loopers, Corn Earworms, Culworms, European Corn Borers, Fall Armyworms, Green Cloverworms, Southern Armyworms, Tobacco Budworms, Vegelable Leafminers.	Apply every 5 to 10 days as needed. Do not apply more than 7 applications per crop. Can be applied up to 1 day prior to harvest
Melons-See Cucurbils Polatoes	Beel Armyworms, Colorado Potato Beelles, Polalo Lealhoppers, Polalo Tuberworms, Cabbage Loopers, Cutworms, European Corn Borers, Polalo Psyllids, Tarnished Plant Buns	Apply as needed. Do not apply more than 12 applications per crop, Can be applied up to 7 days prior to harvest.
Peppers	Vegelables leafminers, Cabbage looper, Flea beelle, Corn earworm, Pepper weevil, European corn borer	Apply as needed. Do not apply more than 8 applications per crop. Can be applied up to 3 days prior to barvest.
Sweet Corn	Corn Earworms, European Corn Borers, Fall armyworms, Southern armyworms	Apply every 5 days and when needed. Do not apply more than 6 applications per season. Can be applied up to 1 day prior to harvest.
Tomaloes	Beel and Southern Armyworms, Tomalo Pinworms, Tomalo Fruitworms, Horn Worms, Granulale Cutworms, Colorado Polato Beelles, Cabbage Loopers	Apply as needed for uniform coverage Do not apply more than 6 applications per season. Can be applied up to day of harvest.

DOMESTIC ANIMALS
USE ONLY ON DOGS, COMPANION LIVESTOCK (CATTLE, HORSES, SWINE) AND POULTRY. Consult a velerinarian before using this product

USE ONLY ON DIOGS, COMPANION LIVESTOCK (CATTLE, HORSES, SWINE) AND POULTRY. Consult a veterinarian before using his product on debiliated, segd, regenant, invising, or medicated animals.

Dogs: To control (xill) Fleas, Ticks and Lice: Use product only outside or in a well-ventilated area. Wearing household latex or rubber gloves, dust entire animal avoiding pets eyes, nose, mouth and genital rates. Rub or brush pets hair to work dust down to the skin. Repeal every three weeks. This product should not be applied by children. Do not use on pregnant animals. Do not use on puppies under 12 weeks rage. Dust pet bedding and dophouse using a shaker can or other dust applicator.

deployee using a shaker can or other dust applicator.

Cattle, Horses: Can be used in dust bags, shaker can and mechanical dust applicator. To control (kill) Hornfles, Lice, Face Files: Place contents of his package in A commercially available dust blag, suspend bag in areas frequented by animals or in gateways or lanes through which the animals pass daily for water, feed, or minerals. Bags may also be placed in fusting sheets or in front of mineral feeders. For milking cows, bags may be suspended in the evil intrough which the cowns leave the milking barn. The bags should have great on the milking barn. The bags should have great of the interest of the first part of the use of the first part of the first

bedding may be treated. Do not slaughter animals within 5 days of treatment.

Poultry: To control (kill) Northern Fowl Mites apply 1 lb, per 100 birds. Ensure thorough treatment of vent area.

STORAGE AND DISPOSAL

Posticide Storage; Store and transport in an upright position. Store in a cool dry area inaccessible to children or pets.

Posticide Disposal: If partly filled — Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outlood or drain.

Container Disposal: If empty — Nonverillable container. Do not reuse or refit this container. Place in trash or offer for recycling if available.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Sensitivities may occur after using ANY pesticide product for gets. If signs of sensitivity occur, bathe your pet with mild seap and rinse with large amounts of water. It signs continue, consult a veterinarian immediately. Do not allow pets which are not listed for use on this tabel to contact treated areas.

DO NOT USE ON CATS. May be toxic or potentially total if applied to or ingested by cats. Accidental application to cats and/or grooming with a recently treated dog may result in tremors and/or uncoordinated muscle movement. If this occurs, immediate veterianzy cate should be provided.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to agratic organisms, including fish and inventebroies. To protect the environment, do not allow p to enter or run off into storm drains, drainage ditches, quiters or surface waters. Applying this product in calm weather when rain predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. This pringing look to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming or weeds while bees are actively visiting the treatment area.

Distributed by: Orgill, Inc. 3742 Tyndale Drive P.O. Box 140 Memphis TN, 38101

