



STERILEX[®] ULTRA STEP

Using PerQuat[®] Technology

Floor Sanitizer

Foot Pan, Boot Sanitizer

Floor sanitizing and deodorizing performance

For Use in Non-Food Contact Areas of Dairy, Food, Meat and Beverage Processing Facilities

Animal Premise Sanitizer

Active Ingredients:

N-Alkyl (C ₁₄ 95%, C ₁₂ 3%, C ₁₆ 2%) dimethyl benzyl ammonium chloride	1.9%
Sodium Carbonate Peroxyhydrate.....	12.1%
Other Ingredients	86.0%
TOTAL:.....	100.0%

KEEP OUT OF THE REACH OF CHILDREN

DANGER

PRECAUTIONARY STATEMENTS

Hazard To Humans And Domestic Animals

DANGER. Corrosive. Causes irreversible eye damage and skin irritation. Harmful if swallowed or absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Wear protective eyewear (goggles, face shield, or safety glasses), coveralls worn over short-sleeved shirt and short pants, socks, chemical resistant footwear and waterproof gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

Sterilex Ultra Step is a moisture activated sanitizer specifically designed to improve environmental sanitation on surfaces where conditions favor microorganism growth and cause sanitation concern.

Sterilex Ultra Step is for use on hard, nonporous surfaces in: federally inspected food processing facilities, federally inspected meat and poultry plants, dairies, wineries, breweries, beverage manufacturing facilities, dairy farms, swine farms, poultry and turkey farms, farm premises, hatcheries, refrigerated trucks, kennels, pet animal quarters, zoos, pet shops, animal laboratories, veterinary facilities, animal care facilities, food service facilities, kitchens, grocery stores and deli areas.

Net Content 10 lbs.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SANITIZING OF HARD, NON-POROUS, INANIMATE, NON-FOOD CONTACT SURFACES:

Sterilex Ultra Step eliminates 99.9% of the following bacteria in 5 minutes on hard, non-porous, inanimate, non-food contact surfaces according to the Standard Test Method for Efficacy of Sanitizers Recommended for Inanimate Non-Food Contact Surfaces:

Enterobacter aerogenes (ATCC 13048)	Staphylococcus aureus (ATCC 6538)	Salmonella enterica (ATCC 10708)
Escherichia coli (ATCC 11229)	Listeria monocytogenes (ATCC 15313)	Salmonella enterica enteritidis (ATCC 13076)

SANITIZING DIRECTIONS FOR NON-FOOD CONTACT SURFACES:

Add a solution of 4.9 – 6.1 ounces Sterilex Ultra Step per gallon of water (or equivalent use dilution) to sanitize hard, non-porous surfaces. Apply sanitizer use solution to pre-cleaned, hard, non-porous surfaces with a cloth, mop, sponge, sprayer, foaming or by immersion. For sprayer applications, use a coarse pump or trigger sprayer. Spray 6-9 inches from surface. Do not breathe spray. Treated surfaces must remain wet for 5 minutes, then wipe with sponge, mop or cloth or allow to air dry. Use within 8 hours after powder has been dissolved in water.

(For food processing or other facilities that have installed entryway sanitizing systems)

ENTRYWAY SANITIZING SYSTEMS AND FOOTBATHS: To prevent cross contamination on treated surfaces, apply a solution of 4.9 – 6.1 ounces Sterilex Ultra Step per gallon of water (or equivalent use dilution). The entire entryway path must be covered with the footbath. Use within 8 hours after powder has been dissolved in water.

ENTRYWAY SHOE FOAM DIRECTIONS: Apply a foam layer that is approximately 0.5 to 2 inches (1.2 cm to 5 cm) thick at all entrances using a foam generating machine or aerator and a solution containing 4.9 – 6.1 ounces Sterilex Ultra Step per gallon of water (or equivalent use dilution). Scrape waterproof shoes and boots. Walk through foam. Boots must remain wet for at least 5 minutes prior to entering area. Foamed entranceways must be washed and the foam replaced at least daily or when it (foam/area) appears dirty. Foam should be applied to entrances of buildings, hatcheries, production and packaging rooms to prevent cross contamination from area to area in the packaging and storage areas of food plants. Use within 8 hours after powder has been dissolved in water

APPLICATION INSTRUCTIONS FOR USE AS A FLOOR SURFACE, FLOOR MAT, FOOT PAN AND BOOT

SANITIZER: After cleaning and sanitizing equipment, apply Sterilex Ultra Step evenly to the floor, foot pan, floor mat, containment device and/or drain surface, at a rate of 3.7 – 7.4 oz. (105 – 210 grams) per 100 square feet of floor surface containing 0.25 fl. oz. – 0.97 fl. oz. (7.4 ml – 28 ml) of water per square foot. Product may be re-applied as needed during the production process. Sterilex Ultra Step does not become active until it comes in contact with moisture.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal

PESTICIDE STORAGE: Keep out of children's reach. Store at room temperature out of direct sunlight, DO NOT store above 30°C (or 86°F).

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Fill the container ¼ full with water and recap. Shake for 10 seconds. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Dispose of rinsate as pesticide waste. If rinsate cannot be used, follow pesticide disposal instructions. Offer for recycling, if available or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. If not triple rinsed these containers are acute hazardous wastes and must be disposed in accordance with local, state and federal regulations.

Manufactured for
STERILEX CORPORATION
111 Lake Front Drive, Hunt Valley, MD 21030
Phone: 800-511-1659 or 443-541-8800

MADE IN USA

EPA ESTABLISHMENT: 067441-IL-001
EPA REGISTRATION: 63761-10

Revision 6376110-20160812



STERILEX® ULTRA STEP

Using PerQuat® Technology

Floor Sanitizer

Foot Pan, Boot Sanitizer

Floor sanitizing and deodorizing performance

For Use in Non-Food Contact Areas of Dairy, Food, Meat and Beverage Processing Facilities

Animal Premise Sanitizer

Active Ingredients:

N-Alkyl (C ₁₄ 95%, C ₁₂ 3%, C ₁₆ 2%) dimethyl benzyl ammonium chloride.....	1.9%
Sodium Carbonate Peroxyhydrate.....	12.1%
Other Ingredients	86.0%
TOTAL:.....	100.0%

KEEP OUT OF THE REACH OF CHILDREN

DANGER

PRECAUTIONARY STATEMENTS

Hazard To Humans And Domestic Animals

DANGER. Corrosive. Causes irreversible eye damage and skin irritation. Harmful if swallowed or absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Wear protective eyewear (goggles, face shield, or safety glasses), coveralls worn over short-sleeved shirt and short pants, socks, chemical resistant footwear and waterproof gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

Sterilex Ultra Step is a moisture activated sanitizer specifically designed to improve environmental sanitation on surfaces where conditions favor microorganism growth and cause sanitation concern.

Sterilex Ultra Step is for use on hard, nonporous surfaces in: federally inspected food processing facilities, federally inspected meat and poultry plants, dairies, wineries, breweries, beverage manufacturing facilities, dairy farms, swine farms, poultry and turkey farms, farm premises, hatcheries, refrigerated trucks, kennels, pet animal quarters, zoos, pet shops, animal laboratories, veterinary facilities, animal care facilities, food service facilities, kitchens, grocery stores and deli areas.

Net Content 100 lbs.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SANITIZING OF HARD, NON-POROUS, INANIMATE, NON-FOOD CONTACT SURFACES:

Sterilex Ultra Step eliminates 99.9% of the following bacteria in 5 minutes on hard, non-porous, inanimate, non-food contact surfaces according to the Standard Test Method for Efficacy of Sanitizers Recommended for Inanimate Non-Food Contact Surfaces:

Enterobacter aerogenes (ATCC 13048)	Staphylococcus aureus (ATCC 6538)	Salmonella enterica (ATCC 10708)
Escherichia coli (ATCC 11229)	Listeria monocytogenes (ATCC 15313)	Salmonella enterica enteritidis (ATCC 13076)

SANITIZING DIRECTIONS FOR NON-FOOD CONTACT SURFACES:

Add a solution of 4.9 – 6.1 ounces Sterilex Ultra Step per gallon of water (or equivalent use dilution) to sanitize hard, non-porous surfaces. Apply sanitizer use solution to pre-cleaned, hard, non-porous surfaces with a cloth, mop, sponge, sprayer, foaming or by immersion. For sprayer applications, use a coarse pump or trigger sprayer. Spray 6-9 inches from surface. Do not breathe spray. Treated surfaces must remain wet for 5 minutes, then wipe with sponge, mop or cloth or allow to air dry. Use within 8 hours after powder has been dissolved in water.

(For food processing or other facilities that have installed entryway sanitizing systems)

ENTRYWAY SANITIZING SYSTEMS AND FOOTBATHS: To prevent cross contamination on treated surfaces, apply a solution of 4.9 – 6.1 ounces Sterilex Ultra Step per gallon of water (or equivalent use dilution). The entire entryway path must be covered with the footbath. Use within 8 hours after powder has been dissolved in water.

ENTRYWAY SHOE FOAM DIRECTIONS: Apply a foam layer that is approximately 0.5 to 2 inches (1.2 cm to 5 cm) thick at all entrances using a foam generating machine or aerator and a solution containing 4.9 – 6.1 ounces Sterilex Ultra Step per gallon of water (or equivalent use dilution). Scrape waterproof shoes and boots. Walk through foam. Boots must remain wet for at least 5 minutes prior to entering area. Foamed entranceways must be washed and the foam replaced at least daily or when it (foam/area) appears dirty. Foam should be applied to entrances of buildings, hatcheries, production and packaging rooms to prevent cross contamination from area to area in the packaging and storage areas of food plants. Use within 8 hours after powder has been dissolved in water

APPLICATION INSTRUCTIONS FOR USE AS A FLOOR SURFACE, FLOOR MAT, FOOT PAN AND BOOT

SANITIZER: After cleaning and sanitizing equipment, apply Sterilex Ultra Step evenly to the floor, foot pan, floor mat, containment device and/or drain surface, at a rate of 3.7 – 7.4 oz. (105 – 210 grams) per 100 square feet of floor surface containing 0.25 fl. oz. – 0.97 fl. oz. (7.4 ml – 28 ml) of water per square foot. Product may be re-applied as needed during the production process. Sterilex Ultra Step does not become active until it comes in contact with moisture.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal

PESTICIDE STORAGE: Keep out of children’s reach. Store at room temperature out of direct sunlight, DO NOT store above 30°C (or 86°F).

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Completely empty bag into application equipment by shaking and tapping sides and bottom to loosen clinging particles. If not emptied in this manner, the bag may be considered an acute hazardous waste and must be disposed in accordance with local, state and federal regulations. When completely empty, offer for recycling if available or dispose of bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Manufactured for
STERILEX CORPORATION
111 Lake Front Drive, Hunt Valley, MD 21030
Phone: 800-511-1659 or 443-541-8800

MADE IN USA

EPA ESTABLISHMENT: 067441-IL-001
EPA REGISTRATION: 63761-10

Revision 6376110-20160812



STERILEX® ULTRA STEP

Using PerQuat® Technology

Floor Sanitizer

Foot Pan, Boot Sanitizer

Floor sanitizing and deodorizing performance

For Use in Non-Food Contact Areas of Dairy, Food, Meat and Beverage Processing Facilities

Animal Premise Sanitizer

Active Ingredients:

N-Alkyl (C ₁₄ 95%, C ₁₂ 3%, C ₁₆ 2%) dimethyl benzyl ammonium chloride.....	1.9%
Sodium Carbonate Peroxyhydrate.....	12.1%
Other Ingredients	86.0%
TOTAL:.....	100.0%

KEEP OUT OF THE REACH OF CHILDREN

DANGER

PRECAUTIONARY STATEMENTS

Hazard To Humans And Domestic Animals

DANGER. Corrosive. Causes irreversible eye damage and skin irritation. Harmful if swallowed or absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Wear protective eyewear (goggles, face shield, or safety glasses), coveralls worn over short-sleeved shirt and short pants, socks, chemical resistant footwear and waterproof gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

Sterilex Ultra Step is a moisture activated sanitizer specifically designed to improve environmental sanitation on surfaces where conditions favor microorganism growth and cause sanitation concern.

Sterilex Ultra Step is for use on hard, nonporous surfaces in: federally inspected food processing facilities, federally inspected meat and poultry plants, dairies, wineries, breweries, beverage manufacturing facilities, dairy farms, swine farms, poultry and turkey farms, farm premises, hatcheries, refrigerated trucks, kennels, pet animal quarters, zoos, pet shops, animal laboratories, veterinary facilities, animal care facilities, food service facilities, kitchens, grocery stores and deli areas.

Net Content 400 lbs.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SANITIZING OF HARD, NON-POROUS, INANIMATE, NON-FOOD CONTACT SURFACES:

Sterilex Ultra Step eliminates 99.9% of the following bacteria in 5 minutes on hard, non-porous, inanimate, non-food contact surfaces according to the Standard Test Method for Efficacy of Sanitizers Recommended for Inanimate Non-Food Contact Surfaces:

Enterobacter aerogenes (ATCC 13048)	Staphylococcus aureus (ATCC 6538)	Salmonella enterica (ATCC 10708)
Escherichia coli (ATCC 11229)	Listeria monocytogenes (ATCC 15313)	Salmonella enterica enteritidis (ATCC 13076)

SANITIZING DIRECTIONS FOR NON-FOOD CONTACT SURFACES:

Add a solution of 4.9 – 6.1 ounces Sterilex Ultra Step per gallon of water (or equivalent use dilution) to sanitize hard, non-porous surfaces. Apply sanitizer use solution to pre-cleaned, hard, non-porous surfaces with a cloth, mop, sponge, sprayer, foaming or by immersion. For sprayer applications, use a coarse pump or trigger sprayer. Spray 6-9 inches from surface. Do not breathe spray. Treated surfaces must remain wet for 5 minutes, then wipe with sponge, mop or cloth or allow to air dry. Use within 8 hours after powder has been dissolved in water.

(For food processing or other facilities that have installed entryway sanitizing systems)

ENTRYWAY SANITIZING SYSTEMS AND FOOTBATHS: To prevent cross contamination on treated surfaces, apply a solution of 4.9 – 6.1 ounces Sterilex Ultra Step per gallon of water (or equivalent use dilution). The entire entryway path must be covered with the footbath. Use within 8 hours after powder has been dissolved in water.

ENTRYWAY SHOE FOAM DIRECTIONS: Apply a foam layer that is approximately 0.5 to 2 inches (1.2 cm to 5 cm) thick at all entrances using a foam generating machine or aerator and a solution containing 4.9 – 6.1 ounces Sterilex Ultra Step per gallon of water (or equivalent use dilution). Scrape waterproof shoes and boots. Walk through foam. Boots must remain wet for at least 5 minutes prior to entering area. Foamed entranceways must be washed and the foam replaced at least daily or when it (foam/area) appears dirty. Foam should be applied to entrances of buildings, hatcheries, production and packaging rooms to prevent cross contamination from area to area in the packaging and storage areas of food plants. Use within 8 hours after powder has been dissolved in water

APPLICATION INSTRUCTIONS FOR USE AS A FLOOR SURFACE, FLOOR MAT, FOOT PAN AND BOOT

SANITIZER: After cleaning and sanitizing equipment, apply Sterilex Ultra Step evenly to the floor, foot pan, floor mat, containment device and/or drain surface, at a rate of 3.7 – 7.4 oz. (105 – 210 grams) per 100 square feet of floor surface containing 0.25 fl. oz. – 0.97 fl. oz. (7.4 ml – 28 ml) of water per square foot. Product may be re-applied as needed during the production process. Sterilex Ultra Step does not become active until it comes in contact with moisture.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal

PESTICIDE STORAGE: Keep out of children’s reach. Store at room temperature out of direct sunlight, DO NOT store above 30°C (or 86°F).

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Completely empty bag into application equipment by shaking and tapping sides and bottom to loosen clinging particles. If not emptied in this manner, the bag may be considered an acute hazardous waste and must be disposed in accordance with local, state and federal regulations. When completely empty, offer for recycling if available or dispose of bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Manufactured for
STERILEX CORPORATION
111 Lake Front Drive, Hunt Valley, MD 21030
Phone: 800-511-1659 or 443-541-8800

MADE IN USA

EPA ESTABLISHMENT: 067441-IL-001
EPA REGISTRATION: 63761-10

Revision 6376110-20160812