


Multi-Purpose
Fungicide



Monterey
Fungi-MaxTM
Brand

**Use on Ornamentals,
Lawns, Fruit and
Nut Trees, Berries
and Vegetables**



Active Ingredient:
Myclobutanil2.0%
Other Ingredients:98.0%
Total:100.0%

EPA Reg. No. 42750-196-54705
EPA Est. No. 48498-CA-1

ALB-081717N) 0416/0817(02)

KEEP OUT OF REACH OF CHILDREN

CAUTION See Attached Booklet for Precautionary
Statements and Directions for Use

NET CONTENTS: 1 PINT/473.16 mL

Monterey **Fungi-Max**[™] Brand

- **Do-It-Yourself Disease Control**
- **Use on Ornamentals, Lawns, Fruit and Nut Trees, Berries and Vegetables**
- **Systemic Action**
- **Protects For Up To 2 Weeks**



⚠ WARNING: Reproductive Harm – www.P65Warnings.ca.gov

EPA Reg. No. 42750-196-54705 EPA Est. No. 48498-CA-1

Manufactured For:

LAWN AND GARDEN PRODUCTS, INC.

P.O. Box 35000 • Fresno, CA 93745

(559) 499-2100 • info@montereylawngarden.com

Fungi-Max[™] is a trademark
of Lawn and Garden Products, Inc.

(ALB-081717N) 0416/0817(02)

KEEP OUT OF REACH OF CHILDREN
CAUTION

FIRST AID

IF IN EYES

Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED

Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

For transportation or medical emergency, please call 1-800-424-9300.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear protective eyewear. Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

USER SAFETY RECOMMENDATIONS

- User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- User should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area.

PRODUCT FACTS

What It Does:	Prevents and controls all Major Plant Diseases including Anthracnose, Brown Patch, Copper Spot, Dollar Spot, Leaf Spot (including Septoria, Zonate), Melting Out, Crown Rot, Leaf Smuts, Powdery Mildew, Rust, Red Thread, Spring Dead Spot, Necrotic Ring Spot and Summer Patch
Where To Use:	Outdoors on roses, flowers, ornamentals, shrubs, trees, lawns, fruit and nut trees, berries and vegetables.
When To Use:	Start Treating When Grass Begins to Turn Green in Spring or When Signs of Diseases are First Noticed.

**STOP. READ ENTIRE LABEL
BEFORE USE.**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For home use only. Shake well before using.

Keep pets and people other than applicator off treated areas until spray has dried.



WHEN TO USE

This product is highly effective when it is used to prevent diseases or applied at the first sign of disease. Start treating when grass begins to turn green in the spring or when signs of diseases are first noticed. For best disease control, treat every 2 weeks.

RESTRICTIONS

- Do not apply more than 13 treatments per year.
- In Nassau and Suffolk Counties in New York State, do not apply more than 3 times per given area per year.
- Do not treat lawns or plants during periods of stress, (i.e. heat, drought, flooding, insects, excessive fertilizer, herbicides or pesticide use, etc.)

- Do not apply when temperatures are above 85°F as injury to lawn can occur.
- Only use hose-end sprayers for broadcast lawn application. For application to individual plants and lawn spot treatment only use conventional pump up sprayers at labeled rates.
- Apply to the lawn after mowing, or allow the area sprayed to dry completely prior to mowing.
- If it rains before product has dried, reapply.
- If disease is already present, this product will not eliminate it.

If you need more information, contact your local County Extension Service.

LAWNS

MONTEREY FUNGI-MAX™ BRAND prevents and cures Anthracnose, Brown Patch, Copper Spot, Dollar Spot, Leaf Spot (including Septoria, Zonate), Melting Out, Crown Rot, Leaf Smuts, Powdery Mildew, Rust, Red Thread, Spring Dead Spot, Necrotic Ring Spot and Summer Patch on home lawns.

Use 2 2/3 fl. oz./half gallon of water to treat 250 sq. ft. Start treating when grass begins to turn green in the spring or when signs of diseases are first noticed. For best disease control treat every 2 weeks.

HOW TO USE

Determine the size of the lawn to be treated. To calculate square feet, multiply length by width. Start at the farthest point to be treated and work backward so you don't walk over treated area. Walk at a steady pace and spray using side to side (arc) motion, slightly overlapping the treated area.

Rainproof — rain or watering 2 hours after application will not wash away effectiveness. If rainfall occurs within 2 hours of application, the product should be reapplied.

ROSES, FLOWERS, ORNAMENTAL SHRUBS & TREES

MONTEREY FUNGI-MAX™ BRAND prevents and controls Black spot, Blight (blossom, cercospora leaf, ovulinia petal), Leaf spot (including cercospora), Powdery mildew, Rust (including needle, fusiform), Scab (including fusicladium) and other diseases on a wide variety of roses, flowers, trees and ornamental shrubs.

See table for a complete list of plants.

HOW TO TREAT

Use 3/4 fl. oz./gallon of water. Apply to the point when spray just begins to runoff the leaves. Apply once every 2 weeks throughout the season or until conditions favoring disease development (hot, moist weather) are no longer present. Allow spray to dry before watering.

USE TIPS

1. Treat roses on a 7 to 10 day schedule. If black spot is not a problem, a 10 to 14 day schedule is recommended.
2. When treating azaleas, mountain laurel or rhododendrons, begin spraying when flowers start to show color.
3. When treating irises, increase use rate to 1½ fl. oz./gallon of water.
4. For Douglas fir, loblolly pine and slash pine use 2¼ fl. oz./gallon of water. Begin treating in early spring and continue treating every 2 to 3 weeks until threat of infection has passed.

<p style="text-align: center;">PLANT</p> <p style="text-align: center;">(Number following plant refers to the specific Use Tip above.)</p>	<p style="text-align: center;">DISEASES</p>
<p>Rose (1)</p>	<p>Black spot, Powdery mildew, Rust</p>
<p>Azalea (2), Mountain laurel (2), Rhododendron (2)</p>	<p>Petal blight (ovulinia spp.), Powdery mildew</p>
<p>Abelia, Acalypha (Copper-leaf), Catalpa, Flowering cherry</p>	<p>Cercospora leaf spot</p>
<p>Hackberry, Honeysuckle, Privet, Trumpet creeper, Zinnia</p>	<p>Powdery mildew</p>
<p>Ageratum, Anemone, Angelica, Ash, Aster, Birch, China aster, Chokeberry, Chrysanthemum, Columbine, Cornflower, Dianthus, Flowering quince, Flowering maple (Abutilon), Four o'clock, Fuchsia, Iris (3), Marigold, Moonflower, Ninebark, Russian olive, Smoke-tree (Gothus)</p>	<p>Rust</p>

<p style="text-align: center;">PLANT</p> <p style="text-align: center;">(Number following plant refers to the specific Use Tip above.)</p>	<p style="text-align: center;">DISEASES</p>
<p>Achillea (Yarrow), Alder, Amelanchier (Juneberry, Shadebush), Barberry, Carnation, Delphinium, Gaillardia, Gardenia, Seranium, Hollyhock, Mock-orange, Pansy, Petunia, Photinia, Salvia, Snap-dragon, Viburnum</p>	<p>Powdery mildew, Rust</p>
<p>African Violet, Begonia, Bittersweet, Buckeye, California poppy, Chestnut (Horse), Cosmos, Crepe myrtle, Dahlia, Elm, Euonymus, Gourd (Ornamental), Grape leaf ivy, Lilac, Locust, Maple, Oak, Poinsettia, Spirea, Sycamore, Walnut, Willow</p>	<p>Powdery mildew</p>
<p>Flowering almond</p>	<p>Blossom blight (Monilinia spp.)</p>

(continued)

PLANT (Number following plant refers to the specific Use Tip above.)	DISEASES
Amorpha (False indigo), Bellflower, Buttonbush, Fleabane, Phlox, Sunflower	Cercospora leaf spot, Powdery mildew, Rust
Calendula, Hydrangea, Leucothoe	Cercospora leaf spot
Flowering crabapple, Flowering pear, Hawthorn	Powdery mildew, Rust, Scab
Dogwood	Anthracnose, Powdery mildew
Douglas fir (4)	Needle rust
Loblolly pine, Siasirpine (4)	Fusiform rust
Pyracantha	Fusicladium scab

FRUITS AND NUTS

Almond (blossom blight, rust, and shothole control; anthracnose suppression)

Mix 1/3 fl. oz. of MONTEREY FUNGI-MAX™ BRAND with 1 gallon of water and uniformly apply to all parts of the tree to point of runoff. For anthracnose, blossom blight, and shothole, apply at pink bud stage (when about 5% of the buds have bloomed). If conditions favoring disease development (hot, moist weather) exist, reapply at full bloom and when the petals fall. For rust, apply MONTEREY FUNGI-MAX™ BRAND about 8 weeks after flowering if conditions favoring disease development exist. Reapply every 21 days so long as disease symptoms persist.

- Do not apply within 90 days of harvest.
- Do not apply more than 3 times per season.

Apple and Mayhaw (powdery mildew, rusts and scab control)

Mix 1/2 fl. oz. of MONTEREY FUNGI-MAX™ BRAND with 1 gallon of water and uniformly apply to all parts of the tree to point of runoff. Reapply every 7 to 10 days so long as disease symptoms persist. For powdery mildew and rusts, apply at pink bud stage when the buds are opening and the pink petals are visible. For scab, apply MONTEREY FUNGI-MAX™ BRAND when the first green leaf tips are visible.

- Do not apply within 14 days of harvest.
- Do not apply more than 10 times per season.

STONE FRUITS

For all stone fruits, MONTEREY FUNGI-MAX™ BRAND may be applied up to the day of harvest.

Apricot and Nectarine (brown rot blossom blight, brown rot, powdery mildew, and shothole control)

Mix 1/3 fl. oz. of MONTEREY FUNGI-MAX™ BRAND with 1 gallon of water and uniformly apply to all parts of the tree to point of runoff. Apply at early red bud stage (the tips of the petals are visible and the flowers are still closed) before infection occurs. Reapply every 7 to 14 days so long as disease symptoms persist.

- Do not apply more than 7 times per season.

Cherry (brown rot blossom blight, brown rot, leaf spot, and powdery mildew control)

Mix 1/3 fl. oz. of MONTEREY FUNGI-MAX™ BRAND with 1 gallon of water and uniformly apply to all parts of the tree to point of runoff. Apply when single flower buds are visible (but still closed), on short stalks and before infection occurs. Reapply every 7 to 14 days so long as disease symptoms persist. For leaf spot, make an additional application of MONTEREY FUNGI-MAX™ BRAND after harvest.

- Do not apply more than 7 times per season.

Peach (brown rot blossom blight, brown rot, powdery mildew, and rust control)

Mix 1/3 fl. oz. of MONTEREY FUNGI-MAX™ BRAND with 1 gallon of water and uniformly apply to all parts of the tree to point of runoff. For brown rot blossom blight, brown rot, and powdery mildew, apply at early pink bud stage (the tips of the petals are visible and the flowers are still closed) before infection occurs. Reapply every 7 to 14 days so long as disease symptoms persist. For rust, apply MONTEREY FUNGI-MAX™ BRAND about 8 weeks after flowering if conditions favoring disease development (hot, moist weather) exist. Reapply every 21 days so long as disease symptoms persist.

- Do not apply more than 7 times per season.

Plum and Prune (brown rot blossom blight, brown rot, and rust control)

Mix 1/3 fl. oz. of MONTEREY FUNGI-MAX™ BRAND with 1 gallon of water and uniformly apply to all parts of the tree to point of runoff. For brown rot blossom blight and brown rot, apply when the first green leaf tips are visible and before infection occurs. Reapply every 7 to 14 days so long as disease symptoms persist. For rust, apply MONTEREY FUNGI-MAX™ BRAND about 8 weeks after flowering if conditions favoring disease development (hot, moist weather) exist. Reapply every 21 days so long as disease symptoms persist.

- Do not apply more than 7 times per season.

Strawberry (leaf blight, leaf spot, and powdery mildew)

Mix 1 fl. oz. of MONTEREY FUNGI-MAX™ BRAND with 1 gallon of water and uniformly apply to all parts of the plant to point of runoff. MONTEREY FUNGI-MAX™ BRAND may be applied up to the day of harvest. Apply when the disease first appears or when conditions favoring disease development (hot, moist weather) exist. Reapply every 14 to 21 days so long as disease symptoms persist.

- Do not apply more than 6 times per season.
- Do not plant a new crop in the same place as the crop that was treated within 30 days of the last application.

BERRIES

For all berries, MONTEREY FUNGI-MAX™ BRAND may be applied up to the day of harvest.

Blackberry*, **Raspberry***, **Loganberry** (cane and leaf rust, leaf spot, orange rust, powdery mildew, and yellow rust control)

Mix 1/2 fl. oz. of MONTEREY FUNGI-MAX™ BRAND with 1 gallon of water and uniformly apply to all parts of the plant to point of runoff. Apply at bud break and reapply every 10 to 14 days so long as disease symptoms persist.

- Do not apply more than 4 times per season.

Currant (powdery mildew control)

Mix 1 fl. oz. of MONTEREY FUNGI-MAX™ BRAND with 1 gallon of water and uniformly apply to all parts of the plant to point of runoff. Apply before bloom, through bloom and up to 2 weeks post-bloom.

- Do not apply more than 8 times per season.

Gooseberry (anthracnose and powdery mildew control)

Mix 1 fl. oz. of MONTEREY FUNGI-MAX™ BRAND with 1 gallon of water and uniformly apply to all parts of the plant to point of runoff. For anthracnose, apply when the first leaf has completely unfolded. If conditions favoring disease development (hot, moist weather) exist, reapply every 10 to 14 days. For powdery mildew, apply MONTEREY FUNGI-MAX™ BRAND before bloom, through bloom, and up to 2 weeks post-bloom.

- Do not apply more than 8 times per season.

*Use on the following members of the caneberry group including varieties and/or hybrids: Bingleberry, black satin berry, boysenberry, cherokee blackberry, Chesterberry, Cheyenne blackberry, coryberry, Darrowberry, dewberry, Oregon evergreen berry, phenomenal berry, rangeberry, ravenberry, rossberry Shawnee blackberry, youngberry, Himalayaberry, Dirkson Thornless, hullberry, Lavacaberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, nectarberry

Grapes (anthracnose, black rot and powdery mildew control)

Mix 1 fl. oz. of MONTEREY FUNGI-MAX™ BRAND with one gallon of water and uniformly apply to all parts of the plant to point of runoff. For anthracnose and black rot, apply when new shoots are 1 to 3 inches in length. Reapply every 14 days so long as disease symptoms persist. For powdery mildew, apply MONTEREY FUNGI-MAX™ BRAND when shoots are 12 to 18 inches in length (before bloom). Reapply every 21 days so long as disease symptoms persist.

- Do not apply within 14 days of harvest.
- Do not apply more than 5 times per season.

Peppermint and Spearmint (powdery mildew and rust control)

Mix 1 fl. oz. of MONTEREY FUNGI-MAX™ BRAND with 1 gallon of water and uniformly apply to all parts of the plant to point of runoff. Apply in the early spring when the plants have broken dormancy. Reapply every 14 to 21 days so long as disease symptoms persist.

- Do not apply within 30 days of harvest.
- Do not apply more than 3 times per season.

VEGETABLES

Asparagus (rust control)

Mix 1 fl. oz. of MONTEREY FUNGI-MAX™ BRAND with one gallon of water and uniformly apply to all parts of the plant to point of runoff. Do not apply to harvestable spears. In all locations, apply to the developing ferns after harvest. Reapply every 14 days so long as disease symptoms persist.

In all states, except California:

- MONTEREY FUNGI-MAX™ BRAND may be applied within 180 days of harvest.
- Do not apply more than one time per growing season.

In California only:

- MONTEREY FUNGI-MAX™ BRAND may be applied within 30 days of harvest.
- Do not apply more than four times per year.

Cucurbits (powdery mildew control)

Includes use on the following members of the cucurbits group:

acorn squash	honeydew melon
balsam apple	hubbard squash
balsam pear	mango melon
bitter melon	muskmelon
butternut squash	ornamental gourd
calabaza	persian melon
cantaloupe	pineapple melon
casaba	pumpkin
Chinese cucumber	Santa Claus melon
Chinese waxgourd	scallop squash
citron melon	snake melon
crenshaw melon	spaghetti squash
crookneck squash	straightneck squash
cucumber	summer squash
edible gourd	vegetable marrow
gherkin	watermelon
golden pershaw melon	winter squash
honey balls	zucchini

Mix 1 fl. oz. of MONTEREY FUNGI-MAX™ BRAND with one gallon of water and uniformly apply to all parts of the plant to point of runoff. Apply MONTEREY FUNGI-MAX™ BRAND when the disease first appears. Reapply every 7 to 10 days as long as disease symptoms persist.

- MONTEREY FUNGI-MAX™ BRAND may be applied up to and including the day of harvest.
- Do not apply more than five times per season.
- Do not plant a new crop in the same place as the crop that was treated within thirty days of the last application.

Snap Bean (pod tip rot and rust control)

Mix 1 fl. oz. of MONTEREY FUNGI-MAX™ BRAND with 1 gallon of water and uniformly apply to all parts of the plant to point of runoff.

For rust, apply when the disease first appears. For pod tip rot, apply when pods begin to develop. For both diseases, if conditions favoring disease development (hot, moist weather) exist, reapply every 7 to 10 days.

- MONTEREY FUNGI-MAX™ BRAND may be applied up to and including the day of harvest.
- Do not apply more than 4 times per season.
- Do not plant a new crop in the same place as the crop that was treated within 30 days of the last application.

Tomato (powdery mildew control)

Mix 3/4 fl. oz. of MONTEREY FUNGI-MAX™ BRAND with 1 gallon of water and uniformly apply to all parts of the plant to the point of runoff. Apply when the disease first appears or if conditions favoring disease development (hot, moist weather) exist. Reapply every 21 days so long as disease symptoms persist.

- MONTEREY FUNGI-MAX™ BRAND may be applied up to and including the day of harvest.
- Do not apply more than 4 times per season.
- Do not plant a new crop in the same place as the crop that was treated within 30 days of the last application.

NOTE: This product is non-staining to most home siding, depending on age and cleanliness. However, before using in areas where the spray may contact home siding (vinyl siding in particular) test in an inconspicuous area and recheck in a few hours. Do not use if any staining is observed.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool, dry area away from heat and open flame.

PESTICIDE DISPOSAL: If Partly Filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Monterey
Fungi-Max™
Brand



KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS

CAUTION. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear protective eyewear. Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

See leaflet for complete Precautionary Statements,
Directions for Use & Storage and Disposal

 **WARNING:** Reproductive Harm – www.P65Warnings.ca.gov

Manufactured For:

LAWN AND GARDEN PRODUCTS, INC.

P.O. Box 35000 • Fresno, CA 93745

(559) 499-2100 • info@montereylawnngarden.com

Fungi-Max™ is a trademark
of Lawn and Garden Products, Inc.



NET CONTENTS:
1 PINT/
473.16 mL