


covers **5,000** sq. ft. (50 FT. x 100 FT.)

2217-640-50931_Lawn Weed Control_20170505_4_50931_.pdf



երերերերերերերերերերերերեր PRESS HERE TO RESEAL բերերերերերերերերերերերերերերեր





BROADLEAF WEED CONTROL





• CONTROLS DANDELION, CHICKWEED, PLANTAIN, CLOVERS, WILD ONION **OVER 200 LISTED BROADLEAF WEEDS**

 For use on turf Apply anytime - Spring - Summer - Fall





*2,4-dichlorophenoxyacetic acid.....

INERT INGREDIENTS

***Dicamba: 3,6-dichloro-o-anisic acid



ACTIVE INGREDIENTS:

**(+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid 0.31%



.... 0.13%

... 98.19%



THIS PRODUCT CONTAINS:

Isomer Specific by AOAC Methods

Contains single isomer form of Mecoprop-p.



*27.4 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.

**6.2 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton.

TRIMEC ® is a registered trademark of PBI/Gordon Corporation

***2.6 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.









FOR THE BEST LAWN IN TOWN



9198-AL-001A 9198-OH-2B







EPA REG. NO. 2217-640-50931

Code on bag indicates

manufacturing location.

covers 5,000 SQ. FT.







Distributed by: Jonathan Green, Inc. POB 326 Farmingdale, NJ 07727

Carpetweed

Canada thistle

Burning nettle

Buttonweed

Buttercup

Burweed

Burdock

Burclover

DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTIAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE LISE OF HANDI INE OF THIS PROPILICY HOPEOFFERD. EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY FOR USE ONLY AS DIRECTED. LIMITED WARRANTY AND DISCLAIMER

www.jonathangreen.com Sign up to receive timely lawn tips

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. area out of reach of children and domestic animals. PESTICIDE STORAGE: Store in original container in a dry place. Store in locked Do not contaminate water, food, or feed by storage or disposal. STORAGE AND DISPOSAL



4. Do not wash from weed leaves for 1 to 2 days

sponld be made.

LAWN WEED CONTROL 5M

nroughout the lawn. covers 5,000 sq. ft. of lawn area. To avoid gaps and streaks, first spread border of lawn and then fill in center with uniform, parallel passes. Walk at a steady pace so as to apply evenly Spreading Hints: These granular settings are approximate. Calibrate your spreader on 100 square feet and adjust to apply at the rate of 1 pround of product per 500 square feet. This bag Owner's Manual. Never apply by hand. INTERINCES LUISA OCCUL QUE LO CONQUINOU DI SDESGORI' SDEGO ST MUICU SDINIGO SUO LONGUIUESS DI SLES DESCOL DE SOLE LO CHECK SDESCOL DELOLE DEGLINISMO OCCUL. QUE LO CONQUINIO IL DECESSISTA CONSONI IMPORTANT: It is important to follow the directions as indicated on this quality lawn, care product as over application may cause injury to your lawn. Spreader settings are approximate. True Temper of the border complete each the edge 2byker 9 - b eusnice

product will spreader before 72 - 92 LasyGreen sunting off the ittle overlap of Lawn Pro, SpeedyGreen 1000-2000-3000 9 - 9 he center. A Deluxe, EdgeGuard, Standard, Basic, vorking toward the lawn, α Broadcast EZ butside and of each end of etarting on the Scotts esults, apply in a circular path along the edge Republic EZ 540 porder strips gnaker First, apply 2 For best Prize Lawn Lil' Foot 2-1/2 Vigoro Prize Lawn CBR III gtp Path Precision Greenline (Red Devil) ۷- 9 Scotts PF4 Spreader **2breader** Ortho Scotts Even Green Rotary LESCO (numbered) Scotts AccuGreen 1000-2000-3000 2/L-9 LESCO (lettered) Э Republic EZ 530 K-Gro EZ 540 Precision Greenline (Red Devil) Greenview Jet-Spred SS1 b/L-9 Ortho 13 EV-N-Spread Earthway K-Gro EZ 530 91 Cyclone B1 Greenview Drop-Spred XP1 91 7 - E Central 600 Earthway 15 Agri-Fab 9 Central 550 6 - 8 Hand Crank 2 Bandini Black Beauty Pro-Green Agri-Fab Green-Meter Green-Meter Jonathan Green Jonathan Green 2.0 LBS. PER 1,000 SQ. FT. Spreaders Spreadèrs **SPREADER SETTINGS*** Rotary

and other broadleaf bebberweed Yellowflower Yellow rocket Yarrow Wormseed Woolly plantain Woolly morningglory Woolly croton Woodsorrel Wild vetch Wild sweet potato Wild strawberry Wild rape Wild radish Wild parsnip noino bliW Wild mustard Wild marigold Wild lettuce Wild geranium Wild garlic Wild four-o'clock Wild carrot Wild buckwheat Wild aster White mustard Western saisify Western clematis Wavyleaf bullthistle Virginia pepperweed Virginia creeper Virginia buttonweed Venice mallow Velvetleaf

Tumble pigweed

Tumble mustard

Trailing crownvetch

ISNWGGD Tansy mustard Tansy ragwort Tall vervain Tall nettle lndia mock Strawberry, Stinging nettle Spurweed Spotted spurge Spotted catsear Spiny sowthistle гріпу соскієвиг Spiny amaranth Speedwell Spanishneedles Sowthistle Sorrel Smooth pigweed Smooth dock Smallflower galinsoga Slender plantain Shepherdspurse Sheep sorrel Scotch thistle Scarlet pimpernel Russian thistle Russian pigweed Rough fleabane Rough cinquefoil Redstem filaree Redroot pigweed Red sorrel Ragweed Purslane, common Puncturevine Prostrate vervain Prostrate spurge Prostrate pigweed Prostrate knotweed Prickly sida Prickly lettuce

Prairie sunflower 90j1009 Рокемеед Poison oak Vvi nosio9 Plantain Plains coreopsis Pineywoods bedstraw Pigweed Pepperweed Pennycress Pearlwort Parsnip Parsley-piert Uriental cockiebur Orange hawkweed Nettle Narrowleaf vetch Mustard Musk thistle Mugwort

Narrowleaf plantain Mouseear hawkweed Morningglory Milk vetch Mexicanweed Matchweed Mallow resbegess Lambsquarters Косћіа Knotweed Knawel Deewnosmil Horsenettle Ноагу vervain Hoary plantain Ноагу сгеѕѕ Henbit Hemp Hedge mustard Controls over 200 weeds!

Hedge bindweed Catchweed bedstraw Heath aster Heartleaf drymary Carolina geranium Healall Намкмееd Hairy fleabane рәәмшпр Ground ivy Goldenrod Galinsoga Frenchweed Florida pusley Fiddleneck False sunflower False flax False dandelion Dollarweed Dogfennel Dogbane

Bullnettle Bull thistle **Bulbous** buttercup Buckhorn plantain Buckhorn Broomweed Broadleaf plantain Broadleaf dock Bristly oxtongue Dichondra Brassbuttons Dandelion Bracted plantain **Daisy**, охеуе RIUG VETVAIN Daisy fleabane Blue lettuce Daisy, English Bloodflower milkweed спиу доск BIBSINT DASSAIN baawbuJ Blackseed plantain Creeping Jenny Common mullein Black medic Cocklebur Black-eyed Susan Cockle Bitterweed Clover, white Biffercress, hairy Clover, sweet Siffer wintercress Clover, strawberry Bird vetch Clover, red Bindweed, field Betony, Florida Clover, hop Clover, crimson Beggarticks Cinquefoil Bedstraw Chicory Austrian fieldcress Сһіскwеед, тоиѕееаг Спіскиева, соттоп SWEETCIOVER Catsear Annual yellow

instructions. Never place unused product down any indoor or outdoor drain. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal

> walkways, etc.) One bag covers 5,000 sq. ft. footsge. Deduct any non-lawn areas (house, gardens, after application. At this time a thorough watering of your lawn by its width. This equals your total square

COVERAGE: coverage you need Total Square Feet Non-Lawn Areas Total Square Feet Width (Feet) Length (Feet)

remain on the leaf surface of the weeds. lightly with water to allow the granules to adhere and to If grass is not moist at time of application, sprinkle preferably in the morning when dew is on the grass. 3. Apply when weeds are young and actively growing, to sustain moisture until the next watering (see step 4). 2. Water lawn thoroughly at least 1 day before application 1. Mow lawn to normal height 1 to 2 days before application. The maximum seasonal rate is 4.0 pounds of product per 1,000 sq.ft. number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum broadcast application rate is 2.0 pounds of product per 1,000 sq.ft. per application. The maximum

fescues. Do not apply this product on carpetgrass, dichondra, St. Augustinegrass, or established turf containing Established turf of cool season grasses includes Kentucky bluegrass, perennial ryegrass, tall fescue, and turf-type tall For use on established turf of warm season grasses includes bermudagrass, centipedegrass, and zoysiagrass. and weeds are in a state of poor growth. A second application may be made in 30 days. refreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. Possible INSTRUCTIONS: Jonathan Green Lawn Weed Control controls a wide range of lawn weeds including dandelion, apply when raining or when rain is expected that day.

back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet sprubs, since injury may result. • Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and During application, close spreader openings when turning.

• Apply when the air is calm. Do not apply to newly seeded lawns. Do not exceed specified dosages for any area. and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have • Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people PRECAUTIONS AND RESTRICTIONS: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR USE

groundwater contamination. Application around a cistern or well may result in contamination of drinking water this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of bodies or drainage systems. driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a

waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect **ENVIRONMENTAL HAZARDS** going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556 Have the product container or label with you when calling a poison control center or doctor, or Do not give anything by mouth to an unconscious person. Do not induce vomiting unless told to do so by a poison control center or doctor. swallowed: | • Have person sip a glass of water if able to swallow. Call a poison control center or doctor immediately for treatment advice.

Call a poison control center or doctor for treatment advice. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. If in eyes: Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Call a poison control center or doctor for further treatment advice. inhaled: give artificial respiration, preferable mouth-to-mouth, if possible. Move person to fresh air. If person is not breathing, call 911 or ambulance, then · Call a poison control center or doctor for treatment advice. or clothing: • Rinse skin immediately with plenty of water for 15 - 20 minutes. If on skin | • Take off contaminated clothing. **GIA T2RI7**

before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear long-sleeved shirt and long Avoid breathing dust. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and CAUTION - Harmful if absorbed through skin. Harmful if inhaled. Avoid contact with skin, eyes, or clothing. HAZARDS TO HUMANS AND DOMESTIC ANIMALS PRECAUTIONARY STATEMENTS

pants, socks, shoes, and gloves. Remove and wash contaminated clothing before reuse.

PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY. READ THE ENTIRE LABEL FIRST. OBSERVE ALL













Gatnip



Aster

MEEDS







