

Cut or Tear here to open

Jonathan

Green

Easy-to-use! Resealable bag!

Jonathan Green

LAWN DISEASE CONTROL SPRING, SUMMER, FALL

ALSO PROVIDES BROAD SPECTRUM SYSTEMIC CONTROL OF SOIL BORNE DISEASES

MAY BE USED ON **COOL AND WARM SEASON GRASSES**

Jonathan Green

LARGE BROWN PATCH LEAF SPOT DOLLAR SPOT RED THREAD


7.5 LBS. NET WEIGHT (3.40 Kg)

ACTIVE INGREDIENT: 97.92%

CAUTION: KEEP OUT OF REACH OF CHILDREN 100.00% EPA Reg. No. 53883-183-50931 EPA Est. No. 075527-PA-12



FUNGUS CONTROL



Spreader settings given are approximate and variation can occur between identical models. User should calibrate spreader by accurately weighing a few pounds into your spreader and applying to a measured area. Then adjust spreader setting if necessary to assure correct rate of application.

For Treatment: Apply at 14-day intervals when disease first appears. Rotations may be used

as bart of the three-application program.

SPREADER SETTINGS

өрод әциодриврүү

For Prevention: Using a 21-day reapplication interval, make three applications starting in Summer Patch For Prevention: Apply in spring and fall. For Treatment: Make two applications, the first when the disease first appears and the second 14-days later. simnotiints ogolitsU tum2 aqint2 Root Disease Bipolaris, Curvularia, and Exserohilum Leaf, Crown, and Drechslera spp. NOTE: For severe cases, rotations with cholorothalonil, iprodione, or mancozeb (ProtectTM) Leaf Spot Apply when disease first appears and make applications at 14-day intervals as needed. Рупсиаліа дпѕеа Continue applications at 14-day intervals as needed. Make a preventative application prior to the expected onset of disease development. Gray Leaf Spot (Blast) Fusarium Blight mivisorm and f. triticum Apply when disease first appears. Make repeat applications at 14-day intervals. Make two treatments at 21-day intervals in late fall to early winter. Make the last application just prior to first permanent snow cover. bloM won2 suningo2 appears and repeat applications every 14-days. Spring Dead Spot $55-60^{\circ}F_{\nu}$ whichever occurs first. For Treatment: Apply in early Spring when disease first Mecrotic Ring Spot application in early Spring when disease first appears or when soil temperatures reach For Prevention: Apply in the fall prior to cessation of turt growth. Make a second Cool Season Brown Patch 5 - 2.I Баеитаппотусеѕ дгатіпіѕ чаг. аачепае agronomic instructions to maintain plant vigor. baeumannomyces graminis var. graminis Take-All-Patch suppression. Use higher rates under most severe disease expression. Follow proper Apply in mid-July or when disease symptoms first appear and repeat at 14-day intervals for **Bermudagrass Decline** Anthracnose, foliar Colletotrichum graminicola Anthracnose, basal Alletotrichum graminicola intervals. For Treatment: Apply at first sign of disease and continue applications at 14-day intervals. NOTE: May be rotated with chlorothalonil or triadimefon. For Prevention: Once soil temperature reaches 60°F, make two applications at 14-day Throughout the expected disease period, continue making applications at 14-day intervals. For best results, make the first application before the turf is completely dormant. Pink Snow Mold
Michrodochium nivale Make first application prior to cessation of turt growth activity in middle-late Fall or early Winter, or in Spring during periods of thaw, or when disease symptoms first appear. hovin minisut Astad bizyoZ

Gloeocercospora sorghi

Red Thread

Loeitsaria fuciformis

Tusarium Patch

Fusarium ivinle Copper Spot Ascochyta Leaf Blight 14-days. NOTE: May be rotated with chlorothalonil, iprodione, or mancozeb (ProtectTM). ratde Brown Patch Pollar Spot Moellerodiscus, Lanzia, Sclerotinia Patch listed. For Treatment: Apply when disease first appears and repeat applications every For Prevention: Apply prior to the expected development of the disease using the lowest rate Application Rate (Lbs./1000 sq. ft.) Disease Specific Instructions Refer to the following table for specific rates and timing depending on the disease being treated. When treating under severe disease pressure, use the highest recommended rate listed below.

for any and all losses, injuries, or damage resulting from or in any way arising from the use, handling, or application of this product, whether in contract, warranty, tort, negligence, strict liability, or otherwise, shall be damages not exceeding the purchase price paid for this product or, at Jonathan Green, Inc. election, the replacement of this product. Except, as expressly provided herein, Jonathan Green, Inc. makes no warranties, guaranties, or representations of any kind, either expressed or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage. The exclusive remedy of any buyer or user of this product a particular purpose, use or eligibility of the product for any particular product. may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Jonathan Green, Inc. To the extent consistent with applicable law, Jonathan Green, Inc. shall in no event be liable for consequential, special, or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences The following limitations of warranty shall apply to the extent that they are consistent with applicable law. Jonathan Green, Inc. warrants that following limitations of warranty shall apply to the extent that the characteristics on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. This product is not guaranteed to be compatible with any other pesticides. It is accordance with all sicks is because the compatible with any other pesticides. It is

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain. If empty: Place in trash or offer for recycling if available. Pesticide Disposal: Nonrefillable container. Do not reuse or refill this container. Pesticide Storage: Store in original container in a dry place. Store in locked area out of reach of children and domestic animals. STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. pressure, use the highest specified rate listed to the right. Refer to the following table for specific rates and timing depending on the disease being treated. When treating under severe disease

• For best results, activate the application within 24 hours with irrigation or natural rainfall. If activating by irrigation, apply 0.10 - 0.25 Apply under dry or moist leaf/soil conditions. **ENOITSUI NOITAZIJ99A** this label, Jonathan Green Lawn Fungus Control is not phytotoxic to any of the above-mentioned grasses. Jonathan Green Lawn Fungus Control may be used on all cool and warm season grasses such as Bentgrass, Bluegrass, Bermudagrass, Prescue, Ryegrass, St. Augustinegrass, Coysiagrass, or their mixtures in residential turf sites. When used in accordance with the instructions in this lebel to a state of the control to the cont

HOMEOWNER TURF APPLICATIONS \bullet Do not apply application closer than fourteen (14) days from previous treatment. • DO NOT exceed a maximum of twelve (12) pounds of product per 1000 square feet per season from any thiophanate-methyl • DO NOT exceed a maximum of three (3) pounds of product per 1000 square feet per application

MAXIMUM APPLICATION RATES AND MINIMUM RETREATMENT INTERVALS identification and advice on other suitable fungicide or disease control strategies. be ineffective and you should promptly contact your State Agricultural Experiment Station or State Cooperative Extension Service for disease specified precautions are followed. A tolerant strain of fungi may be present if a Jonathan Green Lawn Fungus Control treatment is found to contribute to the development of disease tolerance. Jonathan Green Lawn Fungus Control can remain useful for disease control as long as In order to prevent the development of tolerant strains of fungi, Jonathan Green Lawn Fungus Control should only be used with other fungus control with products contraining the active ingredient thiabendazole because thiabendazole has a similar chemistry and mode of action and can products contraining the active ingredient thiabendazole because thiabendazole has a similar chemistry and mode of action and can

 \bullet This product is not guaranteed to be compatible with any other pesticides. • Thoroughly clean all application equipment after each use. • Use only properly calibrated application equipment when applying this product. Do not feed clippings to livestock or poultry.

 Do not graze animals on treated turf. • Do not apply by hand or with handheld rotary spreader (e.g. belly grinder) \bullet Do not apply this product through any type of irrigation system. Jonathan Green Lawn Fungus Control is a broad-spectrum fungicide for use on a wide variety of turf diseases. Jonathan Green Lawn Fungus Control may be applied using ground based, hand-operated (push) or mechanical (power) equipment; do not apply this product by hand or with hand-held rotary spreader (e.g., bellygrinder). If a rate range is given, use the higher rate under conditions of severe disease pressure. Contact your State Extension service for recommendations for application schedules appropriate to your area.

Call a poison control center or doctor immediately for treatment advice.

 Mave person sip a glass of water if able to swallow.
 Do not induce vomitting unless told to do so by the poison control center or doctor.

 Call a poison control center or doctor for treatment advice. If in eyes

• Hold eye open and rinse slowly and gently with water for 15-20 minutes.

• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

GENERAL INFORMATION HOMEOWNER TURE APPLICATIONS contact any person or pet. Keep people and pets out of the area during application. Do not enter or allow others to enter the treated areas until It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will directly

Do not apply directly to water or areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wear long-sleeved shirt, long pants, and shoes with socks. ИОІТИАЭ

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS For spill or transportation emergencies call CHEMTREC (1-800-424-9300). Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall (866) 897-8050 for emergency medical treatment information. HOT LINE NUMBER If on skin • Take off contaminated clothing.

or clothing • Rinse skin immediately with plenty of water for 15-20 minutes.

• Call a poison control center or doctor for treatment advice. • Call a poison control center or doctor for further treatment advice. If inhaled

• Move person to fresh air.

• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible. Do not give anything by mouth to an unconscious person.

