

LAWN WEED CONTROL

Kills Dandelions, Clovers and Other Major Lawn Weeds

Kills 250+ Weeds

Kills the Weeds, Not the Lawn Covers up to 5,000 square feet







Net Wt. 14 lbs (6.35 kg)



LAWN WEED CONTROL 250



KEEP OUT OF REACH OF CHILDREN CAUTION SEE BACK PANEL FOR FURTHER PRECAUTIONS, FIRST AID INFORMATION AND ANALYSIS DESCRIPTION

Net Wt. 14 lbs (6.35 kg)

ì	· · · · · · · · · · · · · · · · · · ·	ada i
報	ACTIVE INGREDIENTS:	a deligi
	2, 4-D, 2-ethylhexyl ester	. 1,108
\$1 31	Mecoprop-p acid	. 0.167
	Dicamba acid	
复复	OTHER INGREDIENTS	98,654
医溶液	TOTAL	100,000
	THIS PRODUCT CONTAINS:	and the
ij	14.70 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.735%	
ě	E CAS 1928-43-4.	
800年	3.34 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.167%	i Le i
	CAS 16484-77-8	1
9	1.42 lb 3, 6-dichloro-o-anisic acid equivalent per ton or 0.071%.	
i di	CAS 1918-00-9.	
級	Isomer Specific by AOAC Methods	BASE LIE



AWN WEED CONTROL

Kills
250+



PERT LAWN WEED CONTROL 250+

Kills



STOP READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap a water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For informa-tion on this product, contact the National Pesticide Information Center, 1-800-828-7378, Monday - Friday, 8:00 am to 12:00 pm PST. You may also contact the National Poison Control Center, 1-800-222-1222, day or night, for emergency medical treatment information. FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

ENVIRONMENTAL HAZARDS

perticled is toxic to fish and aqualic invertebrates and may adversely affect non-target plants. To protect the emironment, do not allow pasticide to enter or run off into storm ins. drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does tallow or wash pesticide off the returnent area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to

prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

 Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production or for research purposes.

This product is a combination of three proven weed killers—2,4-D, mecoprop-p, and dicamba. This herbicide controls a wide range of lawn weeds, including Dandelion, Chickweed, Knotweed, Plantain, Henbit and Spurge.

USE PRECAUTIONS

Do not use on Dichondra, Carpetgrass, 'Floratam' St. Augustinegrass, Creeping Bentgrass or Turfgrass where desirable clovers are present. Do not apply this product to newly seeded grasses until well established or after the second or third mowing. Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees.

Do not exceed specified dosage for any area and do not apply this product within the drip line of, or over surface roots of trees, shrubs,

and other ornamentals.

Reseeding or overseeding areas may occur three (3) weeks after the application of this product.

Do not use this product on or near desirable plants and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of the roots of desirable trees and shrubs, since injury may result.

uniconies) within the unip line or the roots of desirable trees and shrubs, since injury may result.

Do not apply directly to or near water, storm drains, gutters, sewers or drainage directles. Do not apply within 25 feet of rivers, fishponds, lakes, streams, reservoirs, marshes, estuaries, bays and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. For prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

Which CAPPLY
Apply when the air is calm to ensure uniform coverage and avoid spreading granules onto flowers, vegetables and ornamental shrubbery.
Possible retreatment may be necessary if lawn is heavily overrun by weeds, if adverse low moisture conditions prevail or if weeds are in a state of poor growth.

HOW MUCH TO APPLY

For use on Kentucky Bluegrass, Fescue spp., Perennial Ryegrass, common Bermudagrass, Bahiagrass, Zoysiagrass, St. Augustinegrass, Centipedegrass, hybrid Bermudagrass, and Colonial Bentgrass. Apply 2.8 pounds of product per 1,000 square feet of furfgrass. A second application may be made in approximately 30 days. The maximum number of broadcast applications per treatment site is 2 per year.

APPLICATION INSTRUCTIONS

- Mow lawn to normal height 1 to 2 days before application.
 Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).
 Apply when Broadleaf weeds are young and actively growing, preferably in the morning when dew is on the grass. At the time of application, moisture on the weed leaves from dew, rainfall or irrigation/watering aids in control. If grass is not moist at time of
- application, sprinkle lightly with water before applying product.

 4. Avoid applying when temperatures are consistently above 90°F.

 5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

SPREADER SETTIN	IGS
BROADCAST SPREAD	ER
Scotts* SpeedyGreen*	3.75
Scotts* Basic	3.75
Scotts* Lawn Pro	3.75
Scotts* EdgeGuard* Deluxe	3.75
Scotts* EdgeGuard* Mini	3.75
DROP SPREADER	

Scotts* Accugreen*

IOW TO APPLY - SPREADERS & CALIBRATION

lways apply with a rotary or drop spreader to distribute the granules evenly. Never apply by

OVEXAGE: to determine now mucroy to need to apply, intuliply to register by our events, swidth. This equals your total square footage. Deduct all non-lawn areas (house, gardens tc.). One bag covers up to 5,000 square feet.

MPORTANT: It is important to follow directions as indicated on this quality lawn care proc

; overapplication may cause injury to your lawn. Spreader settings are approximate. ifferences may occur due to condition of spreader, speed at which applied and roughness of

- Ifferences may occur due to condition of spreader, speed a twinch applied and roughness rea treated, Be sure to check spreader before beginning, adjusting if necessary. Consult Nemer's Manual. Never apply by hand.
 For rotary spreaders: Operate at intervals corresponding to the swath width.
 For drop spreaders: Operate at intervals corresponding to the swath width.
 For drop spreaders: Overlap the wheel tracks to ensure complete coverage.
 For best results, apply this product uniformly and avoid streaks, kighs or overlaps. Walk at a steady pace to ensure even coverage. Close the spreader opening when turning to change distinctions or when stronging.

Expert Gardener® Lawn Weed Control 0-0-1

GUARANTEED ANALYSIS
Soluble Potash (K,O)1.0%
Derived from: Muriate of Potash.

CALIBRATION INSTRUCTIONS AND SPREADER SETTINGS

Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended rate under actual application conditions.

To provide proper distribution, calibrate spreader before application.

BROADCAST SPREADERS: For best results, apply in a circular path starting on the outside and working toward the center

overlaping slightly to ensure complete coverage. The apply along length of lawn, overlapping slightly to ensure complete coverage. Then apply along length of lawn, overlapping slightly and shutting off the spreader before each turn at edge of border strips.

WEEDS CONTROLLED

Aster, heath Aster, white heath Aster, purple Aster, white prairie Artichoke Austrian fieldcress Betony, Florida Bindweed, Field Black-eyed Susan Blue lettuce Bristly oxtongu Broomwe Buckhern Builnettle Burweed, lawn (spurweed) Buttercup, bulbous Catsear, spotted (false dandelion) Chickweed, common

Cocklebur, oriental Cocklebur, spiny Dock broadleaf Mugwort Mustard, ball Mustard, black Mustard, black Mustard, blue Mustard, hedge Mustard, Indian, Mustard, tansy Mustard, tumble Mustard, white False sunflower Fiddleneck Filaree, redstem Filaree, whitesten Reabane, daisy (a Mustard, wormseed Heabane, rough Galinsoga, hairy Galinsoga, smallflower Nettle, stinging (burning) Parsnip Pearlwort Pennycress (frenchweed) Pepperweed, perennia Hawkweed, yellow Pigweed, russian Pigweed, smooth Pigweed, tumble Pineapple weed

Plantain, woolly Puncturevine Pusley brazil Pusley, brazil
Pusley, florida
Pusley, large flower
Ragweed, bur
Ragweed, commor
Ragweed, landeaf
Ragweed, western
Redstem filaree Rough cinquefoil Sorrel, red (sheep) Speedwell, bi-lobe Speedwell, common Speedwell, com Speedwell, purstane Speedwell, slender Speedwell, snow Speedwell, snow Speedwell, thymeleaf Spiny amaranth Spurge, spotted Spurge, prostrate Strawberry, India mock Thistle, blessed Thistle, bull Thistle, canada Thistle, flodman (prairie) Thistle, platte Inistle, platte
Thistle, plumeless
Thistle, russian
Thistle, scotch
Thistle, wavyleaf
Thistle, yellow (hor
Trailling crownvetch
Velvetleaf
Venoric blue Vervain, blue Vervain, hoary Vetch, bird Vetch, hairy Vetch, milk Vetch, narro Vetch, wild Virginia creeper Western clematis Western salsify Wild aster Wild buckweat Wild carrot Wild four-o-dock Wild marigolo Wild mustard (yellow rocket) Wild parsnip Wild radish Wild sweet potato Wild violet Woodsorrel, violet Woodsorrel, vellow (Oxalis) Woolly croto Wormseed Yarrow, commo Yarrow, wester

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container: Do not reuse or refill this container. If empty: Place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place any unused product down any indoor or outdoor drain.

Keep this product off sidewalks, patios, driveways, etc. This fertilizer may stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders, which scatter fertilizer beyond lawn edge and onto concrete.

F525

EPA Est. No. 59144-GA-1 (A), 32802-MO-1 (B), 37351-CO-1(C), 8378-IN-1 (D), 9404-FL-1 (E), 961-PA-01 (F), 961-IL-01(G), 35497-OR-1 (H), 49991-TX-2 (Q), 69759-OR-002 (R), 7401-TX-01 (S)

When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials, Information regarding the contents and levels of metals in this product is available on the Internet at http://www.aapfco.org/metals.htm

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSIV DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.



green, EdgeGuard and Scotts are register trademarks of OMS Investments, Inc. peedyGreen is a registered trademark of Everris International B.V.

Distributed by Gro Tec, Inc. P O. Box 290, Madison, GA 30650



LAWN WEED CONTROL 250

Burdock

Kills



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

If in Eyes

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this product, contact the National Pesticide Information Center, 1-800-858-7378, Monday - Friday, 8:00 am to 12:00 pm PST. You may also contact the National Poison Control Center, 1-800-222-1222, day or night, for emergency medical treatment Information. FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production or for research purposes.

This product is a combination of three proven weed killers—2,4-D, mecoprop-p, and dicamba. This herbicide controls a wide range of lawn weeds, including Dandelion, Chickweed, Knotweed, Plantain, Henbit and Spurge.

USE PRECAUTIONS

Do not use on Dichondra, Carpetgrass, 'Floratam' St. Augustinegrass, Creeping Bentgrass or Turfgrass where desirable clovers are present. Do not apply this product to newly seeded grasses until well established or after the second or third mowing.

CALIBRATION INSTRUCTIONS AND SPREADER SETTINGS

Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended rate under actual application conditions

To provide proper distribution, calibrate spreader before application.

BROADCAST SPREADERS: For best results, apply in a circular path starting on the outside and working toward the center overlapping slightly to ensure complete coverage.

DROP SPREADERS: Apply 2 strips along each end. Then apply along length of lawn, overlapping slightly and shutting off the spreader before each turn at edge of border strips.

WEEDS CONTROLLED

Clover, vellow sweet Horsenettle Amaranth, purple Clover, white sweet Amaranth slender Horseweed Aster, heath Clover, white limsonweed Cockle Johnny-Jumpup violet Aster, white heath Aster numle Cocklebur Cocklebur, oriental Knotweed, prostrate Aster, white orairie Artichoke Cocklebur, spiny Kochia Austrian fieldcress Common mulleir Lambsquarters Bedstraw, Catchweed Corn Spurry Lespedeza Redstraw smooth Creeping Beggarweed Mallow, alkali Mallow, bristly Creeping Jenny (creeping charlie) Bedstraw, Northern Bedstraw, pineywoods Mallow, common **Cupid shaving Brush** Beggartricks, hairy Mallow, dwarf Cudweed Mallow, venice Betony, Florida Daisy, English Bindweed, Field Marcela Daisy, oxeye Bindweed, Hedge Matchweed Dandelion, common **Bitter Wintercress** Mexicanweed Dandelion false Bittercress, hairy Moneywort Day flower Bitterweed Morningglory, bigroot Deadnettle Black-eyed Susan Morningglory, ivyleaf Dichondra Morningglory, Mexican Black Medic Dock, broadleaf Blood flower milkweed Morningglory, red Dock, curly Morningglory, tall Blue lettuce Dogbane Brassbuttons Morningglory, woolly Dogfennel Mugwort Bristly oxtonau Dollarweed (Pennywort) Mustard, ball Broomweed Dove weed Mustard, black Buckhorn Elderberry Bullnettle Mustard, blue Falseflax Burclover Mustard, hedge False sunflower Mustard, Indian. Fiddleneck Burweed, lawn (spurweed) Mustard, tansy Filaree, redstem Buttercup, bulbous Mustard, tumble

Plantain, blackseed Strawberry, India mock Plantain, bracted Plantain broadleaf Plantain, buckhorn Plantain, hoary Plantain narrowleaf Plantain, slender Plantain, woolly Poison ivy Poison oak Pokeweed Prairie Sunflower Prickly lettuce (compass plant) Prickly sida Prostrate knotweed **Puncturevine** Purslane, common Pusley, brazil Pusley, florida Pusley, large flower Ragweed, bur Ragweed, commor Ragweed, lancleaf Ragweed, western Redstem filaree Rough cinquefoil Scarlet pimpernel Shenberd's purse Smartweed, ladysthumb Smartweed, pale

Smartweed, pennsylvania

Smooth chaff-flower

Tansy ragwort Tanweed Thistle, blessed Thistle, bull Thistle, canada Thistle, flodman (prairie) Thistle, musk Thistle, platte Thistle, plumeless Thistle russian Thistle, scotch Thistle, wavyleaf Thistle, yellow (horrible) Trailing crownvetch Velvetleaf Vervain, blue Vervain, hoary Vervain, prostrate Vervain, tall Vetch, bird Vetch, hairy Vetch, milk Vetch, narrowlead Vetch, wild Virginia creeper Western clematis Western salslfv Wildaster Wild buckweat Wild carrot Wild four-o-clock

Do not use on Dichondra, Carpetgrass, 'Floratam' St. Augustinegrass, Creeping Bentgrass or Turfgrass where desirable clovers are present. Do not apply this product to newly seeded grasses until well established or after the second or third mowing.

Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees.

Do not exceed specified dosage for any area and do not apply this product within the drip line of, or over surface roots of trees, shrubs, and other ornamentals.

Reseeding or overseeding areas may occur three (3) weeks after the application of this product.

Do not use this product on or near desirable plants and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of the roots of desirable trees and shrubs, since injury may result.

Do not apply directly to or near water, storm drains, gutters, sewers or drainage ditches. Do not apply within 25 feet of rivers, fishponds, lakes, streams, reservoirs, marshes, estuaries, bays and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

WHEN TO APPLY

Apply when Broadleaf weeds are actively growing in Spring or early Fall.

Apply when the air is calm to ensure uniform coverage and avoid spreading granules onto flowers, vegetables and ornamental shrubbery, Possible retreatment may be necessary if lawn is heavily overrun by weeds, if adverse low moisture conditions prevail or if weeds are in a state of poor growth.

HOW MUCH TO APPLY

For use on Kentucky Bluegrass, Fescue spp., Perennial Ryegrass, common Bermudagrass, Bahiagrass, Zoysiagrass, St. Augustinegrass, Centipedegrass, hybrid Bermudagrass, and Colonial Bentgrass. Apply 2.8 pounds of product per 1,000 square feet of turfgrass. A second application may be made in approximately 30 days. The maximum number of broadcast applications per treatment site is 2 per year.

APPLICATION INSTRUCTIONS

- 1. Mow lawn to normal height 1 to 2 days before application.
- 2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).
- 3. Apply when Broadleaf weeds are young and actively growing, preferably in the morning when dew is on the grass. At the time of application, moisture on the weed leaves from dew, rainfall or irrigation/watering aids in control. If grass is not moist at time of application, sprinkle lightly with water before applying product.
- 4. Avoid applying when temperatures are consistently above 90°F.
- 5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

SPREADER SET III	103		
BROADCAST SPREADER			
Scotts* SpeedyGreen*	3.75		
Scotts* Basic	3.75		
Scotts* Lawn Pro	3.75		
Scotts* EdgeGuard* Deluxe	3.75		
Scotts* EdgeGuard* Mini	3.75		
DROP SPREADER			

Scotts® Accugreen®

SPREADER SETTINGS HOW TO APPLY - SPREADERS & CALIBRATION

Always apply with a rotary or drop spreader to distribute the granules evenly. Never apply by

COVERAGE: To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to 5,000 square feet.

IMPORTANT: It is important to follow directions as indicated on this quality lawn care product as overapplication may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult Owner's Manual. Never apply by hand.

- · For rotary spreaders: Operate at intervals corresponding to the swath width.
- · For drop spreaders: Overlap the wheel tracks to ensure complete coverage.
- For best results, apply this product uniformly and avoid streaks, skips or overlaps. Walk at a steady pace to ensure even coverage. Close the spreader opening when turning to change directions or when stopping.

Expert Gardener® Lawn Weed Control 0-0-1

F525

.1.0%

GUARANTEED ANALYSIS

Soluble Potash (K.O).. Derived from: Muriate of Potash.

mustara, ingian, Smartweed, pale wild back wear Fiddleneck Burweed, Jawn (spurweed) Mustard, tansv Smartweed, pennsylvania Wild carrot Filaree, redstem Buttercup, bulbous Mustard tumble Smooth chaff-flowe Wild four-o-clock Filaree, whitestern Buttercup, creeping Mustard, white Smoothdock Wild garlic Fleabane, daisy (annual) Buttonweed, common (poorjoe) Mustard, wormseed Sorrel, red (sheep) Wildlettuce Fleabane, rough Buttonweed, Virginia Nettle, stinging (burning) Sowthistle, annual Wild marigold Galinsoga, hairy Carolina geranium (wild) Sowthistle, spiny Nettle, tall Wild mustard (vellow rocket) Galinsoga, smallflower Carnetweed Nettle wood Spanish needles Wild onion Goldenrod Catnip Parsley-piert Speedwell, bi-lobed Wild parsnip Ground ivv Catsear, spotted (false dandelion) Parsnip Speedwell, common Wild radish Groundsel Chamber bitter Pearlwort Speedwell, corn Wild rape Chickweed, common Gumweed Pennycress (frenchweed) Speedwell ivyleaf Wild strawberry Hairy fleabane Chickweed, mouseear Pepperweed, perennial Speedwell, persian Wild sweet potato Hawkweed, meadow Chickweed, sticky Pepperweed, virginia (Peppergrass) Speedwell, purslane Wild violet Hawkweed, mouseear Chicory Speedwell, slender Woodsorrel, creeping Hawkweed, orange Cinquefoil Pigweed, prostrate Speedwell, snow Woodsorrel violet Hawkweed, yellow Pigweed, redroot Clover, crimson Speedwell thymelea Woodsorrel, yellow (Oxalis) Healall Pigweed, russian Clover, alvce Spiny amaranth Woolly croton Heartleaf drymary Pigweed, smooth Clover, hop Spurge, spotted Wormseed Clover, rabbit foot Hemo Pigweed, tumble Spurge, prostrate Yarrow, common Henbit Pineapple weed Clover, red Spurweed Yarrow, western Hoary cress Plains coreopsis (tickseed) Clover, strawberry Star of Bethlehem Yarrow, yellow

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container: Do not reuse or refill this container. If empty: Place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place any unused product down any indoor or outdoor drain.

Keep this product off sidewalks, patios, driveways, etc. This fertilizer may stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders, which scatter fertilizer beyond lawn edge and onto concrete.

EPA Reg. No. 2217-884-59144

EPA Est. No, 59144-GA-1 (A), 32802-MO-1 (B), 37351-CO-1(C), 8378-IN-1 (D), 9404-FL-1 (E), 961-PA-01 (F), 961-IL-01(G), 35497-OR-1 (H), 49991-TX-2 (O), 69759-OR-002 (R), 7401-TX-01 (S) Letter in parenthesis is first letter of lot number

When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials. Information regarding the contents and levels of metals in this product is available on the Internet at http://www.aapfco.org/metals.htm

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL. CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.



Accugreen, EdgeGuard and Scotts are registered trademarks of OMS Investments, Inc. SpeedyGreen is a registered trademark of

> Distributed by Gro Tec, Inc. P O. Box 290, Madison, GA 30650

Expert Gardener® is a registered trademark of Wal-Mart Stores, Inc., Bentonville, AR 72716