WEED&FEED...

Kills Dandelions, Clovers and Other Major Lawn Weeds

Kills the Weeds, Not the Lawn

Covers up to 5,000 Square Feet

28-0-3



KEEP OUT OF REACH OF CHILDREN CAUTION

13 lbs (5.90 kg)

14.70 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.735%

3.34 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.167%. 1.42 lb 3, 6-dichloro-o-anisic acid equivalent per ton or 0.071%. CAS 1918-00-9. Isomer Specific by AOAC Methods



WEED & FEED.

28-0-3



verbai statements or agreements at the point of sale or elsewhere. Tull. The terms of this Limited Warranty and Discialmer cannot be varied by any written or nuobened immediately to the point of purchase, and the purchase price will be refunded in HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPL MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF Accugreen, EdgeGuard and Scotts are registered trademarks of OMS Investments, Inc. FOR USE ONLY AS DIRECTED. THE MANUFACTURER NEITHER MAKES NOR INTENDS LIMITED WARRANTY AND DISCLAIMER

and levels of metals in this product is available on the Internet at Association of American Plant Food Control Officials. Information regarding the contents When applied as directed, this product meets the guidelines for metals adopted by the $\,$

Letter in parenthesis is first letter of lot number. 961-PA-01 (F), 961-IL-01(G), 35497-OR-1 (H), 49991-TX-2 (Q), 69759-OR-002 (R), 7401-TX-01 (S)

EPA Est. No, 59144-GA-1 (A), 32802-MO-1 (B), 37351-CO-1(C), 8378-IN-1 (D), 9404-FL-1 (E), practices. Check with your county or city government to determine if there are local regulations for fertilizer use.

your lawn. Check with your local Cooperative Extension Agency to obtain specific information on local turf best management exbected. Apply this product only to your lawn, and sweep any product that lands in the driveway, sidewalk, or street, back onto Apply only to actively growing turf. Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is impossible to prevent. Be careful when using rotary spreaders, which scatter fertilizer beyond lawn edge and onto concrete.

with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be Keep this product off sidewalks, patios, driveways, etc. This fertilizer may stain concrete and painted surfaces. If it comes in contact

recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place any unused product AND CONTAINER HANDLING: Nonvefillable container: Do not reuse or refill this container. If empty: Place in trash or offer for bags containing unused product in a locked storage area inaccessible to children and domestic animals. PESTICIDE DISPOSAL Do not contaminate water, food or feed by storage and disposal. PESTICIDE STORAGE: Store in original container and keep any

			STORAGE AND DISPOSAL	
Yarrow, yellow	Star of Bethlehem	Plains coreopsis (tickseed)	Ноагу сгеss	Сіочет, strawberту
Yarrow, western	pəəmındç	Pineapple weed	Henbit	Clover, red
Хаггом, соттоп	Spurge, prostrate	Pigweed, tumble	Нетр	Clover, rabbit foot
Wormseed	Spurge, spotted	Pigweed, smooth	Heartleaf drymary	СІолет, hop
Woolly croton	Spiny amaranth	Pigweed, russian	Heabill Action of the second o	СІолет, аlусе
Woodsorrel, yellow (Oxalis)	Speedwell, thymeleaf	Pigweed, redroot	Hawkweed, yellow	Clover, crimson
Woodsorrel, violet	speedwell, snow	Pigweed, prostrate	Намкмеед, отапде	Cinquefoil
Woodsorrel, creeping	Speedwell, slender	(Peppergrass)	Hawkweed, mouseear	Сһісогу
J9loiv bliW	Speedwell, purslane	Pepperweed, virginia	Намкмеед, теадом	Chickweed, sticky
Wild sweet potato	Speedwell, persian	Pepperweed, perennial	Hairy fleabane	Chickweed, mouseear
Wild strawberry	Speedwell, ivyleaf	Pennycress (frenchweed)	beewmue	Сһіскмеед, соттоп
Wild rape	Speedwell, corn	Pearlwort	Groundsel	Chamber bitter
Wild radish	Speedwell, common	Parsnip	Ground ivy	Catsear, spotted (false dandelion)
Wild parsnip	Speedwell, bi-lobed	Parsley-piert	Goldenrod	qinteO
noino bliW	Splanish needles	Nettle, wood	Galinsoga, smallflower	Carpetweed
Wild mustard (yellow rocket	Sowthistle, spiny	Nettle, tall	Galinsoga, hairy	Carolina geranium (wild)
bloginsm bliW	Sowthistle, annual	Nettle, stinging (burning)	Fleabane, rough	Buttonweed, Virginia
Wild lettuce	Sorrel, red (sheep)	Mustard, wormseed	Fleabane, daisy (annual)	Buttonweed, common (poorjoe)
Wild garlic	гиоограск	Mustard, white	Filaree, whitestem	Buttercup, creeping
Wild four-o-clock	Smooth chaff-flower	Mustard, tumble	Filaree, redstem	Buttercup, bulbous
Wild carrot	Smartweed, pennsylvania	Mustard, tansy	Fiddleneck	Burweed, lawn (spurweed)
Wild buckweat	Smartweed, pale	Mustard, Indian,	False sunflower	Burdock
Wild aster	Smartweed, ladysthumb	Mustard, hedge	Falseflax	Burclover
Western salsify	Shepherd's purse	Mustard, blue	Elderberry	Bullnettle
Western dematis	Scarlet pimpernel	Mustard, black	Dove weed	впскрош
Virginia creeper	Rough cinquefoil	Mustard, ball	Dollarweed (Pennywort)	Broomweed
Vetch, wild	Redstem filaree	Мидумог	Dogfennel	Bristly oxtongue
Vetch, narrowleaf	Ragweed, western	Morningglory, woolly	Dogbane	Brassbuttons
Vetch, milk	Ragweed, lancleaf	Morningglory, tall	роск, сипу	Blue lettuce
Уетсћ, ћаігу	увадмеед, соттоп	МогпіпддІогу, rèd	Dock, broadleaf	Blood flower milkweed
Vetch, bird	Ragweed, bur	Morningglory, Mexican	Dichondra	Black Medic
Vervain, tall	Pusley, large flower	Morningglory, ivyleaf	Deadnettle	gJack-eyed Susan
Vervain, prostrate	Pusley, florida	Morningglory, bigroot	Day flower	Bitterweed
Vervain, hoary	Pusley, brazil	Moneywort	Solat ,noilebned	Bittercress, hairy
Vervain, blue	Purslane, common	Mexicanweed	Dandelion, common	Bitter Wintercress
Velvetleaf	Puncturevine	Matchweed	Daisy, oxeye	Bindweed, Hedge
Trailing crownvetch	Prostrate knotweed	Marcela	Daisy, English	Bindweed, Field
Thistle, yellow (horrible)	Prickly sida	Asiney, welleM	pəəmpn	Betony, Florida
Thistle, wavyleaf	Prickly lettuce (compass plant)	Mallow, dwarf	Cupid shaving Brush	Beggartricks, hairy
Thistle, scotch	Prairie Sunflower	Mallow, common	(creeping charlie)	Bedstraw, pineywoods
Thistle, russian	Pokeweed	Mallow, bristly	Çreeping Jenny	Bedstraw, Northern
Thistle, plumeless	Poison oak	Mallow, alkali	Creeping Beggarweed	Bedstraw, smooth
Thistle, platte	yvi nozio¶	геzəpədsə	Corn Spurry	Bedstraw, Catchweed
Thistle, musk	Plantain, woolly	Lambsquarters	Common mullein	szerian fieldcress
Thistle, flodman (prairie)	Plantain, slender	Kochia	Cocklebur, spiny	Artichoke
Thistle, canada	Plantain, narrowleaf	Knotweed, prostrate	Cocklebur, oriental	Aster, white prairie
Thistle, bull	Plantain, hoary	Knawel	Cocklebur	Aster, purple
Thistle, blessed	Plantain, buckhorn	təloiv quqmul-ynndol	Cockle	Aster, white heath
Tanweed	Plantain, broadleaf	beewnosmil	Clover, white	Aster, heath
Tansy ragwort	Plantain, bracted	Horseweed	Clover, white sweet	Amaranth, slender
Strawberry, India mock	Plantain, blackseed	Horsenettle	Clover, yellow sweet	Amaranth, purple

WEEDS CONTROLLED spreader before each turn at edge of border strips.

DROP SPREADERS: Apply 2 strips along each end. Then apply along length of lawn, overlapping slightly and shutting off the BROADCAST SPREADERS: For best results, apply in a circular path starting on the outside and working toward the center Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended rate under actual application conditions. CALIBRATION INSTRUCTIONS AND SPREADER SETTINGS

lirections or when stopping. steady pace to ensure even coverage. Close the spreader opening when turning to change For best results, apply this product uniformly and avoid streaks, skips or overlaps. Walk at a For drop spreaders: Overlap the wheel tracks to ensure complete coverage. For rotary spreaders: Operate at intervals corresponding to the swath width. wher's Manual. Mever apply by hand.

Differences may occur due to condition of spreader, speed at which applied and roughness of as overapplication may cause injury to your lawn. Spreader settings are approximate. IMPORTANT: It is important to follow directions as indicated on this quality lawn care product etc.). One bag covers up to 5,000 square feet. its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, COVERAGE: To determine how much you need to apply, multiply the length of your lawn by Always apply with a rotary or drop spreader to distribute the granules evenly. Never apply

HOW TO APPLY - SPREADERS & CALIBRATION 5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

4. Avoid applying when temperatures are consistently above 90°F. time of application, sprinkle lightly with water before applying product. 2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).

3. Apply when Broadleaf weeds are young and actively growing, preferably in the morning when dew is on the grass. At the time of application, moisture on the weed leaves from dew, rainfall or irrigation/watering aids in control. If grass is not moist at

1. Mow lawn to normal height 1 to 2 days before application. **SNOITOURTSNI NOITADIJ99A**

treatment site is 2 per year.

Scotts® Accugreen®

DROP SPREADER

Scotts® EdgeGuard® Mini 4.5

Scotts® EdgeGuard® Deluxe 4.5

BROADCAST SPREADER

SPREADER SETTINGS

Scotts® SpeedyGreen®

Augustinegrass, Centipedegrass, hybrid Bermudagrass, and Colonial Bentgrass. Apply 2.6 pounds of product per 1,000 square feet of turfgrass. A second application may be made in approximately 30 days. The maximum number of broadcast applications per For use on Kentucky Bluegrass, Fescue spp., Perennial Ryegrass, common Bermudagrass, Bahiagrass, Zoysiagrass, St. Possible retreatment may be necessary if lawn is heavily overrun by weeds, if adverse low moisture conditions prevail or if weeds are

Apply when Broadleaf weeds are actively growing in Spring or early Fall. Apply when the air is calm to ensure uniform coverage and avoid spreading granules onto flowers, vegetables and ornamental

product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. fishponds, lakes, streams, reservoirs, marshes, estuaries, bays and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent Do not apply directly to or near water, storm drains, gutters, sewers or drainage ditches. Do not apply within 25 feet of rivers, Do not use this product on or near desirable plants and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of the roots of desirable trees and shrubs, since injury may result. Reseeding or overseeding areas may occur three (3) weeks after the application of this product.

Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees.

Do not exceed specified dosage for any area and do not apply this product within the drip line of, or over surface roots of trees, shrubs, and other ornamentals. Do not apply this product to newly seeded grasses until well established or after the second or third mowing.

Do not use on Dichondra, Carpetgrass, Floratam's St. Augustinegrass, Creeping Bentgrass or Turfgrass where desirable clovers **USE PRECAUTIONS**

of lawn weeds, including Dandelion, Chickweed, Knotweed, Plantain, Henbit and Spurge. In addition, a 28-0-3 fertilizer provides a This product is a combination of three proven weed killers—2,4-D, mecoprop-p, and dicamba. This herbicide controls a wide range

• Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production or for out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets

PRECAUTIONS AND RESTRICTIONS It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR USE the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treated area of the lawn or garden will help to This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm

nformation. FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300. information on this product, contact the National Pesticide Information Center, 1-800-858-7378, Monday - Friday, 8:00 am to 12:00 pm PST. You may also contact the National Poison Control Center, 1-800-222-1222, day or night, for emergency medical treatment Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For Call a poison control center or doctor for treatment advice. Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the foilet. Hazards to Humans and Domestic Animals PRECAUTIONARY STATEMENTS

STOP READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

	Derived from: Urea, Polymer-Coated Urea and Muriate of Potash. *5.6% Slow-Release Urea Mitrogen From Polymer-Coated Urea
%0'E	Soluble Potash (K ₂ O)
	negoriiV hea Witrogen
%0.82	
700 00	
	GUARANTEED ANALYSIS
EZ	



CUTOFF 12.875"