

STORAGE AND DISPOSAL
PESTICIDE STORAGE: Store product in original container in a cool, dry, locked storage area out of reach of children and pets.
PESTICIDE DISPOSAL AND CONTAINER HANDLING:
 Non-refillable containers. Do not reuse or refill this container.
If Empty: Place in trash or offer for recycling, if available.
If Partly Filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: May be harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum using tobacco, or using the toilet.

FIRST AID
If on Skin or Clothing:
 • Take off contaminated clothing.
 • Rinse skin immediately with plenty of water for 15-20 minutes.
 • Call a poison control center or doctor for treatment advice.
If in Eyes:
 • Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 • Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER:
 Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-225-2883 for general information and/or medical information.

ENVIRONMENTAL HAZARDS: This product is extremely toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.
NOTICE: To the extent consistent with applicable law, buyer assumes all risks of use, storage or handling of this product not in accordance with directions.
The Ortho® Guarantee: If for any reason you, the consumer, are not satisfied with this product, mail us original proof of purchase to obtain a full refund of your purchase price.
QUESTIONS OR COMMENTS? Call 1-800-225-2883



INSECT KILLER FLOWER & VEGETABLE GARDEN DUST

**KILLS INSECTS
 KILLS ON CONTACT**

**KILLS INSECTS
 FOR UP TO
 8
 MONTHS**

**INSECTS KILLED: APHIDS, WHITEFLIES,
 CABBAGE LOOPER & OTHER LISTED PESTS**



**KEEP OUT OF REACH OF CHILDREN
 CAUTION** See back panel for additional precautionary statements
**MANTENER FUERA DEL ALCANCE DE LOS NIÑOS
 PRECAUCIÓN** Consulte el panel posterior para conocer los avisos de precaución adicionales

Active Ingredients:
 Deltamethrin (S)-alpha-cyano-3-phenoxybenzyl (1R,3R)-3-(2,2-dibromovinyl)-2,2-dimethyl-
 cyclopropanecarboxylate 0.05%
Other Ingredients 99.95%
Total 100.00%

NET CONTENTS/ PESO NETO 1.75 LB (793 g)



**FOR USE ON HOME
 VEGETABLE GARDENS
 AND ORNAMENTALS**

**ORTHO® INSECT KILLER
 FLOWER & VEGETABLE GARDEN DUST**
 Controls a variety of insect pests in vegetable gardens and on flowers.

DIRECTIONS FOR USE:
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use in commercial food/feed handling establishments, restaurants or other establishments where food/feed is commercially prepared, processed or served.

IMPORTANT: Read the entire Directions for Use, Precautions and Disclaimer of Warranties before using this product.

Ortho® Insect Killer Flower & Vegetable Garden Dust is a ready-to-use insecticide dust which exhibits effective knockdown and long-lasting control against the pests listed when used according to label directions.


This product can be used for long-lasting pest control in and around homes and their immediate surroundings.

**WHERE TO USE:
 SHRUBS AND ORNAMENTALS:** Apply when insects or their damage first appear and repeat at weekly intervals if necessary. Apply 5 ounces per 1,000 square feet; or thoroughly apply to place a thin layer of dust on both upper and lower sides of leaves, stems and flowers.

ROSES, PETUNIAS, CHRYSANTHEMUMS, GERANIUMS, CARNATIONS AND OTHER RELATED FLOWERS: To control many leaf and flower chewing insects, apply as soon as the insect appears and repeat at 7 day intervals or as long as insects continue to appear. Apply a thin layer of dust on both upper and lower sides of leaves, stems and flowers.

HOME VEGETABLE GARDENS: Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering.

Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering.



Sold By:
 The Ortho Group
 P.O. Box 190 Mansville, OH 43040
 EPA Reg. No.: 73327-14-239
 EPA Est. No. 084175-TX-001

VEGETABLES	TARGET PESTS	REMARKS	Application Rates: Ounces of Dust /100 sq ft	Max Applications /Growing Season
Bulb Vegetables – garlic, leeks, onions, shallots	Cutworms, stink bugs, thrips	Do not apply more than recommended amount 0.112 lb of active ingredient per acre in one growing season. Allow 5 days between applications. Do not apply within one day of harvest.	2.13	4
Cucurbit Vegetables – cantaloupes, citrom melons, cucumbers, gourd, pumpkin, squash, watermelon	Armyworm, cabbage looper, Colorado potato beetle, corn earworm, cucumber beetles, cutworms, diamondback moth (larvae), flea beetles, grasshoppers, green peach aphid, leafhoppers, leafminers, melonworm, pickleworm, stink bugs, tarnished plant bug, tobacco budworm, tomato pinworm, vegetable weevil (adult), western plant bug, whitefly (adult)	Do not apply more than recommended amount 0.168 lb of active ingredient per acre in one growing season. Allow 3 days between applications. Do not apply within 3 days of harvest.	2.13	6
Fruiting Vegetables – eggplant, peppers, tomatillo, tomato	Aphid, armyworm, cabbage looper, Colorado potato beetle, cucumber beetle, cutworms, diamondback moth, European corn bore, flea beetles, imported cabbageworm, leafminers, pepper weevil, stink bug, tarnished plant bug, thrips, tomato fruitworm (corn ear worm), tomato pinworm, variegated cutworm, whitefly	Do not apply more than recommended amount 0.168 lb of active ingredient per acre in one growing season. Allow 5 days between applications. Do not apply within 1 day of harvest.	2.13	6
Root Vegetables – carrot, celery, chervil (rump-rooted)	Armyworm, cabbage looper, Colorado potato beetle, cutworms, European corn bore, flea beetles, green peach aphid, leafhoppers, lygus bug (plant bug), pea aphid, tarnished plant bug	Do not apply more than recommended amount 0.14 lb of active ingredient per acre in one growing season. Allow 3 days between applications. Do not apply within 3 days of harvest. Other than aphids, applications may not be made to root vegetables whose tops or leaves are to be harvested for human consumption.	2.13	5
Tuberous & Corn Vegetables – potato, sweet potato, singes, yam bean, true yam	Armyworm, cabbage looper, Colorado potato beetle, cutworms, European corn bore, flea beetles, green peach aphid, leafhoppers, lygus bug (plant bug), pea aphid, tarnished plant bug	Do not apply more than recommended amount 0.14 lb of active ingredient per acre in one growing season. Allow 3 days between applications. Do not apply within 3 days of harvest. Applications may not be made to tuberous or corn vegetables whose tops or leaves are to be harvested for human consumption.	2.13	5

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store product in original container in a cool, dry, locked storage area out of reach of children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING:

Non-refillable container. Do not reuse or refill this container.

If Empty: Place in trash or offer for recycling, if available.

If Partly Filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum using tobacco, or using the toilet.

FIRST AID

If on Skin or Clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

If in Eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER:

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-225-2883 for general information and/or medical information.

ENVIRONMENTAL HAZARDS: This product is extremely toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

NOTICE: To the extent consistent with applicable law, buyer assumes all risks of use, storage or handling of this product not used in accordance with directions.

The Ortho® Guarantee: If for any reason you, the consumer, are not satisfied with this product, mail us original proof of purchase to obtain a full refund of your purchase price.

QUESTIONS OR COMMENTS? Call 1-800-225-2883



INSECT KILLER FLOWER & VEGETABLE GARDEN DUST

KILLS INSECTS

KILLS ON CONTACT

INSECTS KILLED: APHIDS, WHITEFLIES,
CABBAGE LOOPER & OTHER LISTED PESTS

KILLS INSECTS
FOR UP TO

8

MONTHS



KEEP OUT OF REACH OF CHILDREN
CAUTION

See back panel for additional precautionary statements

MANTENER FUERA DEL ALCANCE DE LOS NIÑOS
PRECAUCIÓN

Consulte el panel posterior para conocer los avisos de precaución adicionales

Active Ingredient:

Deltamethrin (S)-alpha-cyano-3-phenoxybenzyl (1R,3R)-3-(2,2-dibromovinyl)-2,2-dimethyl-
cyclopropanecarboxylate

Other Ingredients

Total

0.05%

99.95%

100.00%

NET CONTENTS/ PESO NETO 1.75 LB (793 g)

FPO

SH1461

FOR USE ON HOME
VEGETABLE GARDENS
AND ORNAMENTALS

**ORTHO® INSECT KILLER
FLOWER & VEGETABLE GARDEN DUST**

Controls a variety of insect pests in vegetable gardens and on flowers.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use in commercial food/feed handling establishments, restaurants or other establishments where food/feed is commercially prepared, processed or served.

IMPORTANT: Read the entire Directions for Use, Precautions and Disclaimer of Warranties before using this product.

Ortho® Insect Killer Flower & Vegetable Garden Dust is a ready-to-use insecticide dust which exhibits effective knockdown and long-lasting control against the pests listed when used according to label directions.


This product can be used for long-lasting pest control in and around homes and their immediate surroundings.

WHERE TO USE:

SHRUBS AND ORNAMENTALS: Apply when insects or their damage first appear and repeat at weekly intervals if necessary. Apply 8 ounces per 1,000 square feet; or thoroughly apply to place a thin layer of dust on both upper and lower sides of leaves, stems and flowers.

ROSES, PETUNIAS, CHRYSANTHEMUMS, GERANIUMS, CARNATIONS AND OTHER RELATED FLOWERS: To control many leaf and flower chewing insects, apply as soon as the insect appears and repeat at 7 day intervals or as long as insects continue to appear. Apply a thin layer of dust on both upper and lower sides of leaves, stems and flowers.

HOME VEGETABLE GARDENS: Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering.



Sold By:
The Ortho Group
P.O. Box 190 Marysville, OH 43040
EPA Reg. No.: 73327-14-239
EPA Est. No. 084175-TX-001

VEGETABLES	TARGET PESTS	REMARKS	Application Rates: Ounces of Dust /100 sq ft	Max Applications /Growing Season
Bulb Vegetables – garlic, leeks, onions, shallots	Cutworms, stink bugs, thrips	Do not apply more than recommended amount 0.112 lb of active ingredient per acre in one growing season. Allow 5 days between applications. Do not apply within one day of harvest.	2.13	4
Cucurbit Vegetables – cantaloupes, citron melon, cucumber, gourd, pumpkin, squash, watermelon	Armyworm, cabbage looper, Colorado potato beetle, corn earworm, cucumber beetles, cutworms, diamondback moth (larvae), flea beetles, grasshoppers, green peach aphid, leafhoppers, leafminers, melonworm, pickleworm, stink bugs, tarnished plant bug, tobacco budworm, tomato pinworm, vegetable weevil (adult), western plant bug, whitefly (adult)	Do not apply more than recommended amount 0.168 lb of active ingredient per acre in one growing season. Allow 3 days between applications. Do not apply within 3 days of harvest.	2.13	6
Fruiting Vegetables – eggplant, peppers, tomatillo, tomato	Aphids, armyworm, cabbage looper, Colorado potato beetle, cucumber beetle, cutworms, diamondback moth, European corn borer, flea beetles, imported cabbageworm, leafminers, pepper weevil, stink bugs, tarnished plant bug, thrips, tomato fruitworm (corn earworm), tomato pinworm, variegated cutworm, whitefly	Do not apply more than recommended amount 0.168 lb of active ingredient per acre in one growing season. Allow 5 days between applications. Do not apply within 1 day of harvest.	2.13	6
Root Vegetables – carrot, celeriac, chervil (turnip-rooted)	Armyworm, cabbage looper, Colorado potato beetle, cutworms, European corn borer, flea beetles, green peach aphid, leafhoppers, lygus bug (plant bug), pea aphid, Tarnished plant bug	Do not apply more than recommended amount 0.14 lb of active ingredient per acre in one growing season. Allow 3 days between applications. Do not apply within 3 days of harvest. Other than radishes, applications may not be made to root vegetables whose tops or leaves are to be harvested for human consumption.	2.13	5
Tuberous & Corm Vegetables – potato, sweet potato, ginger, yam bean, true yam	Armyworm, cabbage looper, Colorado potato beetle, cutworms, European corn borer, flea beetles, green peach aphid, leafhoppers, lygus bug (plant bug), pea aphid, Tarnished plant bug	Do not apply more than recommended amount 0.14 lb of active ingredient per acre in one growing season. Allow 3 days between applications. Do not apply within 3 days of harvest. Applications may not be made to tuberous or corm vegetables whose tops or leaves are to be harvested for human consumption.	2.13	5