


# NATRIA®

# Lawn Weed & Disease Control

Concentrate

## Kills Weeds, Down to the Root

## Kills Most Common Lawn Diseases* Fast




*Not for use in California

| Active Ingredient: | |
|---|---|
| Iron HEDTA (FeHEDTA) | 26.52% |
| Other Ingredients | 73.48% |
| Total | 100.00% |

See Results In Hours!

NET CONTENTS
24 FL OZ (709ml)

KEEP OUT OF REACH OF CHILDREN

CAUTION See back panel for additional precautionary statements

50141503b 22010711

# NATRIA®

## Lawn Weed & Disease Control

Concentrate

## PRECAUTIONARY STATEMENTS

**CAUTION Hazards to Humans and Domestic Animals**

Causes slight eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

## FIRST AID ✚

**IF IN EYES**

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **For medical emergency information, you may call toll free 1-877-229-3763.** During other times, call the poison control center 1-800-222-1222.

**NOTICE TO BUYER:** Seller warrants that this product conforms to the chemical description on this label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions. To the extent consistent with applicable law, this warranty does not extend to use of this product contrary to label directions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller. To the extent consistent with applicable law, the Seller shall not be liable for any such use. To the extent consistent with applicable law, the Seller makes no other warranties, either expressed or implied.

**Money Back Guarantee:** If you are not satisfied with this product, we will gladly refund your original purchase price.

**Garantía de devolución de dinero:** Si usted no está satisfecho con este producto, gustosamente le reembolsaremos el monto original del precio de compra.

**EPA Reg. No. 67702-26-92564**

EPA Est. No indicated by 3rd and 4th digits of the batch number on this package.
(47) = 52251-OR-5 (65) = 92564-TX-1 (57) = 71106-GA-3

NEUDORFF Sold under License of W. Neudorff GmbH KG

Natria® is a registered trademark of SBM Life Science Corp.

Distributed by:
SBM Life Science Corp.
1001 Winstead Drive, Suite 500
Cary, NC 27513

PRESS TO RESEAL

50142385Tb 220107V1

7/15/22

## ALMACENAMIENTO Y ELIMINACIÓN

### ALMACENAMIENTO DEL PESTICIDA

Almacene este producto en su recipiente original y manténgalo en un área de almacenamiento seguro fuera del alcance de niños y animales domésticos. Almacene material sin usar, bien cerrado, en su recipiente original solamente, lejos del fuego.

### ELIMINACIÓN DEL PESTICIDA Y MANIPULACIÓN DEL CONTENEDOR:

Contenedor no rellenable. No rellenar o volver a usar este contenedor. **Si está vacío:** Colocar en la basura para su reciclado si fuese posible. **Si está parcialmente lleno:** Llame a su agencia local de desperdicios sólidos o llame a 1 877-229-3724 para recibir indicaciones de eliminación. Nunca colocar el producto sin usar en un drenaje de salida o de entrada.

**MANTENER FUERA DEL ALCANCE DE LOS NIÑOS**

## DECLARACIONES DE PRECAUCIÓN

**PRECAUCIÓN** **Riesgos para Humanos o Animales Domésticos**

Causa irritación leve en los ojos. Evitar contacto con los ojos o ropa. Lavar cuidadosamente con agua y jabón luego de manipularlo y antes de comer, beber, masticar goma de mascar, usar tabaco o ir al baño. El contacto prolongado o repetido con la piel puede causar reacciones alérgicas en algunas personas.

## PRIMEROS AUXILIOS +

### EN CASO DE CONTACTO CON LOS OJOS

• Mantener los ojos abiertos y enjuagar lentamente y cuidadosamente con agua durante 15 o 20 minutos. • Quitar los lentes de contacto, si los tuviera, luego de los primeros 5 minutos, continuar lavando. • Llamar al centro de control de envenenamiento o al médico para asesoramiento sobre el tratamiento.

Tenga el contenedor o etiqueta del producto con usted cuando llame al centro de control de envenenamiento o al médico, o cuando vaya para recibir tratamiento. **Para recibir información médica en caso de emergencia, puede llamar de manera gratuita al 1-877-229-3763.** En otros horarios llamar al centro de control de envenenamiento al 1-800-222-1222

## RIESGOS AMBIENTALES

Para proteger el medio ambiente, no permita que el pesticida ingrese o corra por los drenajes de tormenta, drenaje de aguas pluviales, canaletas o aguas superficiales. Aplique este producto en clima calmo cuando no haya pronóstico de lluvia en las próximas 24 horas para asegurar que el viento o la lluvia no soplen o laven el pesticida fuera del área de tratamiento. Enjuagar el equipo usado para la aplicación en el área tratada ayudará a evitar que el producto ingrese en cuerpos de agua o sistemas de drenaje.

**AVISO AL COMPRADOR:** El vendedor garantiza que este producto corresponde con la descripción química en esta etiqueta y que se adapta a los fines establecidos en las Indicaciones de uso, sujeto a los riesgos inherentes descritos arriba cuando se usa de acuerdo a las indicaciones de uso bajo condiciones de uso normal. Hasta donde corresponda de acuerdo a la ley, esta garantía no aplica si este producto no se usa de acuerdo a las indicaciones o bajo condiciones de uso normal, o bajo condiciones no razonablemente predecibles para el vendedor. Hasta donde lo permita la ley, el vendedor no será responsable de tal uso. Hasta donde lo permita la ley, el vendedor no garantiza otras cosas, ya sean expresas o implícitas.

# NATRIA®

## Lawn Weed & Disease Control Concentrate

*OPEN*
*Resealable Label for Directions & Precautions*

**READ THE LABEL FIRST!**

Indicaciones en español después de las de inglés

## QUICK FACTS

- **5 in 1 Action!!** • **Kills Moss and Weeds, Not Lawns**
- **Derived from Iron** • **Even Works in Cool Weather**
- **Can be Used on New Lawns 1 Week After Grass Emergence**
- **Treats 1,875 sq. ft.**
- **Controls Disease, Weeds, Moss, Algae, Lichens in Turf**

| | |
|---|---|
| **CONTROLS** | Dandelion, Common Chickweed§‡, Clover and other Listed Broadleaf Weeds<br>Moss, Algae*, Lichens* and Liverworts*<br>Rust*, Dollar Spot* and Snow Mold* |
| **WHERE TO USE** | On lawns. (Do not apply to bentgrass) |
| ☎ | For questions or comments, call toll-free 1-877-229-3724. |

*Not for use in California § only approved for suppression in California
‡Minimum application rate in California is 11.5 fl.oz/yd² or 9.8 ga/1000 sq.ft.

**ACTIVE INGREDIENT:**
Iron HEDTA (FeHEDTA) ..... 26.52%
**OTHER INGREDIENTS** ..... 73.48%
**TOTAL** 100.00%

**EPA Reg. No. 67702-26-92564**

EPA Est. No indicated by 3rd and 4th digits of the batch number on this package.
(47) = 52251-OR-5 (65) = 92564-TX-1
(57) = 71106-GA-3

501423851b 220107V1

6 87073 07310 2

**7/15/22**

## DIRECTIONS FOR USE

**It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.**

### USE SITES

Use to selectively control/suppress broadleaf weeds, moss, algae*, lichens*, liverworts* and diseases* growing in lawns.

### PESTS

**CONTROLS OR SUPPRESSES WEEDS SUCH AS, BUT NOT LIMITED TO**:
Black medic (*Medicago lupulina*) §, Broadleaf plantain (*Plantago major*)*, Bull thistle (*Cirsium vulgare*) , Canada thistle (*Cirsium arvense*)*, Common chickweed (*Stellaria media*)§ ‡ ,Creeping buttercup (*Ranunculus repens*), Creeping Charlie/Ground ivy (*Glechoma hederacea*)* , Dandelion (*Taraxicum officinale*), Dovefoot geranium (*Geranium molle*)*, English daisy (*Bellis perennis*) § , False dandelion (*Hypochoeris radicata*)*, Healall (*Prunella vulgaris*)*, Lawn burweed (*Soliva ptero-sperma*)*, Narrow-leaved plantain (*Plantago lanceolata*)*, Oxalis (*Oxalis spp.*)*, Prostrate knotweed (*Polygonum aviculare*)*, Shepherd's-purse (*Capsella bursa-pastoris*)*, Silverweed cinquefoil (*Potentilla anserina*)*, Speedwell spp.(including Persian Speedwell (*Veronica persica*) and Slender speedwell (*Veronica filiformis*)*, White clover (*Trifolium repens*), Wild chamomile (*Matricaria chamomilla*)*, Wild geranium (*Geranium robertianum*)*, moss, algae*, lichens*, liverworts*
**CONTROLS OR SUPPRESSES DISEASES SUCH AS, BUT NOT LIMITED TO**:
Dollar Spot*, Rust*, Snow Mold*

### APPLICATION DIRECTIONS

**FOR BROADCAST TREATMENT**

| Rate of Spray Solution | Rate of Concentrate | Weeds |
|---|---|---|
| 2.8 fl. oz./$yd^2$ or 2.5 ga/1000 $ft^{2\dagger}$ | 12.6 fl. oz/ 1000 $ft^2$ | ***Rate for moss* and easy-to-control weeds:*** Black medic§, Persian speedwell, Slender speedwell*, Wild geranium*, and Moss. ***This rate can also be used for weed seedlings of:*** False dandelion*, English daisy*, Dandelion*, or White clover*. |
| 5.7 fl. oz./$yd^2$ or 4.9 ga/1000 $ft^2$ | 25.2 fl. oz/ 1000 $ft^2$ | ***Standard rate for control of most weed species***: Black medic, Dandelion, English daisy, False dandelion*, Creeping Charlie/Ground ivy*, Prostrate knotweed*, White clover, Bull thistle, Canada thistle*, Narrow-leaved plantain*, Dove's foot geranium*, Oxalis*, Lawn burweed*, and Algae*, Lichens*, Liverworts* and for the SUPPRESSION of Broad-leaved plantain* |

*(contd)*

## DIRECTIONS FOR USE *(cont'd)*

### APPLICATION DIRECTIONS *(cont'd)*

**FOR BROADCAST TREATMENT** *(cont'd)*

| Rate of Spray Solution | Rate of Concentrate | Weeds |
|---|---|---|
| 11.5 fl oz./$yd^2$ or 9.8 ga/1000 $ft^2$ | 50 fl. oz/ 1000 $ft^2$ | ***Rate for tough perennial weeds such as:*** Creeping buttercup, common chickweed§ and broad-leaved plantain* |
| **Rate of Spray Solution** | **Rate of Concentrate** | **Diseases** |
| 2.8 fl. oz./$yd^2$ or 2.5 ga/1000 $ft^{2\dagger}$ | 12.6 fl. oz/ 1000 $ft^2$ | ***Rate for control or suppression of diseases such as*** Dollar spot* , Snow mold*, and Rust*. Use the higher rate when disease pressure is high. |
| 5.7 fl. oz./$yd^2$ or 4.9 ga/1000 $ft^2$ | 25.2 fl. oz/ 1000 $ft^2$ | |

Shake well before using. Mix one part of this product with 24 parts water (5 oz. in1 gallon of water). Apply the mixed solution at a rate of 2.5-10 gallons/1000 $ft^2$ (2.8-11.5 fluid oz/ $yard^2$), or 5-10 gallons/1000 $ft^2$ (5.7-11.5 fl.oz./$yd^2$) in California, using the chart to determine how much diluted spray to use per area. Any standard hand-held or backpack sprayer can be used. Thoroughly spray weeds. Use a coarse nozzle setting to reduce drift. Uniform coverage is important. Repeat treatment in 3 to 4 weeks for best results.

To get the most complete kill of moss, or when moss is very thick, rake out as much moss as you can using a thatching rake or lawn rake before using this product.

To improve the appearance of lawn or turf, rake out dead moss a few days after treatment.

**FOR SPOT TREATMENT**

Shake well before using. Mix one part of this product with 24 parts water (5 oz. in 1 gallon of water). Thoroughly spray weeds to point of runoff using any standard hand-held or backpack sprayer. Use a coarse nozzle setting to reduce drift. Repeat treatment in 3 to 4 weeks for best results.

**FOR PRE-EMERGENT CONTROL* OF BROADLEAF WEEDS AND CRABGRASS**

Begin application when soil temperatures reach 50 to 55 °F. Shake well before using. Mix one part of this product with 24 parts water (5 fl. oz. in 1 gallon of water). Apply 5 to 10 gallons of spray solution per 1,000 sq. ft. depending on weed pressure. Reapply 3 to 6 weeks later for best results.

### WHAT WILL I SEE?

Works quickly; dying weeds will be seen within hours of application. Treated weeds, moss, or algae* will turn brown or black. Occasionally a darkening of the grass leaf blades can

*(cont'd)*

* Not for use in California § only approved for suppression in California

† Lower application rate not for use in California. Minimum application rate in California is 5.7 fl./$yd^2$ or 4.9 ga/1000 sq.ft. ‡Minimum application rate in California is 11.5 fl.oz./$yd^2$ or 9.8 ga/1000 sq.ft.

7/15/22

## DIRECTIONS FOR USE *(cont'd)*

occur after treatment; however the grass will recover within a few days to weeks. May stain thatch (dead grass under lawn); this does not affect the health of the grass.

### USE PRECAUTIONS

Do not apply to drought stressed grass; ensure lawn is well watered prior to application. Do not apply when the daytime temperature will exceed 85ºF. Do not apply to bentgrass. If unsure of grass sensitivity, test spray a small area of your lawn. Do not apply to newly seeded areas prior to grass seed germination. Avoid spraying desirable plants. Any desirable plants or structures accidentally contacted should be rinsed with water. To avoid staining, wait until treated area dries before re-entry. Do not apply if rainfall is expected within 3 hours of application. Do not irrigate within 3 hours of application. Do not apply to the same area more than 4 times a year.

## STORAGE AND DISPOSAL

### PESTICIDE STORAGE

Store this product in its original container and keep in a secure storage area out of reach of children and domestic animals. Store unused material, tightly closed, in original container only, away from open flame.

### PESTICIDE DISPOSAL AND CONTAINER HANDLING

Non-refillable container. Do not reuse or refill this container.
**If Empty:**
Place in trash or offer for recycling if available.
**If Partly Filled:**
Call your local solid waste agency or call 1 877-229-3724 for disposal instructions. Never place unused product down any indoor or outdoor drain.

## PRECAUTIONARY STATEMENTS

### ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

*(cont'd)*

**¡LEER LA ETIQUETA ANTES DE UTILIZAR EL PRODUCTO!**

## DATOS RÁPIDOS

- **Acción 5 en 1**
- **Mata el moho y las malezas, no el césped**
- **Derivado del hierro**
- **Hasta funciona en el clima frío.**
- **Puede usarse en césped nuevo 1 semana después de que sale el césped**
- **Trata 1,875 pies cuadrados**
- **Controla enfermedades, malezas, moho, algas, liquen en el césped**

| | |
|---|---|
| **CONTROLA** | Diente de león, capiqui común§‡, trébol y otras malezas de hoja ancha de la lista<br>Moho, algas*, líquenes* y plantas hepáticas*.<br>Roya*, Mancha del dólar* y Moho de la nieve* |
| **DÓNDE USAR** | En céspedes (no aplicar en agrostis) |
| ☎ | Si tiene preguntas o comentarios, llame al número gratuito 1-877-229-3724. |

**INGREDIENTE ACTIVO:**
Hierro HEDTA (FeHEDTA) ..... 26.52%
**OTROS INGREDIENTES** ..... 73.48%
**TOTAL** 100.00%

*No puede usarse en California §aprobado solamente para eliminación en California ‡La tasa mínima de aplicación en California es 11.5 oz./yardas cuadradas o 9.8 ga/1000 pies cuadrados.

## INDICACIONES DE USO

**Es una violación de la Ley Federal utilizar este producto de manera contraria a la indicada en la etiqueta.**

### LUGARES DE USO

Usar para controlar/eliminar de manera selectiva malezas de hoja ancha, moho, algas*, líquenes*, hepáticas* y enfermedades* que crecen en los jardines.

### PLAGAS

**CONTROLA O EVITA MALEZAS TALES COMO PERO NO LIMITADAS A**:
Mielga negra (*Medicago lupulina*) §, plátano de hoja ancha (*Plantago major*)*, cardo negro

*(continuación)*

7/15/22

## INDICACIONES DE USO *(continuación)*

### PLAGAS *(continuación)*

(*Cirsium vulgare*) , cardo de Canadá (*Cirsium arvense*)*, oreja de ratón común (*Stellaria media*)§ ‡, botón de oro (*Ranunculus repens*), Creeping Charlie/hiedra rastrera (*Glechoma hederacea*)*, diente de león (*Taraxicum officinale*), geranio pie de paloma (*Geranium molle*)*, margarita inglesa (*Bellis perennis*) § , diente de león falso (*Hypochoeris radicata*)*, hierba del carpintero (*Prunella vulgaris*)*, burweed engomada del césped (*Soliva ptero-sperma*)*, plátano de hoja angosta (*Plantago lanceolata*)*, oxalis (*Oxalis spp.*)*, sanguinaria mayor (*Polygonum aviculare*)*, bolsa de pastor (*Capsella bursa-pastoris*)*, potentilla argentina (*Potentilla anserina*)*, especies de verónica (incluyendo verónica pérsica (*Veronica persica*) y Verónica pequeña (*Veronica liformis*)*), trébol blanco (*Trifolium repens*), manzanilla salvaje (*Matricaria chamomilla*)*, geranio salvaje (*Geranium robertianum*)*, moho, algas*, líquenes*, plantas hepáticas*

**CONTROLA O EVITA ENFERMEDADES TALES COMO PERO NO LIMITADAS A**:
Mancha del dólar* , Roya*, Moho de la nieve*

### INDICACIONES DE APLICACIÓN

| Flujo de solución de rocío | Grado de concentrado | Malezas |
|---|---|---|
| 2.8 onzas líqu./yardas$^2$ o 2.5 galones/1000 pies$^2$ | 12.6 onzas líqu./1000 pies$^2$ | ***Flujo para moho* y malezas de control fácil:*** Mielga negra, verónica pérsica, verónica pequeña*, geranio salvaje*, y moho. ***Este flujo puede usarse también para malezas que recién aparecen de:*** diente de león falso*, margarita inglesa*, diente de león* o trébol blanco*. |
| 5.7 onzas líqu./yardas$^2$ o 4.9 galones/1000 pies$^2$ | 25.2 onzas líqu./1000 pies$^2$ | ***Flujo estándar para control de la mayoría de las especies de malezas:*** Mielga negra, diente de león, margarita inglesa, diente de león falso*, Creeping Charlie/hiedra rastrera*, sanguinaria mayor*, trébol blanco, cardo negro, cardo de Canadá*, plátano de hoja angosta* , geranio pie de paloma*, oxalis*, burweed engomada del césped *, y algas*, líquenes*, plantas hepáticas* y para la *ELIMINACIÓN* del plátano de hoja ancha* |
| 11.5 onzas líqu./yardas$^2$ o 9.8 galones/1000 pies$^2$ | 50 onzas líqu./ 1000 pies$^2$ | ***Flujo para malezas perennes resistentes tal como:*** botón de oro, oreja de ratón común§ y plátano de hoja ancha* |
| 2.8 onzas líqu./yardas$^2$ o 2.5 galones/1000 pies$^2$ | 12.6 onzas líqu./1000 pies$^2$ | ***Flujo para control o eliminación de enfermedades tal como:*** la mancha del dólar*, moho de la nieve* y roya*. Usar un |

## INDICACIONES DE USO *(continuación)*

### INDICACIONES DE APLICACIÓN

**PARA TRATAMIENTO POR ESPARCIMIENTO *(continuación)***

| Flujo de solución de rocío | Grado de concentrado | Enfermedades |
|---|---|---|
| 5.7 onzas líqu./yardas$^2$ o 4.9 galones/1000 pies$^2$ | 25.2 onzas líqu./1000 pies$^2$ | flujo más alto cuando la presión de la enfermedad es alta. |

Agitar bien antes de usar. Mezcle una parte de este producto con 24 partes de agua (5 onzas en 1 galón de agua). Aplique la solución mezclada en un flujo de 2.5-10 galones/1000 pies$^2$ (2.8-11.5 onzas líqu./yardas$^2$), o 5-10 galones/1000 pies$^2$ (5.7-11.5 .oz./yardas$^2$) en California, usando la tabla para determinar cuánto rocío diluido usar por área. Puede usar cualquier rociador portátil o mochila estándar. Rocíe las malezas completamente. Use una boquilla de rocío grueso para evitar salpicarse. Es importante cubrir el follaje de manera uniforme. Repetir el tratamiento entre 3 a 4 semanas para lograr mejores resultados. Para eliminar completamente el moho o cuando el musgo es muy grueso, rastrille y quite tanto musgo como pueda con un rastrillo antes de usar este producto. Para mejorar el aspecto del césped o pasto, rastrille y quite el musgo muerto unos días después del tratamiento.

**PARA TRATAMIENTO LOCALIZADO**

Agitar bien antes de usar. Mezcle una parte de este producto con 24 partes de agua (5 onzas en 1 galón de agua). Rociar la maleza completamente hasta que el producto corra usando un rociador estándar portátil o tipo mochila. Use una boquilla de rocío grueso para evitar salpicarse. Repetir el tratamiento entre 3 a 4 semanas para lograr mejores resultados.

**PARA CONTROL PREEMERGENTE* DE MALEZAS DE HOJA ANCHA Y GARRANCHUELO**

Comience la aplicación cuando las temperaturas del suelo lleguen a 50 hasta 55 °F. Agitar bien antes de usar. Mezclar una parte de este producto con 24 partes de agua (5 onzas en 1 galón de agua). Aplicar 5 a 10 galones de la solución por cada 1,000 pies cuadrados dependiendo de la presión de la maleza. Volver a aplicad dentro de 3 a 6 semanas para lograr mejores resultados.

### ¿QUÉ VERÉ?

Trabaja rápidamente, verá cómo mueren las malezas a las horas de su aplicación. Las malezas, moho o algas* tratadas se pondrán de color marrón o negro. Puede ocurrir un oscurecimiento de las hojas del césped luego del tratamiento; sin embargo el césped se recuperará dentro de unos pocos días o semanas. Puede manchar la paja (pasto muerto bajo el césped); pero esto no afectará la salud del césped.

### PRECAUCIONES DE USO

No aplique al césped estresado seco; asegúrese de que el jardín esté bien regado antes de la aplicación. No aplique cuando la temperatura del día exceda los 85º F. No aplicar al agrostis. Si no está seguro sobre la sensibilidad del césped, pruebe el rocío en un área pequeña de su césped. No aplique en áreas recientemente sembradas antes de que germine la semilla del césped. Evitar rociar plantas preferidas. Si accidentalmente el producto entra en contacto con plantas o estructuras, enjuáguelas inmediatamente con agua. Para evitar manchar, espere a que el área tratada se seque antes de volver a ingresar. No aplicar si se espera lluvia dentro de las 3 horas de la aplicación. No regar hasta dentro de las 3 horas de su aplicación. No aplicar en la misma área más de 4 veces por año.