

**KILLS EVEN
TOUGH WEEDS
ROOTS AND ALL**

IMAGE®

Herbicide Consumer Concentrate

ALL-IN-ONE LAWN WEED KILLER*

CONCENTRATE



DANDELION



NUTSEGE



CLOVER



CRABGRASS

KILLS 100+ WEEDS

WON'T HARM LAWNS**

KEEP OUT OF REACH OF CHILDREN

CAUTION See Booklet for Additional Precautionary Information,
First Aid and Complete Directions For Use.

* Controls Listed Annual Grasses, Broadleaf Weeds and Sedges

** When used as directed

300523712 - 3/22 | 1M187 V3

COVERS **6,000** SQ. FT.

Active Ingredients:	By Wt.
Sulfentrazone	1.6%
Quinclorac	5.0%
Other Ingredients	93.4%
	<hr/> 100.0%

NET CONTENTS 24 FL OZ (710 mL)

IMAGE[®]

Herbicide Consumer Concentrate

ALL-IN-ONE LAWN WEED KILLER*

CONCENTRATE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wear long-sleeved shirt, long pants, shoes, socks and waterproof gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to marine/estuarine invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems. This product may impact surface water quality due to runoff of rain water or spray drift. Under certain conditions this product may have a high potential for runoff into surface water for several months after application. This is especially true for poorly draining soils and soils with shallow ground water. The active ingredients in this product are known to leach through soil into groundwater under certain conditions as a result of labeled use. Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

FIRST AID

If swallowed

- Call poison control center or doctor immediately for treatment advice.
- Have a person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

If on skin or clothing

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

If in eyes

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-265-0761 for emergency medical treatment information.



© 2015, 2017, 2020, 2022
Central Garden & Pet Company
300523713 - 3/22 | IM187 V3

Manufactured for:
Central Garden & Pet, Garden Division
1000 Parkwood Circle, Suite 700
Atlanta, GA 30339
1-800-265-0761
gardendivision@central.com

EPA Reg No. 279-3463-90098
EPA Est. No. 59144-GA-1_a, 67572-GA-1_b, 67572-GA-2_c,
74750-GA-1_b, 70815-GA-1_c, 5905-GA-1_i
Subscript used is first letter of Run Code on container.

**KILLS
WEEDS FAST**

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Observe all cautions and limitations on this label. The use of this product not consistent with this label may result in plant injury. Keep container closed to avoid spills and contamination.

APPLICATION INSTRUCTIONS AND PRECAUTIONS

WHERE TO USE	Lawns: Well established turfgrasses. Cool Season Grasses: Fine & Tall Fescue, Perennial Ryegrass, Kentucky & Rough Bluegrass. Warm Season Grasses: Common & Hybrid Bermudagrass, Buffalograss, Centipedegrass, Seashore Paspalum, Zoysiagrass. Do not use on St. Augustinegrass.
KILLS	Sedges, Grasses and Broadleaf Weeds: Yellow Nutsedge, Kyllinga, Crabgrass, Foxtail, Ground Ivy, Wild Violet, Plantain, Dandelion, Clover, Dollarweed, Spurge, Woodsorrel, Knotweed, Chickweed, Curly Dock, Wild Onion, Wild Garlic, and other listed weeds.
WHEN TO USE	When weeds are small and actively growing. For best performance, apply when temperatures are between 55°F - 85°F.
HOW MUCH TO USE	4 fl oz per 1 gallon of water per 1,000 ft ² . Spot spray weed directly until thoroughly covered. Some weeds may require repeat application. (See detailed instructions.)
COVERAGE	24 fl.oz. covers up to 6,000 sq.ft.

See inside booklet for additional Precautionary Information, First Aid and complete Directions For Use.

300523713 - 3/22 | IM187 V3



Manufactured for:
Central Garden & Pet, Garden Division
1000 Parkwood Circle, Suite 700
Atlanta, GA 30339

1-800-265-0761
gardendivision@central.com

EPA Reg No. 279-3463-90098
EPA Est. No. 59144-GA-1_a, 67572-GA-1_b, 67572-GA-2_c,
74750-GA-1_b, 70815-GA-1_e, 5905-GA-1_f
Subscript used is first letter of Run Code on container.

Image, Central Garden & Pet, and Central Garden & Pet with design are trademarks of Central Garden & Pet Company.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Observe all cautions and limitations on this label. The use of this product not consistent with this label may result in plant injury. Keep container closed to avoid spills and contamination.

PRODUCT INFORMATION

This product selectively controls tough weeds like crabgrass, chickweed, dandelion, clover, ground ivy, purple[†] and yellow nutsedge (nutgrass), Kyllinga, wild onion and garlic, broadleaf plantain, purslane, wild violet, spurge, and other listed annual, biennial, and broadleaf weeds in a variety of lawns, including:

Cool Season Grasses	Warm Season Grasses
Tall & Fine Fescue, Kentucky Bluegrass, Rough Bluegrass, Ryegrass (perennial)	Common & Hybrid Bermudagrass, Zoysiagrass, Centipedegrass, Buffalograss, Seashore Paspalum

[†]Not for use in California

This product has demonstrated tolerance on both cool and warm season turfgrasses. However, not all varieties have been evaluated. Some turf varieties are more susceptible to temporary turf discoloration than others. It is recommended to treat a small area prior to treatment of larger areas.

This unique formula causes visual symptoms to develop quickly within 3 to 7 days after application. This product involves uptake by both weed roots and shoots. Complete kill should occur in 2-3 weeks. This product also controls yellow nutsedge tubers (nutlets) in the soil over time.

A postemergence application of this product is improved when adequate soil moisture is present at application.

HOW TO APPLY:

- Measure the entire area to be sprayed. For rectangular areas, multiply the length by width to obtain total square feet.
- Shake well before use.
- Remove measuring cap and fill cap according to chart below.
- Pour into sprayer and add corresponding amount of water.

Amount of Product	Amount of Water	Coverage
1 fl oz	1 Quart (32 oz)	250 sq.ft.
2 fl oz	½ Gallon (64 oz)	500 sq.ft.
4 fl oz	1 Gallon (128 oz)	1000 sq.ft.
6 fl oz	1½ Gallons (192 oz)	1500 sq.ft.
8 fl oz	2 Gallons (256 oz)	2000 sq.ft.

PUMP-STYLE TANK SPRAYERS

- Close sprayer tightly and pressurize with hand pump.
- For spot applications, use a coarse spray nozzle. For larger applications, use a fan spray nozzle.
- Hold tip of wand 12 inches from target area while spraying.

DIAL-STYLE HOSE-END SPRAYERS

- Fill sprayer jar with enough product for the area to be treated. DO NOT ADD WATER.
- Close sprayer, set dial, hook to garden hose and apply.
- Pour any unused product back into the product's original container.

APPLICATION INSTRUCTIONS:

- Best weed control results will be obtained when no rainfall or irrigation occurs within 24 hours after application.
- Listed sedges and grassy weeds: Spray directly on emerged weeds. Image® Herbicide Concentrate All-in-One Lawn Weed Killer® Concentrate is effective against newly emerged and established sedges and grassy weeds.
- Listed broadleaf weeds: Spray directly on newly emerged weeds.
- For best results, apply to weeds when less than 3 inches high.
- Apply in calm weather.
- Begin spraying at the farthest point of the treatment area and work backward to avoid contact with wet treated surfaces.
- Good spray coverage is needed for optimum control.
- Avoid overlap spraying.
- Avoid side to side or swinging arm motion, as it can result in uneven coverage.
- If listed weeds are not controlled by the initial application, apply a second application 4 to 6 weeks after the initial treatment.
- Water lawn with at least 0.5 inches within 7 days (but not sooner than 1 day after application) if no rainfall or irrigation occurs within 7 days after application.

APPLICATION RESTRICTIONS

Keep children and pets out of the treated area until sprays have dried.

Do not spray over the top of flowers, ornamentals (including herbaceous) or garden vegetables.

Do not use on turfgrasses other than those listed on the label.

Do not use in greenhouses.

Do not graze or feed animals with treated grass clippings.

Do not apply to newly planted, plugged, or sodded turfgrass.

APPLICATION PRECAUTIONS

Use only on well-established non-stressed turfgrass. Applications to stressed plants (turfgrass or target weeds) may result in temporary discoloration or poor weed control. Allow turfgrass to develop good root system before applying. If discoloration occurs, treated turfgrass will recover with new growth and discolored leaf tissue will be removed with mowing. To reduce potential for discoloration, do not apply on turfgrass that is weakened by weather, mechanical, chemical, disease or other related stress. Maintain proper cultural practices such as adequate moisture and fertility levels to promote healthy turfgrass growth.

Avoid applying when windy to prevent product drifting to other plants.

Avoid reseeding, winter overseeding or plugging lawns for at least 4 weeks after treatment.

Grass and sedge weed species controlled or suppressed by this product:

Common Name	Scientific Name	Common Name	Scientific Name	Common Name	Scientific Name
Barnyardgrass	<i>(Echinochloa crus-galli)</i>	Foxtail, yellow	<i>(Setaria glauca)</i>	Junglerice	<i>(Echinochloa colona)</i>
Broadleaf Signalgrass	<i>(Brachiaria platyphylla)</i>	Kyllinga, cocks-comb	<i>(Kyllinga squamulata)</i>	Nutsedge, purple'	<i>(Cyperus rotundus)</i>
Crabgrass, Large	<i>(Digitaria sanguinalis)</i>	Kyllinga, green	<i>(Kyllinga brevifolia)</i>	Nutsedge, yellow	<i>(Cyperus esculentus)</i>
Crabgrass, Smooth	<i>(Digitaria ischaemum)</i>	Kyllinga, false green	<i>(Kyllinga gracillima)</i>	Sedge, annual	<i>(Cyperus compressus)</i>
Deergrass	<i>(Panicum cladesimum)</i>	Kyllinga, fragrant	<i>(Kyllinga sesquiflorus)</i>	Sedge, globe	<i>(Cyperus globulosus)</i>
Goosegrass	<i>(Eleusine indica)</i>	Kyllinga, tufted	<i>(Kyllinga pumila)</i>	Sedge, cylindrical	<i>(Cyperus retrorsus)</i>
Foxtail, giant	<i>(Setaria faberi)</i>	Japanese stiltgrass	<i>(Microstegium vimineum)</i>	Sedge, Surinam	<i>(Cyperus surinamensis)</i>
Foxtail, green	<i>(Setaria viridis)</i>			Sedge, Texas	<i>(Cyperus polystachyos)</i>

For optimal control of Goosegrass, Junglerice, or Purple Nutsedge¹, repeat treatment every 3-4 weeks.

¹Not for use in California

Broadleaf weeds controlled or suppressed by this product:

Common Name	Scientific Name	Common Name	Scientific Name	Common Name	Scientific Name
Bedstraw, catchweed	<i>(Galium aparine)</i>	Dollarweed	<i>(Hydrocotyle spp.)</i>	Morning glory, ivyleaf	<i>(Ipomoea hederacea)</i>
Beggarticks ¹	<i>(Bidens frondosa)</i>	(pennywort)		Morning glory,	<i>(Ipomoea wrightii)</i>
Braggweed, Florida	<i>(Desmodium tortuosum)</i>	Eclipta	<i>(Eclipta prostrata)</i>	palmleaf	
Bindweed, field	<i>(Convolvulus arvensis)</i>	Evening primrose	<i>(Oenothera biennis)</i>	Morning glory, pitted	<i>(Ipomoea lacunose)</i>
Bittercress	<i>(Cardamine spp.)</i>	Fiddleneck	<i>(Amsinckia spp.)</i>	Morning glory, tall	<i>(Ipomoea purpurea)</i>
Black medic	<i>(Medicago lupulina)</i>	Filaree	<i>(Erodium spp.)</i>	Parsley piert	<i>(Alchemilla arvensis)</i>
Buttercups	<i>(Ranunculus spp.)</i>	Filaree, redstem	<i>(Erodium cicutarium)</i>	Pigweed, Redroot	<i>(Amaranthus retroflexus)</i>
Buttonweed, common (poorjoe)	<i>(Diodia teres)</i>	Gallsingoa	<i>(Gallsingoa ciliate)</i>	Pigweed, Smooth	<i>(Amaranthus hybridus)</i>
Carolina geranium	<i>(Geranium carolinianum)</i>	Goldenrod	<i>(Solidago spp.)</i>	Pigweed, Tumble	<i>(Amaranthus albus)</i>
Carpetweed	<i>(Mollugo verticillata)</i>	Ground ivy	<i>(Glechmea hederacea)</i>	Pineapple weed	<i>(Matricaria matricariodes)</i>
Chickweed, common	<i>(Stellaria media)</i>	Groundsel, common		Plantain, Broadleaf	<i>(Plantago comanatala)</i>
Chickweed, mouseear	<i>(Cerastium vulgatum)</i>	Henbit	<i>(Lamium amplexicaule)</i>	Plantain, Buckhorn	<i>(Plantago lanceolata)</i>
Cinquefoil	<i>(Potentilla spp.)</i>	Knawel	<i>(Scleranthus annuus)</i>	Plantain, Narrowleaf	<i>(Plantago narrowiflora)</i>
Clover	<i>(Trifolium spp.)</i>	Knotweed, prostrate	<i>(Polygonum aviculare)</i>	Puncturevine	<i>(Tribulus terrestris)</i>
Clover, alyce	<i>(Alysicarpus ovalifolius)</i>	Kochia	<i>(Kochia scoparia)</i>	Purslane, common	<i>(Portulaca oleracea)</i>
Clover, hop	<i>(Trifolium campestre)</i>	Lambsquarters, common	<i>(Chenopodium album)</i>	Pusley, Florida	<i>(Richardia scabra)</i>
Clover, red	<i>(Trifolium pretense)</i>	Lawn burweed (Spurweed)	<i>(Soliva pterosperma)</i>	Redweed	<i>(Melochia corchorifolia)</i>
Cudweed	<i>(Gnaphalium spp.)</i>	Lespedeza, common	<i>(Lespedeza striata)</i>	Shepherd's-Purse	<i>(Capsella bursa-pastoris)</i>
Daisy, English	<i>(Bellis perennis)</i>	London Rocket	<i>(Sisymbrium irio)</i>		
Dandelion	<i>(Taraxacum officinale)</i>	Mallow, common	<i>(Malva neglecta)</i>		
Dandelion, false	<i>(Microseris cuspidate)</i>	Morning glory	<i>(Ipomea spp.)</i>		
Dayflower, common	<i>(Commelina diffusa)</i>	Morning glory, entireleaf	<i>(Ipomoea hederacea var. integruscula)</i>		
Dock, Curly	<i>(Rumex crispus)</i>				

Broadleaf weeds controlled or suppressed by this product (cont'd):

Common Name	Scientific Name	Common Name	Scientific Name	Common Name	Scientific Name
Smartweed, Pennsylvania	(<i>Polygonum pensylvanicum</i>)	Spurge, spotted	(<i>Euphorbia maculate</i>)	Wild onion	(<i>Allium canadense</i>)
Sorrel, Red	(<i>Rumex acetosella</i>)	Star-of-Bethlehem	(<i>Ornithogalum umbellatum</i>)	Woodsorrel, creeping	(<i>Oxalis corniculata</i>)
Speedwell	(<i>Veronica spp.</i>)	Velvetleaf	(<i>Abutilon theophrasti</i>)	Woodsorrel, yellow	(<i>Oxalis stricta</i>)
Speedwell, slender	(<i>Veronica filiformis</i>)	Violet, wild	(<i>Viola pratensis</i>)		
Speedwell, thymeleaf	(<i>Veronica serpyllifolia</i>)	Violet,	(<i>Viola rafeinesquii</i>)		
Spurge, (annuals)	(<i>Euphorbia spp.</i>)	Johnny-Jump-Up			
Spurge, prostrate	(<i>Euphorbia humistrata</i>)	Wild garlic	(<i>Allium vineale</i>)		

*Not for use in California

Certain listed weeds may require more than one application to achieve season-long control.

STORAGE AND DISPOSAL

Pesticide Storage

Keep from freezing. Do not store below 32°F. Keep out of reach of children and animals. Store in original container only. Store in a cool, dry place, preferably in a locked storage area and avoid excess heat.

In case of a spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call Central Garden & Pet, Garden Division: 1-800-265-0761.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Handling

Non-refillable container. Do not reuse or refill this container. **If empty:** Offer for recycling, if available, or dispose of in trash.

If partially filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: To the extent consistent with applicable law, buyer assumes all risks of use, storage or handling of this product not in accordance with label directions.

NOTES

NOTES
