279-3463-90098 Image Consumer Concentrate All-in-One Lawn Weed Killer Concentrate_20230828_16_90098.pdf

KILLS EVEN TOUGH WEEDS ROOTS AND ALL

IMAGE®

Herbicide Consumer Concentrate

ALL-IN-ONE LAWN WEED KILLER*

CONCENTRATE


DANDELION


NUTSEDGE


CLOVER


CRABGRASS

KILLS 100+ WEEDS

WON'T HARM LAWNS**

COVERS 6,000 SQ. FT.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Booklet for Additional Precautionary Information, First Aid and Complete Directions For Use.

Active Ingredients:	By Wt.
Sulfentrazone	1.6%
Quinclorac	5.0%
Other Ingredients	93.4%
	100.0%

*Controls Listed Annual Grasses, Broadleaf Weeds and Sedges

** When used as directed

300534284 - 1/23 | IM198 V3

NET CONTENTS 24 FL OZ (710 mL)

IMAGE®
Herbicide Consumer Concentrate

ALL-IN-ONE
LAWN WEED KILLER*

CONCENTRATE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wear long-sleeved shirt, long pants, shoes, socks and waterproof gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to marine/estuarine invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems. This product may impact surface water quality due to runoff of rain water or spray drift. Under certain conditions this product may have a high potential for runoff into surface water for several months after application. This is especially true for poorly draining soils and soils with shallow ground water.

The active ingredients in this product are known to leach through soil into groundwater under certain conditions as a result of labeled use. Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

FIRST AID

If swallowed	• Call poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-265-0761 for emergency medical treatment information.

STORAGE AND DISPOSAL

Pesticide Storage

Keep from freezing. Do not store below 32°F. Keep out of reach of children and animals. Store in original container only. Store in a cool, dry place, preferably in a locked storage area and avoid excess heat.

In case of a spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call Central Garden & Pet, Garden Division: 1-800-265-0761.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are acutely hazardous.Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Handling

Non-refillable container. Do not reuse or refill this container. If empty: Offer for recycling, if available, or dispose of in trash.

If partially filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

CENTRAL GARDEN & PET

Manufactured for:
Central Garden & Pet, Garden Division
1000 Parkwood Circle, Suite 700
Atlanta, GA 30339

1-800-265-0761
gardendivision@central.com

EPA Reg No. 279-3463-90098
EPA Est. No. 59144-GA-1$_b$, 67572-GA-1$_b$, 67572-GA-2$_c$, 74750-GA-1$_b$, 70815-GA-1$_c$, 5905-GA-1$_c$

Subscript used is first letter of Run Code on container.

© 2015-2023 Central Garden & Pet Company

300534285 – 1/23 | IMI98 V3

KILLS WEEDS FAST

IMAGE®

Herbicide Consumer Concentrate

ALL-IN-ONE LAWN WEED KILLER*

CONCENTRATE

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Observe all cautions and limitations on this label. The use of this product not consistent with this label may result in plant injury. Keep container closed to avoid spills and contamination.

APPLICATION INSTRUCTIONS AND PRECAUTIONS

WHERE TO USE	**Lawns:** **Well established turfgrasses.** **Cool Season Grasses:** Fine & Tall Fescue, Perennial Ryegrass, Kentucky & Rough Bluegrass. **Warm Season Grasses:** Common & Hybrid Bermudagrass, Buffalograss, Centipedegrass, Seashore Paspalum, Zoysiagrass. **Do not use on St. Augustinegrass.**
KILLS	Sedges, Grasses and Broadleaf Weeds: Yellow Nutsedge, Kyllinga, Crabgrass, Foxtail, Ground Ivy, Wild Violet, Plantain, Dandelion, Clover, Dollarweed, Spurge, Woodsorrel, Knotweed, Chickweed, Curly Dock, Wild Onion, Wild Garlic, and other listed weeds.
WHEN TO USE	When weeds are small and actively growing. For best performance, apply when temperatures are between 55°F - 85°F.
HOW MUCH TO USE	4 fl. oz. per 1 gallon of water per 1,000 ft². Spot spray weed directly until thoroughly covered. Some weeds may require repeat application. (See detailed instructions.)
COVERAGE	24 fl.oz. covers up to 6,000 sq.ft.

See inside booklet for additional Precautionary Information, First Aid and complete Directions For Use.

CENTRAL GARDEN & PET

Manufactured for:
Central Garden & Pet, Garden Division
1000 Parkwood Circle, Suite 700
Atlanta, GA 30339

1-800-265-0761
gardendivision@central.com

300534285 – 1/23 | IM198 V3

© 2015-2023 Central Garden & Pet Company

EPA Reg No. 279-3463-90098
EPA Est. No. 59144-GA-1, 67572-GA-1, 67572-GA-2, 74750-GA-1, 70815-GA-1, 5905-GA-1

Subscript used is first letter of Run Code on container.

Image, Central Garden & Pet, and Central Garden & Pet with design are trademarks of Central Garden & Pet Company.

8 13576 00691 4

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Observe all cautions and limitations on this label. The use of this product not consistent with this label may result in plant injury. Keep container closed to avoid spills and contamination.

PRODUCT INFORMATION

This product selectively controls tough weeds like crabgrass, chickweed, dandelion, clover, ground ivy, purple‡ and yellow nutsedge (nutgrass), Kyllinga, wild onion and garlic, broadleaf plantain, purslane, wild violet, spurge, and other listed annual, biennial, and broadleaf weeds in a variety of lawns, including:

Cool Season Grasses
Tall & Fine Fescue, Kentucky Bluegrass, Rough Bluegrass, Ryegrass (perennial)
Warm Season Grasses
Common & Hybrid Bermudagrass, Zoysiagrass, Centipedegrass, Buffalograss, Seashore Paspalum

‡Not for use in California

This product has demonstrated tolerance on both cool and warm season turfgrasses. However, not all varieties have been evaluated. Some turf varieties are more susceptible to temporary turf discoloration than others. It is recommended to treat a small area prior to treatment of larger areas.

This unique formula causes visual symptoms to develop quickly within 3 to 7 days after application. This product involves uptake by both weed roots and shoots. Complete kill should occur in 2-3 weeks. This product also controls yellow nutsedge tubers (nutlets) in the soil over time.

A postemergence application of this product is improved when adequate soil moisture is present at application.

HOW TO APPLY:

- Measure the entire area to be sprayed. For rectangular areas, multiply the length by width to obtain total square feet.
- Shake well before use.
- Remove measuring cap and fill cap according to chart below.
- Pour into sprayer and add corresponding amount of water.

Amount of Product	Amount of Water	Coverage
1 fl. oz.	1 Quart (32 oz)	250 sq.ft.
2 fl. oz.	½ Gallon (64 oz)	500 sq.ft.
4 fl. oz.	1 Gallon (128 oz)	1000 sq.ft.
6 fl. oz.	1½ Gallons (192 oz)	1500 sq.ft.
8 fl. oz.	2 Gallons (256 oz)	2000 sq.ft.

PUMP-STYLE TANK SPRAYERS

- Close sprayer tightly and pressurize with hand pump.
- For spot applications, use a coarse spray nozzle. For larger applications, use a fan spray nozzle.
- Hold tip of wand 12 inches from target area while spraying.

DIAL-STYLE HOSE-END SPRAYERS

- Fill sprayer jar with enough product for the area to be treated. DO NOT ADD WATER.
- Close sprayer, set dial, hook to garden hose and apply.
- Pour any unused product back into the product's original container.

APPLICATION INSTRUCTIONS:

- Best weed control results will be obtained when no rainfall or irrigation occurs within 24 hours after application.
- Listed sedges and grassy weeds: Spray directly on emerged weeds. Image® Herbicide Consumer Concentrate All-in-One Lawn Weed Killer* Concentrate is effective against newly emerged and established sedges and grassy weeds.
- Listed broadleaf weeds: Spray directly on newly emerged weeds.
- For best results, apply to weeds when less than 3 inches high.
- Apply in calm weather.
- Begin spraying at the farthest point of the treatment area and work backward to avoid contact with wet treated surfaces.
- Good spray coverage is needed for optimum control.
- Avoid overlap spraying.
- Avoid side to side or swinging arm motion, as it can result in uneven coverage.
- If listed weeds are not controlled by the initial application, apply a second application 4 to 6 weeks after the initial treatment.
- Water lawn with at least 0.5 inches within 7 days (but not sooner than 1 day after application) if no rainfall or irrigation occurs within 7 days after application.

APPLICATION RESTRICTIONS

Keep children and pets out of the treated area until sprays have dried.
Do not spray over the top of flowers, ornamentals (including herbaceous) or garden vegetables.
Do not use on turfgrasses other than those listed on the label.
Do not use in greenhouses.
Do not graze or feed animals with treated grass clippings.
Do not apply to newly planted, plugged, or sodded turfgrass.

APPLICATION PRECAUTIONS

Use only on well-established non-stressed turfgrass. Applications to stressed plants (turfgrass or target weeds) may result in temporary discoloration or poor weed control. Allow turfgrass to develop good root system before applying. If discoloration occurs, treated turfgrass will recover with new growth and discolored leaf tissue will be removed with mowing. To reduce potential for discoloration, do not apply on turfgrass that is weakened by weather, mechanical, chemical, disease or other related stress. Maintain proper cultural practices such as adequate moisture and fertility levels to promote healthy turfgrass growth.
Avoid applying when windy to prevent product drifting to other plants.
Avoid reseeding, winter overseeding or plugging lawns for at least 4 weeks after treatment.

Grass and sedge weed species controlled or suppressed by this product:

Common Name	Scientific Name	Common Name	Scientific Name
Barnyardgrass	*(Echinochloa crus-galli)*	Kyllinga, false green	*(Kyllinga gracillima)*
Broadleaf Signalgrass	*(Brachiaria platyphylla)*	Kyllinga, fragrant	*(Kyllinga sesquiflorus)*
Crabgrass, Large	*(Digitaria sanguinalis)*	Kyllinga, tufted	*(Kyllinga pumila)*
Crabgrass, Smooth	*(Digitaria ischaemum)*	Japanese stiltgrass	*(Microstegium vimineum)*
Deergrass	*(Panicum cladestinum)*	Junglerice	*(Echinochloa colona)*
Goosegrass	*(Eleusine indica)*	Nutsedge, purple‡	*(Cyperus rotundus)*
Foxtail, giant	*(Setaria faberi)*	Nutsedge, yellow	*(Cyperus esculentus)*
Foxtail, green	*(Setaria viridis)*	Sedge, annual	*(Cyperus compressus)*
Foxtail, yellow	*(Setaria glauca)*	Sedge, globe	*(Cyperus globulosus)*
Kyllinga, cocks-comb	*(Kyllinga squamulata)*	Sedge, cylindrical	*(Cyperus retrorsus)*
Kyllinga, green	*(Kyllinga brevifolia)*	Sedge, Surinam	*(Cyperus surinamensis)*
		Sedge, Texas	*(Cyperus polystachyos)*

For optimal control of Goosegrass, Junglerice, or Purple Nutsedge‡, repeat treatment every 3-4 weeks.
‡Not for use in California

Broadleaf weeds controlled or suppressed by this product:

Common Name	Scientific Name	Common Name	Scientific Name
Bedstraw, catchweed	*(Galium aparine)*	Morning glory	*(Ipomea spp.)*
Beggarticks‡	*(Bidens frondosa)*	Morning glory, entireleaf	*(Ipomoea hederacea var. integriuscula)*
Beggarweed, Florida	*(Desmodium tortuosum)*	Morning glory, ivyleaf	*(Ipomoea hederacea)*
Bindweed, field	*(Convolvulus arvensis)*	Morning glory, palmleaf	*(Ipomoea wrightii)*
Bittercress	*(Cardamine spp.)*	Morning glory, pitted	*(Ipomoea lacunose)*
Black medic	*(Medicago lupuline)*	Morning glory, tall	*(Ipomoea purpurea)*
Buttercups	*(Ranunculus spp.)*	Parsley piert	*(Alchemilla arvensis)*
Buttonweed, common (poorjoe)	*(Diodia teres)*	Pigweed, Redroot	*(Amaranthus retroflexus)*
Carolina geranium	*(Geranium carolinianum)*	Pigweed, Smooth	*(Amaranthus hybridus)*
Carpetweed	*(Mollugo verticillata)*	Pigweed, Tumble	*(Amaranthus albus)*
Chickweed, common	*(Stellaria media)*	Pineapple weed	*(Matricaria matricariodes)*
Chickweed, mouseear	*(Cerastium vulgatum)*	Plantain, Broadleaf	*(Plantago comanalata)*
Cinquefoil	*(Potentilla spp.)*	Plantain, Buckhorn	*(Plantago lanceolata)*
Clover	*(Trifolium spp.)*	Plantain, Narrowleaf	*(Plantago narowiforma)*
Clover, alyce	*(Alysicarpus ovalifolius)*	Puncturevine	*(Tribulus terrestris)*
Clover, hop	*(Trifolium campestre)*	Purslane, common	*(Portulaca oleracea)*
Clover, red	*(Trifolium pretense)*	Pusley, Florida	*(Richardia scabra)*
Clover, white	*(Trifolium repens)*	Redweed	*(Melochia corchorifolia)*
Cudweed	*(Gnaphalium spp.)*	Shepherd's-Purse	*(Capsella bursa-pastoris)*
Daisy, English	*(Bellis perennis)*	Smartweed, Pennsylvania	*(Polygonum pensylvanicum)*
Dandelion	*(Taraxacum officianale)*	Sorrel, Red	*(Rumex acetosella)*
Dandelion, false	*(Microseris cuspidate)*	Speedwell	*(Veronica spp.)*
Dayflower, common	*(Commelina diffusa)*	Speedwell, slender	*(Veronica filiformis)*
Dock, Curly	*(Rumex crispus)*	Speedwell, thymeleaf	*(Veronica serpyllifolia)*
Dollarweed (pennywort)	*(Hydrocotyle spp.)*	Spurge, (annuals)	*(Euphorbia spp.)*
Eclipta	*(Eclipta prostrata)*	Spurge, prostrate	*(Euphorbia humistrata)*
Evening primrose	*(Oenother biennis)*	Spurge, spotted	*(Euphorbia maculate)*
Fiddleneck	*(Amsinckia spp.)*	Star-of-Bethlehem	*(Ornithogalum umbellatum)*
Filaree	*(Erodium spp.)*	Velvetleaf	*(Abutilon theophrasti)*
Filaree, redstem	*(Erodium cicutarium)*	Violet, wild	*(Viola pratincola)*
Galinsoga	*(Galinsoga ciliate)*	Violet, Johnny-Jump-Up	*(Viola rafeinesquii)*
Goldenrod	*(Solidago spp.)*	Wild garlic	*(Allium vineale)*
Ground ivy	*(Glechmea hederacea)*	Wild onion	*(Allium canadense)*
Groundsel, common	*(Senecio vulgaris)*	Woodsorrel, creeping	*(Oxalis corniculata)*
Henbit	*(Lamium amplexicaule)*	Woodsorrel, yellow	*(Oxalis stricta)*
Knawel	*(Scleranthus annuus)*		
Knotweed, prostrate	*(Polygonum aviculare)*		
Kochia	*(Kockia scoparia)*		
Lambsquarters, common	*(Chenopodium album)*		
Lawn burweed (Spurweed)	*(Soliva pterosperma)*		
Lespedeza, common	*(Lespedeza striafa)*		
London Rocket	*(Sisymbrium irio)*		
Mallow, common	*(Malva neglecta)*		

‡Not for use in California
Certain listed weeds may require more than one application to achieve season-long control.

STORAGE AND DISPOSAL

Pesticide Storage

Keep from freezing. Do not store below 32°F. Keep out of reach of children and animals. Store in original container only. Store in a cool, dry place, preferably in a locked storage area and avoid excess heat.

In case of a spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call Central Garden & Pet, Garden Division: 1-800-265-0761.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are acutely hazardous.Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Handling

Non-refillable container. Do not reuse or refill this container. If empty: Offer for recycling, if available, or dispose of in trash.

If partially filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: To the extent consistent with applicable law, buyer assumes all risks of use, storage or handling of this product not in accordance with label directions.

NOTES