



080311

LESCO DIMENSION\* Crabgrass Pre-emergent PLUS FERTILIZER

COVERS 13,000 SQ FT

LESCO®

**DIMENSION\***  
**Crabgrass**  
**Pre-emergent**  
**PLUS FERTILIZER**

**19-0-7**


**N** NITROGEN - For grasses  
Made leaf and stem growth  
and a dark green color.

**P** PHOSPHATE - Contains  
no phosphorus

**K** POTASH - Promotes  
strong roots and  
improves turf hardiness.

Apply in late winter  
through early spring to  
prevent annual weeds.

PREVENTS CRABGRASS  
AND OVER 20 OTHER  
ANNUAL GRASS AND  
BROADLEAF WEEDS



Crabgrass Control  
MADE EASY!

ACTIVE INGREDIENT  
Diflufenican, S,S'-dimethyl 2-(difluoromethyl)-4-(3-methylpropyl)-6-(trifluoromethyl)-3,5-pyridinedicarboxylate 89.85%  
OTHER INGREDIENTS 10.15%  
TOTAL 100.00%

Product protected by U.S. Patent No. 4,682,184. Other patents pending.

KEEP OUT OF REACH OF CHILDREN  
**CAUTION**  
**PRECAUCIÓN**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail).

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.  
Read the entire label before using this product. Use only according to label instructions.

LESCO DIMENSION\* Crabgrass Pre-emergent PLUS FERTILIZER

080311



Net Wt. 50 lb (22.7 kg)

LESCO® **DIMENSION\*** Crabgrass  
Pre-emergent  
PLUS FERTILIZER

**LESCO® DIMENSION\* Crabgrass Pre-emergent PLUS FERTILIZER**

**GUARANTEED ANALYSIS**

Total Nitrogen (N)	19.00%
19.00% Urea Nitrogen (N*)	
Soluble Potash (K <sub>2</sub> O)	7.00%
Sulfur (S) Total	1.70%
1.70% Free Sulfur (S)	
Derived from: Polymer Coated Sulfur Coated Urea, Urea, Muriate of Potash, Chlorine (Cl), Malt	5.25%
*5.00% Slowly Available Urea Nitrogen from Polymer Coated Sulfur Coated Urea	F 1560

Read the entire label before using the product. Use only according to label instructions

**KEEP OUT OF REACH OF CHILDREN  
CAUTION  
PRECAUCIÓN**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail).

**FIRST AID**

- If swallowed**
  - Call a poison control center or doctor immediately for treatment advice.
  - Have person sip a glass of water if able to swallow.
  - Do not induce vomiting unless told to by the poison control center or doctor.
  - Do not give anything by mouth to an unconscious person.
- If in eyes**
  - Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
  - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
  - Call a poison control center or doctor for treatment advice.
- If on skin or clothing**
  - Take off contaminated clothing.
  - Rinse skin immediately with plenty of water for 15 to 20 minutes.
  - Call a poison control center or doctor for treatment advice.
- If inhaled**
  - Move person to fresh air.
  - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
  - Call a poison control center or doctor for further treatment advice.

**HOT LINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact CHEMTREC at 1-800-424-9300 for emergency medical treatment information.

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION:** Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact with handling materials may cause allergic reaction in some individuals.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment/washwaters.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner that is inconsistent with its labeling. Not for use on turf being grown for sale or other commercial use as soil, or for commercial seed production, or for research purposes.

**GENERAL INFORMATION**  
This product is a premium quality combination product that provides both weed control and the required nutrients for optimum turfgrass growth all in a single product.

This product is a premium quality combination product that provides both weed control and the required nutrients for optimum turfgrass growth all in a single product. It contains Dimension herbicide, a selective herbicide that can be applied up to 4 weeks later than other crabgrass preventers and provides up to 4 months control of crabgrass and other grasses and broadleaf weeds when applied prior to their germination in lawns and ornamental turfgrasses. Apply to well established lawns and ornamental turfgrasses listed below. (See "USE ON THE FOLLOWING TURFGRASSES" section for a list of the tolerant turfgrasses.) This product will not harm nearby established ornamentals when used according to directions given below.

**USE PRECAUTIONS**  
This product should only be applied to lawns and ornamental turfgrasses that are well established. Use of this product on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, pests, excessive temperatures, or lack of moisture, may increase the chances for turfgrass injury. Do not apply to lawns or turfgrasses that are under stress.

**USE ON THE FOLLOWING TURFGRASSES**

- This product can be applied to the following turfgrasses when used as directed:
- |                                                                                                                                                                         |                                                                                                                                                                                   |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Cool-Season Turfgrasses</b>                                                                                                                                          | <b>Warm-Season Turfgrasses</b>                                                                                                                                                    |
| <ul style="list-style-type: none"> <li>Creeping Bentgrass</li> <li>Fine Fescue</li> <li>Kentucky Bluegrass</li> <li>Perennial Ryegrass</li> <li>Tall Fescues</li> </ul> | <ul style="list-style-type: none"> <li>Bahiagrass</li> <li>Bermudagrass</li> <li>Buffalograss</li> <li>Centipedegrass</li> <li>St. Augustinegrass</li> <li>Zoysiagrass</li> </ul> |

This product should be applied in the spring or fall prior to weed seed germination in established and newly germinated lawns and ornamental turfgrasses. For control of other grass and broadleaf weeds, this product must be applied as a preventative treatment before their germination. Only crabgrass will be controlled with delayed applications after germination.

**Tips for Improved Weed Control:** For best results, apply this product within a few days after mowing and delay further mowings after the application until the lawn or ornamental turfgrass has been watered or received significant rainfall to dissolve the fertilizer granules. Improved weed control may be observed when treated lawn or ornamental turfgrass is watered or receives rainfall within a few days after application of this product.

**RESEEDING, OVERSEEDING OR SPRIGGING**

Reseeding, overseeding, or sprigging of treated areas with this product should be delayed until 12 weeks from the time of application. Reseeding, overseeding, or sprigging before 12 weeks after application may prevent establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practice such as soil cultivation, irrigation and fertilization should be followed.

However, if overseeding with perennial ryegrass in the fall, overseed 8 weeks or later after a single application. For best results, use mechanical or power seeding equipment (sift seeders) designed to give good seed to soil contact.

To establish a new lawn of Kentucky Bluegrass, Perennial Ryegrass and/or Tall Fescue, this product may be applied two weeks after the first sign of germination. For other tolerant turfgrasses listed on this label, wait until the grass is well established (2 mowings) before applying this product.

- DO NOT apply to flowers, vegetables or fruit trees.
- DO NOT use clippings from treated turf for mulching around flowers, vegetables or fruit trees.
- DO NOT apply this product to pastures.
- Keep people and pets off treated areas until dust has settled.
- DO NOT apply this product within 8 weeks of seeding.
- DO NOT apply more than 3.82 pounds of this product per 1,000 sq. ft. per application.
- In the State of New York, make only one application per year. In other States do not make more than 3 applications per year spaced at least 2-4 months, apart according to "Region".

**SUGGESTED SPREADER SETTINGS** Recommended Rates in lb per 1,000 sq. ft.

Spreader	3.82 lb
LESCO Commercial (80 pound hopper and larger) Calibration Gauge	#15
Vigoro® 4300	4
Scotts® Broadcast/Rotary Spreaders (Scotts® Basic®, Standard™, Deluxe EdgeGuard® or SpeedyGreen™)	6

**STORAGE AND DISPOSAL**

Storage: Store this product only in its original container in a dry, cool, secure storage area. Do not contaminate water, foodstuffs, feed or seed by storage or disposal.  
Empty Bag Disposal: Do not reuse bags. Dispose of emptied bag(s) in a sanitary landfill, or by incineration.  
Partly Filled Bag Disposal: Call your local solid waste agency or (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.  
Spill: In case of spill, sweep up material and dispose of material according to "Partly Filled Bag Disposal" directions listed above.

**WEEDS CONTROLLED**

This product will provide control of crabgrass and other grasses and broadleaf weeds listed below when applied prior to germination in established lawns and ornamental turfgrasses.

Grass Weeds:	Broadleaf Weeds:		
Annual Bluegrass	Goosegrass	Buttercups	Creeping Oxalis
Annual Ryegrass	Green Foxtail	Buttercup Oxalis	Henbit
Barnyardgrass	Kikuyugrass (seedling)	Carolina Geranium	Parsley-piert
Crabgrass	Perennial Ryegrass	Chickweed	Pineappleweed
Crownvetch	Summer Annual Crabgrass (seedling)	Common Lespedeza	Prostrate Spurge
Yellow Foxtail	Yellow Foxtail	Common Purslane	Spotted Spurge
		Corn Speedwell	Yellow Woodsoot

**WHEN TO APPLY - Application Timings**

This product may be applied either in the spring or in the fall for the control of weeds. When this product is applied in the fall, a sequential or second application should be made the following spring to provide season-long control of weeds.

If this product is applied in the fall, DO NOT reseed, overseed or sprig turfgrasses before 8 weeks after application of this product due to potential turfgrass injury to the newly reseeded, overseeded or sprigged turfgrasses.

When establishing a new lawn of Kentucky Bluegrass, Perennial Ryegrass, and/or Tall Fescue, this product should be applied prior to 8 weeks before seeding, but may be applied two weeks to 8 weeks after the first sign of grass germination.

**LENGTH OF WEED CONTROL**  
This product may provide up to 4 months of weed control if applied according to label directions. The actual length of control will vary depending on use rate, weather conditions, condition of the turfgrass, intensity of weed pressure, and the location or "Region" within the United States where this product is being applied. Please check the map below to determine which "Region" of the U.S. that applies to your particular use area, and find the number of pounds of this product needed in your area based on the length of weed control given in the Recommended Use Rates For This Product.

**POST-EMERGENCE CRABGRASS CONTROL**

For post-emergence crabgrass control, apply this product at the recommended rates when crabgrass is not beyond 3-4 leaf stage. Applications will control existing crabgrass up to the 4 leaf stage and prevent further growth.



- Northern Zone: March to April
- Southern Zone: January to March

**HOW TO APPLY**

**APPLICATION EQUIPMENT AND INSTRUCTIONS**  
This product can be applied with a rotary or drop type spreader designed to apply granular fertilizers. For best results, apply the product evenly and uniformly over the turfgrass area avoiding streaking, skips or overlaps. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions. Adjust the spreader settings such that the spreader delivers the appropriate application rates of this product as recommended below.

A more uniform application can be made by spreading one-half of the recommended amount of this product over the turfgrass area and then applying the remaining one-half of this product at a right angle to the previous application direction. Check application equipment frequently to ensure equipment is functioning properly and applying a uniform distribution of product.

**RECOMMENDED USE RATES FOR THIS PRODUCT**

Region of U.S.	-- -- LENGTH OF WEED CONTROL -- -- (Pounds of this product to apply per 1,000 sq ft of turfgrass)	
	3-4 Months Control	Early Post Emergence Control
North	2.67 (1.74 lb. ai/acre)(lb/acre)	3.82 (0.25 lb. ai/acre)
South/West	3.82 (0.25 lb. ai/acre)	3.82 (0.25 lb. ai/acre)

\*Use the higher rate in areas of high weed pressure.  
A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 sq. ft. per growing season. When used as directed, this product will supplement your regular fertilizer program by providing 0.5 pounds of nitrogen per 1,000 square feet at the 2.67 lb rate or 0.73 pounds of nitrogen per 1,000 square feet at the 3.82 lb rate and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.  
\*Based on Inert Fertilizer claimed.

**CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**  
**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully and completely. It is impossible to eliminate all risks associated with the use of this product. To the extent consistent with applicable law, Buyer and/or User assume all risks of ineffectiveness or other unintended consequences or damages that may result from conditions outside of the control of LESCO, Inc., including but not limited to, such factors as manner of use or application, weather or weather conditions outside the range considered normal at the application site or for the time period in which the product is applied, the presence of other materials, incompatible products, or other influencing factors which are beyond the control of LESCO, Inc., to the extent consistent with applicable law, all such risks shall be assumed by Buyer and/or User, and Buyer and/or User agrees to hold LESCO, Inc., harmless for any claims relating to such factors.

LESCO, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with the Directions for Use under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of LESCO, Inc. and Buyer and/or User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LESCO, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE BUYER AND/OR USER AND THE EXCLUSIVE LIABILITY OF LESCO, INC. FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT) SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LESCO, INC., REPLACEMENT OF THE PRODUCT, OR IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL LESCO, INC. BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

LESCO, Inc. offers this product, and Buyer and/or User accepts it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of LESCO, Inc.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatoryaffairs.com>

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept., 1301 East 9th Street, Suite 1300, Cleveland, OH 44114-1849, referring to the item number found on this bag.

LESCO and Poly Plus are registered trademarks and the sweeping design is a trademark of LESCO Technologies, LLC. Dimension is a registered trademark of Dow AgroSciences. Vigoro is a registered trademark of Unifac Industries Corporation. Scotts, Scotts Basic, Deluxe EdgeGuard and Speedy Green are registered trademarks and Scotts Standard is a trademark of the Scotts Company.

Covers 13,000 Sq. Ft.  
Net Wt. 50 lb (22.7 kg)

EPA Reg. No. 10404-86  
EPA Est. No. 82757-OH-001 (M), 82757-FL-001 (S)  
First letter in lot code indicates manufacturing site.

(101603)  
REV. 2/27/08 KY #080311

Distributed by  
LESCO, Inc.  
1301 East 9th Street  
Cleveland, OH 44114-1849

FOR PRODUCT ASSISTANCE, CALL LESCO AT 1-800-321-5325  
OR VISIT US AT WWW.LESCO.COM



FREIGHT SHIPPING BAG  
Meeting Requirements of  
APPLICABLE FREIGHT CLASSIFICATION  
Guaranteed by  
HOOD PACKAGING CORPORATION  
TYLER, TEXAS

01323 084317  
18 X 29