HOOD • LESCO PRE-M 0.86% 0-0-7 • FRONT
BLACK, BLUE, GREEN, YELLOW

7 58073 52388 8

052388

COVERS 12,500 SQ FT

LESCO®

PRE-M® 0.86%
Crabgrass Pre-emergent
PLUS POTASH

0-0-7

N NITROGEN – Contains no nitrogen.

P PHOSPHATE – Contains no phosphorus.

K POTASH – Promotes strong roots and improves turf hardiness.

Pre-emergent control of crabgrass, goosegrass, foxtail, chickweed, oxalis, spurge, knotweed and *Poa annua.*

PREVENTS ANNUAL GRASSY AND BROADLEAF WEEDS

Crabgrass Control MADE EASY!

ACTIVE INGREDIENT
Dimethylpyridinyl)methyl]-2,6-dinitrobenzenamine*..... 0.86%
OTHER INGREDIENTS..... 99.14%
TOTAL..... 100.00%

*Licensed under one or more of the following U. S. Patents: 5,281,742; 4,586,441.

KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUCIÓN
Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

TM

7 58073 52388 8

Net Wt. 50 lb (22.7 kg)

LESCO® PRE-M® 0.86% Crabgrass Pre-emergent PLUS POTASH

LESCO® PRE-M® 0.86% Crabgrass Pre-emergent PLUS POTASH

LESCO® PRE-M® 0.86% Crabgrass Pre-emergent PLUS POTASH 052388

HOOD • LESCO PRE-M 0.86% 0-0-7 • BACK
BLACK, YELLOW

LESCO®

PRE-M® 0.86% Crabgrass Pre-emergent PLUS POTASH

GUARANTEED ANALYSIS

Soluble Potash (K_2O)	7.00%
Derived from: Muriate of Potash.	
Chlorine (Cl) Max.	5.25%

Made in U.S.A. Produced under the following U.S. Patent 5,750,130

PRE-EMERGENT WEED CONTROL For Use on Turfgrasses
Not for use on turf being grown for sale or for other commercial use as sod, or for commercial seed production, or for research purposes.

KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUCION
Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes eye irritation. Harmful if absorbed through skin. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Avoid breathing dust.

FIRST AID	
If swallowed	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact CHEMTREC at 1-800-424-9300 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. DO NOT apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. DO NOT contaminate water by cleaning of equipment or disposal of equipment washwaters or rinsate.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in possession of the user at time of pesticide application.
- Observe all cautions and limitations in this leaflet and on the labels of products used in combination with this product. Do not use this product other than in accordance with the instructions set forth on this label. The use of this product not consistent with this label can result in injury to turf, animals, or persons. Keep container closed to avoid spills and contamination.
- DO NOT apply this product through any type of irrigation system.
- Do not enter or allow others to enter the treated area until dusts have settled.
- Do not apply this product in a way that will contact any person or pets, either directly or by drift.
- Do not allow dust to drift from the application site and contact people, structures people occupy at the time and the associated property, parks and recreation areas, nontarget crops, aquatic and wetland areas, woodlands, pastures, rangelands, or animals.

APPLICATION RATES
RATE AND APPLICATION CHART

Turfgrass Species	Weeds	Rates of PRE-M 0.86% Crabgrass Pre-emergent Plus Potash	Comments:
NORTHERN GRASSES Bentgrasses Perennial ryegrass Fine fescue Tall fescue	crabgrass foxtail *Poa annua* barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed evening primrose hop clover	4.0 to 5.3 lbs/1,000 sq ft as the initial application prior to weed germination in the spring.	Apply a repeat application of 2.7 to 4.0 lbs/1,000 sq ft after a minimum of 4 weeks for extended control or where heavy weed infestations are expected.
	goosegrass	4.0 to 5.3 lbs/1,000 sq ft in the spring prior to weed germination.	Apply a repeat application of 4.0 lbs/1,000 sq ft.
	cudweed *Poa annua* chickweed lawn burweed henbit corn speedwell	4.0 to 5.3 lbs/1,000 sq ft.	Apply in late summer or early fall prior to weed germination.
Bentgrass and perennial ryegrass *Poa annua* (1/2 inch height or taller)	crabgrass foxtail *Poa annua* barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed evening primrose hop clover	4.0 lbs/1,000 sq ft in the spring prior to weed germination.	Apply a repeat application of 2.7 to 4.0 lbs/1,000 sq ft after a minimum of 4 weeks for extended control or where heavy weed infestations are expected.
	goosegrass	4.0 lbs/1,000 sq ft prior to weed germination in the spring.	Apply a repeat application of 4.0 lbs/1,000 sq ft after a minimum of 4 weeks.
	cudweed *Poa annua* chickweed lawn burweed henbit corn speedwell	4.0 lbs/1,000 sq ft.	Apply in late summer or early fall prior to weed germination.
SOUTHERN GRASSES Bermudagrass* St. Augustinegrass Bahiagrass Zoysiagrass Centipedegrass Tall fescue	crabgrass foxtail *Poa annua* barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed evening primrose hop clover	4.0 to 5.3 lbs/1,000 sq ft prior to weed germination in the spring.	Apply a repeat application of 2.7 to 4.0 lbs/1,000 sq ft after a minimum of 4 weeks if necessary.
	goosegrass	4.0 lbs/1,000 sq ft prior to weed germination in the spring. Apply a repeat application 4.0 to 5.3 lbs/ 1,000 sq ft after a minimum of 4 weeks.	An additional application of 4.0 lbs/1,000 sq ft may be applied for extended goosegrass control a minimum of 4 weeks after the second application.
	cudweed *Poa annua* chickweed lawn burweed henbit corn speedwell	4.0 to 5.3 lbs/1,000 sq ft.	Apply in late summer or early fall prior to weed germination.

*Not for use on greens or tees

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.
PESTICIDE STORAGE: Store this product only in its original container in a dry, cool, secured storage area. Do not contaminate water, foodstuffs, feed or seed by storage or disposal.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, State or local procedures. Or call [1-800-CLEANUP] for disposal instructions. Never place unused product down any indoor or outdoor drain.
CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill of by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.
SPILL: In case of spill, sweep up material and dispose of material according to "pesticide disposal" directions listed above.

GENERAL INFORMATION

PRE-M 0.86% Crabgrass Pre-emergent Plus Potash is a combination pre-emergent herbicide and fertilizer formulated to provide control of most annual grasses and certain broadleaf weeds found in turfgrass. **PRE-M 0.86% Crabgrass Pre-emergent Plus Potash** is labeled for use on the following turfgrass species: Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, bentgrass, *Poa annua* (except putting greens and tees), bermudagrass, St. Augustinegrass, bahiagrass, centipedegrass, and zoysiagrass.

PRE-M 0.86% Crabgrass Pre-emergent Plus Potash controls weeds as they germinate but will not control established weeds.

Unusually cold, excessively wet, or hot and dry conditions that delay germination or extend germination over a long period of time can reduce weed control.

Applied according to label directions and under normal growing conditions, **PRE-M 0.86% Crabgrass Pre-emergent Plus Potash** will not cause turf injury. Overapplication can result in turf stand loss, turf injury, or soil residues. Uneven application can decrease weed control or cause turf injury.

Seedling diseases, cold weather, excessive moisture, high soil pH, high soil salt concentration, or drought can weaken turfgrass and increase the possibility of turf damage from PRE-M 0.86% Plus Potash.

APPLICATION

Apply product uniformly at the recommended rate using a properly calibrated spreader. The efficacy of **PRE-M 0.86% Crabgrass Pre-emergent Plus Potash** will be improved if the application is followed by one-half inch of rainfall or its equivalent in sprinkler irrigation. This will remove the product from the turfgrass foliage and establish a pre-emergent herbicide barrier in the soil.

Avoid contact of product with porous surfaces such as driveways, stone, or wood. Removing particles will help prevent staining.

SUGGESTED SPREADER SETTINGS

Recommended Rates in lbs per 1,000 sq ft

Spreader	Ground Speed	Spreading Width	2.7#	4.0#*	5.3#**
LESCO Commercial (80 pound hopper or larger)	3 mph	9 ft.			
	LESCO Calibration Gauge		#13	#15	#17
Vigoro® 4300			3	3.5	4
Scotts® Broadcast/Rotary Spreaders (Scotts® Basic®, Standard™, Deluxe EdgeGuard® or SpeedyGreen®)			5	5.5	7

*Delivers 1.6 lb ai/A **Delivers 2.0 lb ai/A

IMPORTANT: Spreader settings are only approximate. Age, condition of spreader, and operator speed can cause wide variation in application rate. Spreader should be calibrated prior to each application by the operator.

The efficacy of this product will be improved if the application is followed by one-half inch of rainfall or its equivalent in sprinkler irrigation. If this product is not activated by rainfall or irrigation within 14 days, erratic weed control may result.

PRECAUTIONS

Use only on well established turfgrass with a dense and uniform stand. On newly planted areas, application should not be made until the turfgrass has filled in and has been mowed at least four times.

On turf that has been thinned or damaged due to winter injury, excessive moisture, etc., allow for turf recovery prior to making an application.

Applications made to overseeded warm-season turfgrasses may cause thinning or injury of the overseeded species.

Do not use on bentgrass or *Poa annua* (greens and tees) or injury may occur.

Delay reseeding or winter overseeding treated turfgrass for at least 2 months following application. Do not sprig turfgrass for 3 months after application.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully and completely. It is impossible to eliminate all risks inherent in the application of this product. Crop injury, ineffectiveness or other unintended consequences may be expected despite the care that can be taken by LESCO, Inc. and/or User. Many factors beyond the control of LESCO, Inc. and/or User affect the application, performance, and/or yield of this product. These factors include but not limited to, such factors as manner of use or application, weather or weather conditions outside the range considered normal at the application site or for the time period in which the product is applied, the presence of other materials, incompatibility of other materials, and other factors outside the control of LESCO, Inc. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and/or User, and Buyer and/or User agrees to hold LESCO, Inc. harmless for any claims relating to such factors.

LESCO, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with the Directions for Use under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of LESCO, Inc. and Buyer and/or User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LESCO, INC. MAKES NO WARRANTIES OR GUARANTIES EXPRESS OR IMPLIED, OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE BUYER AND/OR USER AND THE EXCLUSIVE LIABILITY OF LESCO, INC. FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, NEGLIGENCE, MISUSE, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LESCO, INC. REPLACEMENT OF THE PRODUCT, OR IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL LESCO, Inc., BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

LESCO, Inc. offers this product, and Buyer and/or User accepts it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of LESCO, Inc.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn. RA Dept., 1301 East 9th Street, Suite 1300, Cleveland, Ohio 44114-1849, referring to the number found on this bag.

LESCO and PRE-M are registered trademarks and the sweeping design is a trademark of LESCO Technologies, LLC. Vigoro is a registered trademark of United Industries Corporation. Scotts, Scotts Basic, Deluxe EdgeGuard and Speedy Green are registered trademarks and Scotts Standard is a trademark of The Scott Company.

Covers 12,500 Sq. Ft.

Net Wt. 50 lb (22.7 kg)

EPA REG. NO. 10404-100
EPA EST. NO. 82757-FL-001 (S), 82757-OH-001 (M)
First letter of lot code indicates manufacturing site

(113004)
Rev. 11/3/09 VT **#052388**

F1560
Distributed by
LESCO, Inc.
1301 East 9th Street
Cleveland, OH 44114-1849

7 58073 52388 8

FOR PRODUCT ASSISTANCE, CALL LESCO AT 1-800-321-5325
www.lesco.com

FREIGHT SHIPPING BAG
Meeting Requirements of
APPLICABLE FREIGHT CLASSIFICATION
Guaranteed by
HOOD PACKAGING CORPORATION
TYLER, TEXAS

01762 084424
15 X 28