

44752AL0022


Agri Star®

BIFENAMITE 4SC

**Agricultural Miticide
For Agricultural Use Only**

ACTIVE INGREDIENT:	(% by weight)
bifenazate: hydrazine carboxylic acid, 2-(4-methoxy- [1,1'-biphenyl]-3-yl) 1-methylethyl ester	43.2%
OTHER INGREDIENTS:	56.8%
TOTAL:	100.0%

Contains 4 pounds active ingredient bifenazate per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF INHALED:	<ul style="list-style-type: none"> • Move the person to fresh air. • If the person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For chemical emergency: spill, leak, fire, exposure or accident, call CHEMTREC: 1-800-424-9300.

See inside booklet for additional Precautionary Statements and Directions For Use.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Harmful if inhaled. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

GROUP 20D ACARICIDE

USER SAFETY RECOMMENDATIONS

Users should:

1. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
2. Remove clothing/PPE immediately if pesticide gets inside. Then, wash thoroughly and change into clean clothing.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing them. As soon as possible, wash thoroughly and change clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds, estuarine/marine invertebrates and fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate. This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are foraging the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a dry location.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

[For rigid, non-refillable containers (≤ 5 gallons):] Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake container for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer container for recycling, if available, or puncture and dispose of container in a sanitary landfill, or by incineration.

[For rigid, non-refillable containers (> 5 gallons):] Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer container for recycling, if available, or puncture and dispose of container in a sanitary landfill, or by incineration.

EPA Reg. No. 42750-321

EPA Est. No. 42750-MO-002

Manufactured For:

ALBAUGH, LLC

1525 NE 36th Street • Ankeny, Iowa 50021

PEEL HERE TO OPEN ▲

8



3

NET CONTENTS: 2.5 GALLONS

PF-134941

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Harmful if inhaled. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and Other Handlers Must Wear:

1. Long-sleeved shirt and long pants,
2. Shoes plus socks, and
3. For mixing and loading activities when not using a closed system, wear an apron and chemical-resistant gloves made of: barrier laminate, polyethylene, butyl rubber (>14 mils), nitrile rubber (>14 mils), neoprene rubber (>14 mils), natural rubber (>14 mils), polyvinyl chloride (>14 mils), or Viton (>14 mils).

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

1. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
2. Remove clothing/PPE immediately if pesticide gets inside. Then, wash thoroughly and change into clean clothing.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing them. As soon as possible, wash thoroughly and change clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds, estuarine/marine invertebrates and fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate. This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are foraging the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame. Do not mix or allow to come into contact with oxidizing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exceptions are listed under the USE INSTRUCTIONS for each crop. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

1. Coveralls
2. Shoes plus socks
3. Chemical-resistant gloves made of barrier laminate, polyethylene, butyl rubber (≥14 mils), nitrile rubber (≥14 mils), neoprene rubber (≥14 mils), natural rubber (≥14 mils), polyvinyl chloride (≥14 mils), or Viton (≥14 mils).

USE INFORMATION

BIFENAMITE 4SC provides both knockdown action as well as extended residual control of the following mites when applied directly to foliage as directed by this label:

Avocado red spider	Clover	Pecan leaf scorch	Spruce spider mite
Banks grass	European red*	Persea	Strawberry spider
Brown almond	McDaniel	Sixspotted	Two-spotted spider
Citrus red	Pacific spider	Southern red mite	Willamette

* using maximum rate specified for the crop or crop group.

This prod coverage

This prod

- 1) Fill spr
- 2) Begin
- 3) Fill the

NOTE: TI pH 5.5 - 6

Tank Mixe recomme where all for the int must follo

BIFENAM developm If mite pre est specif applicatio

This prod INSTRUC

This prod canopy or the minim

Chemigat operated tables for TATO, AN

USE RES

- Do NC bifena
- Do NC applie
- When
- Do NC
- Do NC

If mite pr highest sp

* A mitico

This product is appropriate for use in IPM and resistance management programs. Because this product is *not* systemic, effective control requires complete coverage of both upper and lower leaf surfaces.

MIXING INSTRUCTIONS

This product is a flowable suspension concentrate, users must be sure to stir or shake this product well prior to use.

- 1) Fill spray tank with 1/2 the desired amount of water.
- 2) Begin agitation and add the required amount of BIFENAMITE 4SC.
- 3) Fill the tank with the remaining amount of required water and continue agitation until spraying is completed.

NOTE: The stability of this product can be affected by high pH and high temperature. Spray mixtures containing BIFENAMITE 4SC should be kept within pH 5.5 - 6.5.

Tank Mixes: To increase the number of insect and/or mite species controlled, this product may be tank - mixed with other insecticides. *It is strongly recommended that compatibility with other tank-mix partners be confirmed prior to broad-scale applications.* Tank mixtures are only permitted in States where all of the pesticide products used as tank-mix partners are registered. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

USE RATES AND DIRECTIONS

BIFENAMITE 4SC controls the mite species listed above and is especially effective against grass, red and spider mites, primarily in the motile stage of development but does provide ovicidal control of spider mites (Tetranychus spp.) as well. Note that this product does NOT control broad, flat or rust mites. If mite pressure is not heavy, the lowest specified rates listed should be used. If mite pressure is significant or to provide extended residual control, the highest specified rates listed should be used. In order to maximize residual control, applications should be made as soon as mites are detected. For specific application rates, application numbers, and Pre-Harvest Intervals (PHI), refer to the appropriate USE INSTRUCTIONS tables below.

This product may be ground-applied using the following types of equipment: air-blast sprayers, compressed air, or hydraulic ground booms. The USE INSTRUCTIONS tables below indicate the minimum numbers of gallons of spray solution to apply per acre for ground applications.

This product may be applied aerially on certain crops using either a fixed-wing aircraft or a helicopter. A minimum spray volume that ensures complete canopy coverage must be used. Refer to the USE INSTRUCTIONS tables for the minimum numbers of gallons of spray solution to apply per acre (or apply the minimum gallons/acre allowed by your State, which may not be less than the minimum gallons/acre shown on this label).

Chemigation (Cranberry, Mint, Potato, and Timothy, only): Only one chemigation application may be made per calendar year, and the system must be operated at 80% to 100% during the application to apply the minimum amount of water possible. Refer to the USE INSTRUCTIONS - BEARING CROPS tables for the ranges in application rates permitted and to the CHEMIGATION USE PRECAUTIONS AND INSTRUCTIONS FOR CRANBERRY, MINT, POTATO, AND TIMOTHY.

USE RESTRICTIONS

- Do NOT plant another crop within 30 days after last application of this product. Planting within 30 days of last application increases the likelihood of bifenazate residues appearing in rotational crops.
- Do NOT exceed the maximum amount of bifenazate allowed per crop per calendar year. This applies to *all* product(s) containing bifenazate that are applied to the crop in a calendar year.
- When applying to Golden Delicious apples, do NOT tank-mix oil with this product.
- Do NOT use adjuvants in tank mixtures that will be applied to greenhouse tomatoes.
- Do NOT apply to greenhouse tomatoes within 3 days of harvest (3-day PHI).

USE INSTRUCTIONS - BEARING CROPS

(crops that will bear fruit within one year of application)

If mite pressure is not heavy, the lowest specified rates listed should be used. If mite pressure is significant or to provide extended residual control, the highest specified rates listed should be used.

AVOCADO

Application Rate (fl. oz. / Acre)	Minimum Gallons of Water per Acre		Acre Inches of Water for Chemigation	Maximum Number of Applications per Year	Minimum Number of Days Between Applications	Pre-Harvest Interval (PHI) in Days
	Ground	Air				
12 - 16	80	-	-	2	21*	7

* A miticide with a different mode of action should be used between any 2 applications of BIFENAMITE 4SC.

Blackberry

A

* A miticide

A

* Minimum

A

* Minimum

Cucumber
nese prese
apple bals:

A

* Minimum

¹Edible Go
²Muskmeic
honey ballt
³Squash, s
⁴Squash, v
ghetti squa

Cultivars o
(dry), navy
rice bean,

A

* Minimum

** A miticide

complete

CANEBERRY SUBGROUP 13-07A:

Blackberry; loganberry; red and black raspberry and cultivars and/or hybrids of these; Wild Raspberry

Application Rate (fl. oz. / Acre)	Minimum Gallons of Water per Acre		Acre Inches of Water for Chemigation	Maximum Number of Applications per Year	Minimum Number of Days Between Applications	Pre-Harvest Interval (PHI) in Days
	Ground	Air				
12 - 16	50	-	-	2	30*	1

* A miticide with a different mode of action should be used between any 2 applications of BIFENAMITE 4SC.

ept within

CHRISTMAS TREES / CONIFER PLANTATIONS AND NURSERIES

Application Rate (fl. oz. / Acre)	Minimum Gallons of Water per Acre		Acre Inches of Water for Chemigation	Maximum Number of Applications per Year	Minimum Number of Days Between Applications	Pre-Harvest Interval (PHI) in Days
	Ground	Air				
12 - 16	100	10*	-	1	-	-

* Minimum gallons per acre OR the minimum permitted by your state, but not less than shown.

s strongly
in States
registered
ng. Usersstage of
ust mites.
the high-
r specific

The USE

complete
(or apply1 must be
3 CROPS
/INT, PO-

likelihood of

e that are

ontrol, the

arvest
rval
-ll)
ays
7**COTTON**

Application Rate (fl. oz. / Acre)	Minimum Gallons of Water per Acre		Acre Inches of Water for Chemigation	Maximum Number of Applications per Year	Minimum Number of Days Between Applications	Pre-Harvest Interval (PHI) in Days
	Ground	Air				
16 - 24	20	5*	-	1	-	60

* Minimum gallons per acre OR the minimum permitted by your state, but not less than shown.

CUCURBIT VEGETABLES:Cucumbers; Edible Gourds¹; Muskmelon²; Pumpkin; Squash (summer³ and winter⁴), Watermelon; Chayote (fruit) (*Sechium edule*), Chinese waxgourd (Chinese preserving melon) (*Benincasa hispida*), Citrus melon (*Citrullus lanatus* var. *citroides*), Gherkin (*Cucumis anguria*), *Momordica* spp. (includes balsam apple balsam pear, bitter melon, Chinese cucumber

Application Rate (fl. oz. / Acre)	Minimum Gallons of Water per Acre		Acre Inches of Water for Chemigation	Maximum Number of Applications per Year	Minimum Number of Days Between Applications	Pre-Harvest Interval (PHI) in Days
	Ground	Air				
12 - 16	50	10*	-	1	-	3

* Minimum gallons per acre OR the minimum permitted by your state, but not less than shown.

¹Edible Gourd (*Lagenaria* spp.) (includes *hyotan* and *cucuzza*), and (*Luffa acutangula* and *L. cylindrica*) which includes hechima and Chinese okra.²Muskmelon (hybrids and/or cultivars of *Cucumis melo*) includes true cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon.³Squash, summer (*Cucurbita pepo* var. *melo*) includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini.⁴Squash, winter (*Cucurbita maxima*: *C. moschata*) includes butternut squash, calabaza, hubbard squash; (*C. mixta*; *C. pepo*) includes acorn squash, spaghetti squash.**DRIED SHELLED BEAN (EXCEPT SOYBEAN) SUBGROUP:**Cultivars of *Lupinus* spp. include grain lupin, sweet lupin, white lupin, and white sweet lupin; *Phaseolus* spp. include field bean, kidney bean, lima beans (dry), navy bean, and pinto bean; tepary bean; *Vigna* spp., including adzuki bean, blackeyed pea, catjang, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, and urd bean; broad bean (dry); chickpea; guar; lablab bean; and lentil.

Application Rate (fl. oz. / Acre)	Minimum Gallons of Water per Acre		Acre Inches of Water for Chemigation	Maximum Number of Applications per Year	Minimum Number of Days Between Applications	Pre-Harvest Interval (PHI) in Days
	Ground	Air				
16 - 24	20	7*	-	2	14**	7

* Minimum gallons per acre OR the minimum permitted by your state, but not less than shown.

** A miticide with a different mode of action should be used between any 2 applications of BIFENAMITE 4SC.

Eggplant; O
tomato; gar
pepper, chil
hybrids of tApplie
(fl. o
1* Minimum g
1 Pepper (CaS
Grape'; AmApplie
(fl. o
11 Grapes: Th
may enter o
as at least l
tasks nameApplie
(fl. o
1Bean (*Lupin*
beans, navy
catjang, Chi
(garbanzo);
pea, sugar
snow pea; (C
Porto Rico pApplie
(fl. o
1* Minimum g
** A miticideApplie
(fl. o
1* Minimum g
** Refer to U

FRUITING VEGETABLES, GROUP 8-10:

Eggplant; Okra; Peppers (including all varieties of *Capsicum* spp.); Tomatoes (field grown); African eggplant; bush tomato; bell pepper; cocona; currant tomato; garden huckleberry; goji berry; ground cherry (*Physalis* spp.); marjynia; naranjilla; pea eggplant; pepino; pepper (includes bell pepper, nonbell pepper, chili pepper, hot cooking pepper, pimento, sweet pepper), roselle; scarlet eggplant; sunberry; tomatillo; tree tomato; cultivars, varieties and/or hybrids of these.

Application Rate (fl. oz. / Acre)	Minimum Gallons of Water per Acre		Acre Inches of Water for Chemigation	Maximum Number of Applications per Year	Minimum Number of Days Between Applications	Pre-Harvest Interval (PHI) in Days
	Ground	Air				
12 - 16	50	10*	-	1	-	3

* Minimum gallons per acre OR the minimum permitted by your state, *but not less than shown*.

† Pepper (*Capsicum* spp.) includes bell peppers, chili peppers, cooking peppers, pimento, and sweet peppers.

SMALL FRUIT VINE CLIMBING SUBGROUP 13-07F, EXCEPT FUZZY KIWIFRUIT:

Grape¹; Amur River Grape; Gooseberry; Kiwifruit, hardy; Maypop; Schisandra Berry; Cultivars, varieties and/or hybrids of these.

Application Rate (fl. oz. / Acre)	Minimum Gallons of Water per Acre		Acre Inches of Water for Chemigation	Maximum Number of Applications per Year	Minimum Number of Days Between Applications	Pre-Harvest Interval (PHI) in Days
	Ground	Air				
12 - 16	50	-	-	1	-	14

¹Grapes: The REI is 5 days for cane turning, tying and girdling of table grapes. Exception: In addition to the early entry exceptions allowed by WPS, you may enter or allow workers to enter treated areas to perform all tasks other than tying, turning and girdling after 12 hours following applications, as long as at least long pants, long-sleeved shirts shoes and socks are worn. Notify workers of the exception (including when entry is permitted for each of the tasks named in the exception).

HOPS

Application Rate (fl. oz. / Acre)	Minimum Gallons of Water per Acre		Acre Inches of Water for Chemigation	Maximum Number of Applications per Year	Minimum Number of Days Between Applications	Pre-Harvest Interval (PHI) in Days
	Ground	Air				
12 - 24	50	-	-	1	-	14

**LEGUME VEGETABLES SUBGROUP 6A (succulent) SUCCULENT PEAS
and BEANS SUBGROUP 6B; SUCCULENT SHELLED SOYBEAN:**

Bean (*Lupinus* spp.) (includes grain lupin, sweet lupin, white lupin, and white sweet lupin); bean (*Phaseolus* spp.) (includes field bean, kidney bean, lima beans, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean); bean (*Vigna* spp.) (includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean); broadbean (fava); chickpea (garbanzo); guar; jackbean; lablab bean; lentil; pea (*Pisum* spp.) (includes dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea); pigeon pea; soybean (immature seed); sword bean; sugar pea; Chinese pea; pois mange tout; snap pea; ming pea; podded pea; snow pea; China pea; chicharo; shi hia wandou; saya-endo; sugar snap pea; Congo pea; no-eye pea; red gram; arher; gandule; dhal; toor; gunds pea; Porto Rico pea; urher gandul; guandu; pois-d'angole; gungo pea.

Application Rate (fl. oz. / Acre)	Minimum Gallons of Water per Acre		Acre Inches of Water for Chemigation	Maximum Number of Applications per Year	Minimum Number of Days Between Applications	Pre-Harvest Interval (PHI) in Days
	Ground	Air				
16 - 24	20	10*	-	2	14**	3

* Minimum gallons per acre OR the minimum permitted by your state, *but not less than shown*.

** A miticide with a different mode of action should be used between any 2 applications of BIFENAMITE 4SC.

MINT

Application Rate (fl. oz. / Acre)	Minimum Gallons of Water per Acre		Acre Inches of Water for Chemigation	Maximum Number of Applications per Year	Minimum Number of Days Between Applications	Pre-Harvest Interval (PHI) in Days
	Ground	Air				
12 - 24	50	10*	0.1 - 0.2**	1	-	7

* Minimum gallons per acre OR the minimum permitted by your state, *but not less than shown*.

** Refer to USE RATES AND DIRECTIONS and CHEMIGATION USE PRECAUTIONS AND INSTRUCTIONS sections.

Apples; Crab

Appl
(fl.)

Tuberous and
chufa; dashee

Appl
(fl.)

* Minimum ga
** Refer to US
*** A miticide

Strawberry; B
these.

Appl
(fl.)

* Refer to US†
** A miticide v

Apricots; Che
Plum (*Prunus*

Appl
(fl.)

Appl
(fl.)

* Refer to US†

POME FRUIT, GROUP 11-10:

Apples; Crabapples; Pears; Quince; azarole; medlar; pear, Asian; quince, Chinese; quince, Japanese; tejocote; cultivars, varieties and/or hybrids of these.

Application Rate (fl. oz. / Acre)	Minimum Gallons of Water per Acre		Acre Inches of Water for Chemigation	Maximum Number of Applications per Year	Minimum Number of Days Between Applications	Pre-Harvest Interval (PHI) in Days
	Ground	Air				
12 - 16	50	-	-	1	-	7

POTATOES

Tuberous and corm vegetables: arracacha; arrowroot; artichoke, Chinese; artichoke, Jerusalem; canna, edible; cassava, bitter and sweet; chayote (root); chufa; dasheen (taro); ginger; leren; potato; sweet potato; taniar; turmeric; yam bean; yam, true.

Application Rate (fl. oz. / Acre)	Minimum Gallons of Water per Acre		Acre Inches of Water for Chemigation	Maximum Number of Applications per Year	Minimum Number of Days Between Applications	Pre-Harvest Interval (PHI) in Days
	Ground	Air				
16 - 24	20	10*	0.1 - 0.2**	2	14***	14

* Minimum gallons per acre OR the minimum permitted by your state, *but not less than shown*.

** Refer to USE RATES AND DIRECTIONS and CHEMIGATION USE PRECAUTIONS AND INSTRUCTIONS sections.

*** A miticide with a different mode of action should be used between any 2 applications of BIFENAMITE 4SC.

LOW GROWING BERRY SUBGROUP 13-07G;

Strawberry; Bearberry; Bilberry; Blueberry, lowbush; Cloudberry; Cranberry; Lingonberry; Muntries; Partridgeberry; Cultivars, varieties, and/or hybrids of these.

Application Rate (fl. oz. / Acre)	Minimum Gallons of Water per Acre		Acre Inches of Water for Chemigation	Maximum Number of Applications per Year	Minimum Number of Days Between Applications	Pre-Harvest Interval (PHI) in Days
	Ground	Air				
12 - 16	100	-	0.1 - 0.2 * (Cranberry only)	2	21*	1

* Refer to USE RATES AND DIRECTIONS and CHEMIGATION USE PRECAUTIONS AND INSTRUCTIONS sections.

** A miticide with a different mode of action should be used between any 2 applications of BIFENAMITE 4SC.

STONE FRUIT:Apricots; Cherries (sweet & tart); Nectarines; Peaches; Plums/Prunes (*Prunus domestica*, *Prunus* spp.), Chickasaw Plum (*Prunus augustifolia*), Damson Plum (*Prunus domestica* spp. *insititia*), Japanese Plum (*Prunus salicina*), and Plumcot (*Prunus armeniaca* X *P. domestica*)

Application Rate (fl. oz. / Acre)	Minimum Gallons of Water per Acre		Acre Inches of Water for Chemigation	Maximum Number of Applications per Year	Minimum Number of Days Between Applications	Pre-Harvest Interval (PHI) in Days
	Ground	Air				
12 - 16	80	-	-	1	-	3

TIMOTHY - forage and hay fed to horses ONLY.*(Applications are only permitted in Eureka and Humboldt Counties in Nevada)*

Application Rate (fl. oz. / Acre)	Minimum Gallons of Water per Acre		Acre Inches of Water for Chemigation	Maximum Number of Applications per Year	Minimum Number of Days Between Applications	Pre-Harvest Interval (PHI) in Days
	Ground	Air				
16 - 24	20	10	0.1 - 0.2*	1	-	Forage - 0 Hay - 7

* Refer to USE RATES AND DIRECTIONS and CHEMIGATION USE PRECAUTIONS AND INSTRUCTIONS sections.

TOMATOES:

Grown in Greenhouses ONLY

Application Rate (fl. oz. / Acre)	Minimum Gallons of Water per Acre		Acre Inches of Water for Chemigation	Maximum Number of Applications per Year	Minimum Number of Days Between Applications	Pre-Harvest Interval (PHI) in Days
	Ground	Air				
2 - 4	100*	-	-	2	-	3

* Apply 1-4 quarts of mixture per 100 square feet (100-400 gallons per acre) to the foliage using ground spray equipment (including air blast, compressed air, ground boom, hose-end and hydraulic sprayers). The spray volume used is dependent on the size of the plants being treated, with larger plants requiring greater spray volumes.

TREE NUTS: Almonds

Application Rate (fl. oz. / Acre)	Minimum Gallons of Water per Acre		Acre Inches of Water for Chemigation	Maximum Number of Applications per Year	Minimum Number of Days Between Applications	Pre-Harvest Interval (PHI) in Days
	Ground	Air				
12 - 24	50	-	-	1	-	7

TREE NUTS:

Beech nut; Brazil nut; Butternut; Cashew; Chestnut; Chinquapin; Filbert (Hazelnut); Hickory nut; Macadamia nut (bush nut); Pecans; Pistachios; Walnut (black & English)

Application Rate (fl. oz. / Acre)	Minimum Gallons of Water per Acre		Acre Inches of Water for Chemigation	Maximum Number of Applications per Year	Minimum Number of Days Between Applications	Pre-Harvest Interval (PHI) in Days
	Ground	Air				
12 - 24	50	-	-	1	-	14

TROPICAL FRUIT:

Papaya; Star Apple; Black Sapote; Mango; Sapodilla; Canistel; Mamey Sapote; Lychee; Longan; Spanish Lime; Rambutan; Pulasan; Guava; Feijoa; Jaboticaba; Wax Jambu; Starfruit (Carambola); Passionfruit; Acerola; Sugar Apple; Cherimoya; Atemoya; Custard Apple; Ilama; Soursap; Biriba

Application Rate (fl. oz. / Acre)	Minimum Gallons of Water per Acre		Acre Inches of Water for Chemigation	Maximum Number of Applications per Year	Minimum Number of Days Between Applications	Pre-Harvest Interval (PHI) in Days
	Ground	Air				
12 - 16	50	-	-	2	21*	1

* A miticide with a different mode of action should be used between any 2 applications of BIFENAMITE 4SC.

USE INSTRUCTIONS - NON-BEARING CROPS

This product may be used on non-bearing crops (defined as crops that will not bear fruit within one year of application), including all crops listed for the USE INSTRUCTIONS - BEARING CROPS tables as well as the Additional Crops listed below. This use includes application to non-bearing fruit trees and berries in commercial nurseries and tree plantations.

Non-bearing crop applications may be used to control the following mites:

Banks grass	European red*	Southern red mite	Willamette
Brown almond	McDaniel	Spruce spider mite	
Citrus red	Pacific spider	Strawberry spider	
Clover	Pecan leaf scorch	Two-spotted spider	

* using the maximum rate specified for the crop or crop group.

USE RESTRICTIONS FOR NON-BEARING CROPS

- Use ground application equipment ONLY.
- The non-bearing crops listed below have a 12-hour restricted entry interval (REI).

Additional Crops: Berries (blueberry, highbush; elderberry; huckleberry); Citrus (grapefruit, lemons, limes, oranges, tangerines, etc.); Currants; Dates; Figs; Persimmons

Application Rate (fl. oz. / Acre)	Minimum Gallons of Water per Acre	Maximum Number of Applications Per calendar Year
12 - 16	50	1

This product plants, foliage

This product especially effective

Bamboo Sp
Citrus Red M
Clover Mite
European R
Lewis Mite t

NOTE: This

Because this required. W to beneficial

Use Rates

Apply as so use the low residual con

Mix 2-4 fluid feet (100-40 sprayers). T

Coverage of ounces per in a small-sc

USE RESTRICTIONS

- Do NOT
- Do NOT
- Do NOT

CHEMIGATION

- Apply thi
- move irri
- Crop inju
- If you ha
- Do not c
- label-pre
- A persor
- the syste
- The syst
- vent wat
- The pest
- pump.
- The pest
- pump an
- ically or i
- The syst
- The irrig
- to the pc
- Systems
- materials
- Do not a
- Maintain
- To insure
- Applying
- from leaf
- Do NOT
- this prod
- Apply thi

USE INSTRUCTIONS – GREENHOUSES AND SHADEHOUSES

This product may be applied to foliage for knockdown and residual control of mites on ornamental plants (including ornamental bulb crops, bedding plants, foliage and flowering plants, perennials, shrubs and trees). The plants may be grown in the ground or in pots.

This product works most effectively on mites that are motile, but also has ovicidal activity against the *Tetranychus* spp. mites (i.e., Spider Mites) and is especially effective against grass mites, red mites and spider mites. Common mite species controlled by BIFENAMITE 4SC include:

Bamboo Spider Mite *Schizotetranychus celarius* (Banks)
Citrus Red Mite *Panonychus citri* (McGregor)
Clover Mite *Bryobia praetiosa* (Koch)
European Red Mite *Panonychus ulmi* (Koch)
Lewis Mite *Eotetranychus lewisii*

Pacific Mite *Tetranychus pacificus* (McGregor)
Southern Red Mite *Oligonychus ilicis* (McGregor)
Spruce Spider Mite *Oligonychus ununguis* (Jacobi)
Strawberry Mite *Tetranychus turkestanii*
Two-Spotted Spider Mite *Tetranychus urticae* (Koch)

NOTE: This product is not effective in controlling rust mites, broad mites and flat mites.

Because this product is not a systemic insecticide, thorough coverage of both the upper and lower surfaces of the leaves, as well as any fruit present, is required. When used as directed, this product is useful in resistance management and IPM programs due to its selective nature that minimizes impacts to beneficial insects.

Use Rates and Application Information

Apply as soon as mites appear. BIFENAMITE 4SC provides residual control for up to 28 days. Within the range of rates specified in the next paragraph, use the lower rates when mite pressure is low or for preventative applications; and use the higher rates when mite pressure is severe or for extended residual control.

Mix 2-4 fluid ounces of BIFENAMITE 4SC per 100 gallons of water (1/8 – 1/4 teaspoon per gallon of water). Apply 1-4 quarts of the mixture per 100 square feet (100-400 gallons per acre) to the foliage using ground spray equipment (including air blast, compressed air, ground boom, hose-end and hydraulic sprayers). The spray volume used is dependent on the size of the plants being treated, with larger plants requiring greater spray volumes.

Coverage of ornamentals with hard-to-wet foliage may be improved by using an adjuvant to the spray mixture. Silwet 77® or Sylgard 309® at a rate of 4 ounces per 100 gallons of water has been found to be effective. As with any tank mixture, the compatibility and safety of any adjuvants should be tested in a small-scale trial prior to widespread use.

USE RESTRICTIONS:

- Do NOT apply BIFENAMITE 4SC more than two times per calendar year.
- Do NOT apply more than 16 fluid ounces of BIFENAMITE 4SC per acre per calendar year.
- Do NOT apply BIFENAMITE 4SC by aerial application.

CHEMIGATION USE PRECAUTIONS AND INSTRUCTIONS FOR CRANBERRY, MINT, POTATOES AND TIMOTHY

- Apply this product only through sprinkler systems, including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make the necessary adjustments should the need arise.
- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.
- Maintain constant agitation in the supply tank for the duration of the application.
- To insure complete mixing of the spray solution, use a positive displacement pump to inject the solution before a right angle turn in the main line.
- Applying more than the recommended amount of irrigation water per acre may result in decreased performance due to the chemical being flushed from leaf surfaces.
- Do NOT apply this product if there are leaks in any of the fittings or connections, if nozzles do not provide uniform distribution, or if lines containing this product will be drained and dismantled.
- Apply this product for the entire duration of the irrigation process. Calibration accuracy and product distribution will be improved if a larger volume of

- a more dilu
- Poor contr
- Only one c
- the minimu
- Contact St

This product i
controlling mit
may be neces
not demonst

To help prev

- IPM techni
- Ensure all i
- Do not wa
- Always ap
- Unless oth
- of action.
- Because o
- 4SC, when

Common lace
Insidious flow
Predatory mite
Predatory mite
Predatory mite
Predatory mite

The use of the

Do not conta

PESTICIDE

PESTICIDE

CONTAINER

[For rigid, n
promptly afte
flow begins to
or store rinsa
for recycling.

[For rigid, n
promptly afte
Replace and
container on
rinsate into a
for recycling.

a more dilute mixture is injected.

- Poor control of mites may result if sprinkler application patterns do not sufficiently overlap. Excessive overlap may result in crop injury.
- Only one chemigation application may be made per calendar year, and the system must be operated at 80% to 100% during the application to apply the minimum amount of water possible.
- Contact State lead agencies for pesticide regulation for State-specific requirements pertaining to chemigation.

STRATEGIES FOR RESISTANCE MANAGEMENT

This product is effective against mites while safe for predaceous beneficial insects when used as directed. The chemistry of this product also helps in controlling mites that have developed a resistance to other commonly used products. Due to these properties, fewer product applications for mite control may be necessary; and the likelihood of resistance development is reduced. The active ingredient in BIFENAMITE 4SC is a Group 20D Acaricide that has not demonstrated cross-resistance with other commercial miticides, and is suitable for use in rotation with other miticides.

To help prevent resistance from developing, the following strategies should be followed:

- IPM techniques should be incorporated into your insect control program.
- Ensure all treated foliage is thoroughly covered by spray.
- Do not wait to apply until pest pressure is heavy. Scout regularly and apply BIFENAMITE 4SC as soon as mites are detected.
- Always apply BIFENAMITE 4SC as required by this label.
- Unless otherwise indicated in this label, make only one application of BIFENAMITE 4SC per calendar year and rotate to a product with a different mode of action.
- Because of its selectivity, BIFENAMITE 4SC can be used in conjunction with most biological control organisms available for mite control. BIFENAMITE 4SC, when used as directed, does not adversely affect populations of beneficial/predaceous mites and insects including:

Common lacewing (*Chrysopa carnea*)
 Insidious flower bug (*Orius insidiosus*)
 Predatory mite (*Amblyseius fallacis*)
 Predatory mite (*Phytoseiulus persimilis*)
 Predatory mite (*Typhlodromus pyri*)
 Predatory mite (*Zetzellia mali*)

Seven-spotted lady beetle (*Coccinella septempunctata*)
 Six-spotted thrips (*Scolothrips sexmaculatus*)
 Spider mite destroyer (*Stethorus punctum*)
 Western flower thrips (*Frankliniella occidentalis*)
 Western predatory mite (*Typhlodromus occidentalis*)

The use of these organisms in conjunction with BIFENAMITE 4SC is encouraged as a means of reducing the number of chemical applications.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a dry location.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

[For rigid, non-refillable containers (≤ 5 gallons):] Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake container for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer container for recycling, if available, or puncture and dispose of container in a sanitary landfill, or by incineration.

[For rigid, non-refillable containers (> 5 gallons):] Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer container for recycling, if available, or puncture and dispose of container in a sanitary landfill, or by incineration.

The Directions :
Crop injury, ine
conditions, pre
LLC or Seller. .
relating to such

To the extent al
for the purpose
use conditions.
not reasonably
ALLOWABLE E
POSE NOR AN

To the extent al
from the use or
AND THE EXC
CLAIMS BASE
OR HANDLING
LLC OR SELLE

Albaugh, LLC a
which may not

Conditions of Sale and Limitation of Warranty and Liability

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials, resistant strains or other influencing factors in the use of the product, which are beyond the control of Albaugh, LLC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Albaugh, LLC and Seller harmless for any claims relating to such factors.

To the extent allowed by applicable laws, Albaugh, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Albaugh, LLC and Buyer and User assume the risk of any such use. TO THE EXTENT ALLOWABLE BY APPLICABLE LAW, ALBAUGH, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent allowed by applicable laws, in no event shall Albaugh, LLC or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT ALLOWABLE BY APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ALBAUGH, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ALBAUGH, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

Albaugh, LLC and Seller offer this product, and Buyer and User accept it, subject to foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of Albaugh, LLC.



apply

ps in
ontrol
t has

mode

MITE

lent)
the
tank
inner

lent)
ater.
the
the
inner

ACTIVE INGRI
bifenazate: hyd
[1,1'-biphenyl]-
OTHER INGRE

TOTAL:
Contains 4 pou
EPA Reg. No. 4

KEEP

**IF
SWALLOWED:**

IF INHALED:

Have the produ
tor, or going for
call CHEMTREC

S

GROUP 20D ACARICIDE

Agri Star®

BIFENAMITE 4SC

**Agricultural Miticide
For Agricultural Use Only**

ACTIVE INGREDIENT:	(% by weight)
bifenazate: hydrazine carboxylic acid, 2-(4-methoxy-[1,1'-biphenyl]-3-yl) 1-methylethyl ester	43.2%
OTHER INGREDIENTS:	56.8%
TOTAL:	100.0%

Contains 4 pounds active ingredient bifenazate per gallon.

EPA Reg. No. 42750-321

EPA Est. No. 42750-MO-002

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF INHALED:	<ul style="list-style-type: none"> • Move the person to fresh air. • If the person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For chemical emergency: spill, leak, fire, exposure or accident, call CHEMTREC: 1-800-424-9300.

**See inside booklet for Precautionary Statements
and Directions For Use.**

Manufactured For:

ALBAUGH, LLC
1525 NE 36th Street
Ankeny, Iowa 50021



Albaugh®

Your Alternative™

AD112017

Job #134941

ct.
op
jh,
ns

fit
ial
ns
NT
R-

ng
R,
IG
SE
H,

ty.

PROOF

THIS PROOF IS TO BE CHECKED FOR ACCURACY

Please review and approve **Text, Spelling, Copy Placement, Size, Shape, Colors** and **Dieline**.

Authorized signature accepts responsibility for accuracy of all copy, color break and artwork. Cimarron Label is not liable for any discrepancies subsequently identified.

PLEASE NOTE: Due to color variance between printers/monitors, the colors represented by this proof cannot be deemed accurate. Please refer to a color matching system such as the Pantone Matching System for a truer representation of spot colors.

THIS PROOF IS NOT ACCURATE FOR COLOR-MATCH.





Dieline does not print.

 **Cimarron Label**

Experts in Extended Text Labeling

4201 North Westport Ave. • Sioux Falls, SD 57107

Phone: (605) 978-0451 • Fax: (605) 978-0463

DATE	JOB NUMBER	CUSTOMER
12/11/17	134941	ALBAUGH
LABEL SIZE	LEAFLET FOLDED SIZE	LEAFLET FLAT SIZE
6.75" X 6.75"	6.5" X 6.0"	6.5" X 29.6875"
LABEL COLORS	LEAFLET "OUT" COLORS	LEAFLET "IN" COLORS
 BLK	 BLK  187	 BLK
PATTERN VARNISH: <input checked="" type="checkbox"/> YES <input type="checkbox"/> NO		

Form: CS 006L - 11/8/2011

ARTWORK IS APPROVED

REVISED PROOF NEEDED

**WE CANNOT PROCESS
THIS ORDER WITHOUT AN
AUTHORIZED SIGNATURE**

Signed _____ Date _____