

Visible fast acting for the sults in 12 hours enters the plant on contact

Concentrate 🏗



p Out Of Reach Of C

Contents 32 FL OZ (946 ML.

Kills to the Root!

Makes up to 10.5 gallons



- ✓ Kills unwanted weeds and grasses, roots and all
- ✓ Non-Selective: Kills weeds and grasses
- Use along fences; in cracks of walks, drives and patios; and around flower beds, trees and shrubs
- Use to prepare sites: for large plantings, for planting flower beds, fruit trees, ornamentals, vegetables and gardens, and for renovating lawns

Store and transport in an upright position.
EPA Reg. No. 42750-269-4 EPA Est. No. 4-NY-1
Buyers Guarantee Limited to Label Claims.
@Bonide Products, Inc. All Rights Reserved
Distributed by:
BN
Distributed by:
G301 Sutliff Road
Oriskany, NY 13424

Quart

Quart

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not mix or allow contact with an oxidizing agent; a hazardous chemical reaction may occur.

# **HOW IT WORKS**

This product enters plants through foliage and moves systemically to the roots, killing weeds by stopping the function of a substance found in plants (but not humans or animals). Any of this product not absorbed by plants will be deactivated after it hits the soil. Weeds usually yellow and wilt within hours with complete kill in 1 to 2 weeks.

## WHERE TO USE

On patios, walkways, driveways, gravel or mulch beds. Around flowers (flower beds), shrubs or trees. On driveways & walkways. Along fences. In large areas - for lawn replacement, for garden plot preparation\*

\*IMPORTANT: While this product may be used to prepare a vegetable garden for planting, DO NOT USE for spot weed treatment after planting (any) fruits or vegetables.

## **AMOUNT TO USE**

## For best results:

- Add 3-6 fl. oz. (12 TBS) per gallon of water (use the higher rate for larger weeds)
- Apply on warm days in bright sunlight

# MIXING INSTRUCTIONS

# Tank (Pump-Up) Sprayer:

Use a plastic, fiberglass, plastic-lined steel or stainless steel sprayer. Do not mix, store or apply in (with) a galvanized or unlined steel sprayer (container).

For tough established weeds and for best results, add (measure) 6 fl. oz. (12 Tbs) to (per) (1) gallon of water. Spot treat or spray evenly over 200 sq. ft.

# Hose-End Sprayer:

For large areas consider using the Bonide Auto Mix Sprayer. DO NOT add water. Spray evenly over measured area. After spraying, unused product can be poured back into its original container.

## **HOW TO SPRAY**

- Spray the weeds or grasses you want to kill until thoroughly wet.
- When spot treating weeds around desirable plants, shield plants from drift with a sheet (piece) of cardboard or plastic.
- If desirable plants are accidentally sprayed, rinse off immediately with water (or cut off treated area).

**IMPORTANT:** Do not spray plants or grasses you like - they will die. Not recommended for spot weed control in lawns since this product kills lawn grasses.

## WHEN TO APPLY

Apply when weeds are actively growing.

For best results, apply during warm, sunny weather (above

60° F) (to accelerate systemic movement from foliage to roots). Apply when air is calm to prevent drift to desirable plants.

Rainproof Protection: Rain or watering 30 minutes after application will NOT wash away effectiveness.

Apply when weeds are small and before seeds form. Established (hard to kill) weeds may require a repeat application.

## APPLICATION RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Only the person applying this product may be in the area during application.
- Do not (mix, store or) apply with (in) a galvanized or unlined steel (except stainless steel) sprayer, or apply through any irrigation system.
- DO NOT USE for spot weed treatment after planting fruits or vegetables.
- People and pets must not enter treated areas until spray has dried.

## WHEN TO REPLANT

All ornamental flowers, trees and shrubs may be planted 1 day after application. Lawn grasses, herbs, vegetables (all), and fruits may be planted 3 days after application.

# HOW TO CLEAN SPRAYER

To clean sprayer after use, rinse sprayer and all sprayer parts with water 3 times. Spray rinse water on bare soil or gravel. After cleaning, sprayer may be used for other products.

## ADDITIONAL USES

## To Kill Vines

- If vines are growing up poles, fences, or tree trunks with mature bark, cut vines to a height of 3 to 4 feet and spray vines thoroughly.
   If vines are climbing shrubs or tree trunks with green bark, cut
- If vines are climbing shrubs or tree trunks with green bark, cu vines at base and treat as directed for stumps or spray regrowth. If spraying regrowth, shield shrubs and tree trunks from spray drift with a sheet (piece) of cardboard or plastic.

# To Kill Stumps

- Cut living stump close to ground.
- Drive (drill) 4 to 5 holes into freshly cut stump.
- Immediately pour undiluted product into holes.

## ADDITIONAL USES cont.

IMPORTANT: Some trees may share the same root system. Adjacent trees having a similar age, height and spacing may signal a shared root system. Injury may occur to non-treated trees when one or more trees sharing common roots are treated.

## To Kill Kudzu

 For best results, apply in mid to late summer when vines are mature and actively growing.

# To Kill Wild Blackberry

- Spray anytime plants are actively growing.
   Dead capes should be cut down and removed.
- · Reapplication is often required to kill deep roots.

## To Kill Poison Ivy & Poison Oak

- Contact with these plants anytime of the year can cause an allergic reaction.
- Spray anytime plants are actively growing, but at least 4
  weeks before the first killing frost in the fall. Reapply if
  new growth appears.
- Handle dead plants with rubber gloves.
- Dispose of plants and rubber gloves in tightly sealed garbage bags.

## ADDITIONAL USES cont.

## For Lawn Replacement

Use this product to kill a (an unwanted) lawn and (its) weeds before planting a new lawn (or prior to laying a patio, sidewalk, or driveway).

- Skip one mowing before spraying.
- Use 6 fl. oz. (12 Tbs) per gallon of water for each 300 sq.
- ft. Apply evenly over treatment area. If soil is dry, water before application and 2 to 3 days after application.
- If green patches remain after 7 days, reapply.
- Wait (at least) 3 days for weeds and lawn to die. Repeat if necessary. Rake or till to prepare area for planting seed or sod

- For (flower bed and) Garden (Plot) Preparation

   Use this product to kill weeds, grasses and brush before planting flowers, fruits, vegetables, trees or shrubs.
- If soil is dry, water before application and 2 to 3 days after application.
- Use 6 fl. oz. (12 Tbs) per gallon of water for each 300 sq. ft. Apply evenly over treatment area.
- Wait 1 to 3 days after last application before planting garden. (See when to replant section for details)

# KILLS ALL TYPES OF WEEDS & GRASSES

Weeds (including): Artichoke thistle, beggarweed, black medic. blue medic, blue mustard, blue toadflax, brassbuttons, broadleaf plantain, burclover, buttercup. Canada thistle, cattail, chickweed, cocklebur, common groundsel, common mullein, common plantain, common ragweed, creeping beggarweed, creeping-charlie, curly dock, dandelion, dog fennel, evening primrose, false dandelion, fiddleneck, field bindweed (wild morningglory), field pennycress, filaree, Florida pusley, garden spurge, henbit, horsenettle, horseradish, horseweed/marestail, iceplant, knapweed, knotweed, lambsquarters, lantana, little bittercress. London rocket, mallow, mayweed, milkweed. mouseear chickweed, narrowleaf (buckhorn) plantain, nightshade (silverleaf), oldenlandia, oxalis, Pennsylvania smartweed, pennywort, poison hemlock, prickly lettuce, primrose, prostrate spurge, puncturevine, ragweed, red clover, redroot pigweed, shepherdspurse, smooth catsear. smooth pigweed, sowthistle, spotted spurge, tansy ragwort, tansymustard, teaweed, trumpetcreeper, tumble mustard, Virginia creeper, white clover, whitetop, wild barley, wild mustard, wild oats, wild sweet potato, vellow starthistle. and other broadleaf weeds

Grasses (including):Annual bluegrass, annual ryegrass, bahiagrass, barnyardgrass, bentgrass, Bermudagrass, bromegrass, Cogongrass, crabgrass, creeping bentgrass, dallisgrass, fall panicum, fescue, field sandbur, foxtail, goosegrass, guineagrass, Johnsongrass, Kentucky bluegrass, kikuyugrass, lovegrass, maidencane, nimblewill, nutsedge, orchardgrass, pampas grass, perennial ryegrass, purple nutsedge, quackgrass, sandspur, smooth bromegrass, sprangletop, St. Augustinegrass, tall fesque, Texas panicum, Timothy, torpedograss, vaseygrass, witchgrass, yellow nutgrass, zoysia and other grassy weeds.

Brush (Tough Weeds), Unwanted Trees, Shrubs & Vines (including): Alder, ash, (quaking) aspen, blackberry, bluegum eucalyptus, broom (French, Scotch), buckwheat, ceanothus, chamise, cherry, coyote brush, dewberry, elderberry, elm, eucalyptus, giant reed, hawthorn, hazel, honeysuckle, kudzu, locust, madrone, maple, oak, persimmon, poison oak, poison ivy, poplar, raspberry, sage, sagebrush, salmonberry, saltcedar, sassafras, sumac, sweetgum, tan oak, thimbleberry, tree tobacco, wild rose (multiflora), willow and other tough weeds.

### STORAGE AND DISPOSAL

## **DO NOT** contaminate water, food or feed by storage or disposal. Pesticide Storage: Store in original container only in a safe place away from

direct sunlight Pesticide Disposal: Waste resulting from use of this product may be disposed of on site or at an approved waste facility.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Container is not safe for food, feed or drinking water.

# PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS-CAUTION Causes moderate eve irritation, Avoid contact with eves or clothing, Wash thoroughly with soap and water after handling.

#### ENVIRONMENTAL HAZARDS To protect the environment, do not allow pesticide to enter or run off into

storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. This product is toxic to aquatic invertebrates. Do not apply directly to water. Do

not contaminate water when cleaning equipment or disposing of equipment

### washwaters. USER SAFETY RECOMMENDATIONS Users should:

1. Wash hands before eating, drinking, chewing gum, using tobacco, or using

the toilet 2. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

NOTICE: To the extent consistent with applicable law, Buyer assumes all responsibility for safety and use not in accordance with directions.