

WEED & FEED

Covers up to Squar 5,000 Squar



WINTERIZER

WEED & FEED

Encourages Early Spring Green-Up

Weed Control + Fertilizer

Kills Henbit, Dollarweed, White Clover, Plantain & Chickweed



Covers up to 5,000 Square Feet

14 lbs (6.35 kg)

26-0-12



26-0-12

Covers up to 5,000 Square



WINTERIZER WEED & FEED



Covers up to 5,000 Square Feet 14 lbs (6.35 kg) 26-0-12



PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL MARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY

FOR USE ONLY AS DIRECTED. THE MANUFACTURER NEITHER MAKES NOR LIMITED WARRANTY AND DISCLAIMER

Officials. Information regarding the contents and levels of metals in this product is available on the Internet at 49991-TX-2 (Q), 69759-OR-002 (R), 7401-TX-01 (S) Ebe Est. No, 59144-GA-1 (A), 32802-MO-1 (B), 37351-CO-1(C), 8378-IN-1 (D), 9404-FL-1 (E), 961-PA-01 (F), 961-IL-01(G), 35497-OR-1 (H),

Keep this product off sidewalks, patios, driveways, etc. This fertilizer may stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders, which scatter fertilizer beyond lawn edge and onto concrete.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. PESTICIDE DISPOSAL AND CONTAINER HANDLING: Monrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions. Mever place unused product down recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions. Mever place unused product down recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions. Mever place unused product down

JASOGE AND DISPOSAL				
Сіочет, strawberry	Ноагу сгеss	Plains coreopsis (tickseed)	mədəldtəð to rist	Yarrow, yellow
Clover, red	tidn9H	bineapple weed	рәәмлидс	Yarrow, western
Clover, rabbit foot	dшән	Pigweed, tumble	Spurge, prostrate	Хаггом, соттоп
Clover, hop	Heartleaf drymary	Pigweed, smooth	Spurge, spotted	Wormseed
Сіолет, аіусе	Healall	Pigweed, russian	Aznarama ynig2	Woolly croton
Clover, crimson	Hawkweed, yellow	Pigweed, redroot	Speedwell, thymeleaf	Woodsorrel, yellow (Oxalis)
Cinquefoil	Hawkweed, orange	Pigweed, prostrate	wous 'Jlawbaad'	Woodsorrel, violet
Сһісоту	Намкмеед, тоизееаг	(Peppergrass)	Speedwell, slender	Woodsorrel, creeping
Chickweed, sticky	нэмкмеед, теадом	Pepperweed, virginia	Speedwell, purslane	19loiv bliW
Chickweed, mouseear	Hairy fleabane	Pepperweed, perennial	Speedwell, persian	otstoq təəwz bliW
Сһіскмеед, соттоп	рәәмшпо	Pennycress (frenchweed)	Speedwell, ivyleaf	Wild strawberry
Chamber bitter	وَيonuqsej	Pearlwort	Speedwell, corn	Wild rape
Catsear, spotted (false dandelion)	γείο nud ivy	Parsnip	Speedwell, common	Wild radish
qinteO	roldenrod	Parsley-piert	Speedwell, bi-lobed	Wild parsnip
Carpetweed	Galinsoga, smallflower	Mettle, wood	səlbəən dzinaq2	noino bliW
Carolina geranium (wild)	Viled , seposnile d	Mettle, tall	Sowthistle, spiny	Wild mustard (yellow rocket
Buttonweed, Virginia	Heabane, rough	Mettle, stinging (burning)	Sowthistle, annual	Wild marigold
Buttonweed, common (poorjoe)	Heabane, daisy (annual)	Mustard, wormseed	Sorrel, red (sheep)	Wild lettuce
Buffercup, creeping	Filaree, whitestem	Mustard, white	гиоография	Wild garlic
Buttercup, bulbous	Filaree, redstem	Mustard, tumble	Smooth chaff-flower	Wild four-o-clock
Burweed, lawn (spurweed)	Fiddleneck	Mustard, tansy	Smartweed, pennsylvania	Wild carrot
Burdock "	False sunflower	Mustard, Indian,	Smartweed, pale	Wild buckweat
Burclover	X6l192l64	Wnstard, hedge	gmndysybel, ladysthumb	Wild aster
Bulinettle	Elderberry	Mustard, blue	Zyebyerq z bnrze	Western salsify
рискиоки	Dove weed	Mustard, black	Scarlet pimpernel	Western clematis
Broomweed	Dollarweed (Pennywort)	Mustard, ball	Rongh cinquefoil	Virginia creeper
Rustly oxtongue	Vogtennel (Pennerted)	Mugwort	Redstem filaree	Vetch, wild
Brassbuttons	Dogbane	Μοτηιησσίοτу, woolly	Kagweed, western	Vetch, narrowleat
Blue lettuce	роск, сипу	Могліпддіогу, tall	Ragweed, lancleaf	Vetch, milk
Blood flower milkweed	Dock, broadleaf	Могпіпддіогу, red	кадмеед, соттоп	Уетсћ, ћајгу
Black Medic	Dichondra Dock broadlaaf	Morningglory, Mexican	Kagweed, bur	Vetch, bird
Rlack-eyed Susan	Deadnettle	Могліпддіогу, ічуїеат	Pusley, large flower	Vervain, tall
Bitterweed	Vay flower	Morningglory, bigroot	Pusley, florida	Vervain, prostrate
Bittercress, hairy	Dandelion, false	Moneywort	Pusley, brazil	Vervain, hoary
Bitter Wintercress	Dandelion, common	Mexicanweed	Purslane, common	Vervain, blue
Bindweed, Hedge	Uaisy, oxeye	beewdateM	Puncturevine	Velvetleat
Bindweed, Field	Daisy, English	Marcela	Prostrate knotweed	Irailing crownvetch
Betony, Florida		Mallow, venice	Proetrote (motivoed	I histle, yellow (horrible)
Beggarticks, hairy	pəəmpn)	trewb, wolleM	Prickly lettuce (compass plant)	Thistle, wavyleat
Bedstraw, pineywoods Beggstricks, bairy	dering Brush	Mallow, common	Prairie Suntlower	I histle, scotch
Bedstraw, Northern Bedstraw, pinewwoods	Creeping Jenny (creeping charlie)	Mallow, bristly	Pokeweed Prairie Cunflower	I histle, russian
Bedstraw, Morthern	Creeping Beggarweed	Mallow, alkali Mallow brietly	Poison oak	Thistle, plumeless
Bedstraw, Catchweed Bedstraw, smooth	Corn Spurry Creening Beggarweed	Lespedeza Lespedeza	Yvi nozio9 Veo nozio9	Thistle, platte
	niellum nommo)	Lambsquarters	Plantain, woolly	Thistle, musk
Austrian fieldcress				
Artichoke	Cocklebur, spiny	Kochia	Plantain, slender	Thistle, flodman (prairie)
Aster, white prairie	Cocklebur, oriental	Knotweed, prostrate	Plantain, narrowleaf	Thistle, canada
Aster, purple	Cocklebur	Knawel	Plantain, hoary	Thistle, bull
Aster, white heath	Cockle	təloiv quqmul-ynndol	Plantain, buckhorn	Thistle, blessed
Aster, heath	Clover, white	bəəwnosmil	Plantain, broadleaf	Tanweed
Amaranth, slender	Clover, white sweet	Horseweed	Plantain, bracted	Tansy ragwort

WEEDS CONTROLLED

DROP SPREADERS: Apply 2 strips along each end. Then apply along length of lawn, overlapping slightly and shutting off the spreader before each turn at edge of border strips.

SPREADER SETTINGS HOW TO APPLY - SPREADERS & CALIBRATION

1. Mow lawn to normal height 1 to 2 days before application.

2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).

3. Apply when Broadleaf weeds are young and actively growing, preferably in the morning when dew is on the grass. At the time of application, moisture on the weed leaves from dew, rainfall or irrigation/watering aids in control. If grass is not moist at time of application, sprinkle lightly with water before applying product.

4. Avoid applying when temperatures are consistently above 90°F.

5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

SNOITOURIESTIONS

Possible retreatment may be necessary it lawn is heavily overrun by weeds, if adverse low moisture conditions prevail or if weeds are Apply when Broadleaf weeds are actively growing in Spring or early Fall. Apply when the air is calm to ensure uniform coverage and avoid spreading granules onto flowers, vegetables and ornamental

The maximum broadcast application rate is 4 lb of product per 1000 sq.ft. per application. The maximum number of broadcast applications is 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8 lb of product per For use on Kentucky Bluegrass, Fescue spp., Perennial Ryegrass, common Bermudagrass, Bahiagrass, Zoysiagrass, St. Augustinegrass, Centipedegrass, hybrid Bermudagrass, and Colonial Bentgrass. Apply 2.8 pounds of product per 1,000 square feet of turigrass. One 14 lb bag covers 5000 sq. ft. • Do not apply this product to newly seeded grasses until well-established or after the second or third mowing. • Reseeding or overseeding areas may occur three (3) weeks after the application of this product **USE PRECAUTIONS:**

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish
pronds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to
your lawn, and sweep any product landing on the dirveway, sidewalk, gutter, or street back onto the treated area. To prevent
product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that
day.

Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees, shrubs, exceed specified dosage for any area and do not apply this product within the dripline of, or over surface roots of trees, shrubs, Do not use on dichondra, carpetgrass, Floratam' St. Augustinegrass, creeping bentgrass, or turfgrass where desirable clovers are Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research

This product is a combination of three proven weed killers--2,4-D, mecoprop-p, and dicamba. This herbicide controls a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, and spurge in addition, a 26-0-12 fertilizer provides a It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm crains, drainaged dictines, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not she treated area of the lawn or garden will help to

nformation. FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300. pm PST. You may also contact the National Poison Control Center, 1-800-222-1222, day or night, for emergency medical treatment nformation on this product, contact the National Pesticide Information Center, 1-800-858-7378, Monday - Friday, 8:00 am to 12:00 Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For If in Eyes

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. **Hazards to Humans and Domestic Animals** PRECAUTIONARY STATEMENTS

STOP READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.





MEED & LEED **WINTERIZER**

