

Rainfast in 3 hours See results in hours

Can re-seed turf I day after applying

One gallon of concentrate will treat up to 10000 ft<sup>2</sup>

Controls Weeds, Disease, Moss, Algae, and Lichens in turf

People and pets can enter treated area when spray dries

# NOT FOR USE IN CALIFORNIA

EPA Reg. No. 67702-26 EPA Est. No. 48498-CA-I

**KEEP OUT OF REACH OF CHILDREN** 

**NET CONTENTS: 2.5 GALLON** 

CAUTION

Active Ingredient	By weight	
Iron HEDTA (FeHEDTA)	26.52%	
Other Ingredients	73.48%	
Total	100.00%	



SEE INSIDE BOOKLET FOR DIRECTIONS FOR USE AND ADDITIONAL PRECAUTIONARY STATEMENTS

IF IN	-Hold eye open and rinse slowly and gently with water for 15-20 minutes.
EYES	-Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
	-Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. During other times, call the poison control center at 1-800-222-1222.

# PRECAUTIONARY STATEMENTS

FIRST AID

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS. CAUTION:** Causes slight eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

### **ENVIRONMENTAL HAZARDS**

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

# **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read and follow all applicable directions and precautions on this label before using.

SITES: Lawns, turf (use on rights of way or non-crop areas), golf courses, parks, playgrounds, cemeteries, athletic fields

# **PESTS:**

Controls or suppresses weeds such as but not limited to: Black medic (Medicago lupulina), Broadleaf plantain (Plantago major), Bull thistle (Cirsium vulgare), Canada thistle (Cirsium arvense), Common chickweed (Stellaria media), Creeping buttercup (Ranunculus repens), Creeping Charlie/Ground ivy (Glechoma hederacea), Dandelion (Taraxicum officinale), Dovefoot geranium (Geranium molle), English daisy (Bellis perennis), False dandelion (Hypochoeris radicata), Healall (Prunella vulgaris), Lawn burweed (Soliva ptero-sperma), Moss, liverworts, algae and lichens,

**PESTS (Cont.):** Narrow-leaved plantain (*Plantago lanceolata*), Oxalis (*Oxalis spp.*), Persian Speedwell (*Veronica persica*), Prostrate knotweed (*Polygonum aviculare*), Shepherd's-purse (*Capsella bursa-pastoris*), Silverweed cinquefoil (*Potentilla anserina*), Slender speedwell (*Veronica filiformis*), White clover (*Trifolium repens*), Wild chamomile (*Matricaria chamomilla*), Wild geranium (*Geranium robertianum*)

Controls or suppresses diseases such as but not limited to: Dollar Spot, Rust, Snow Mold

# **Application Directions:**

Rate of Spray Solution	Rate of Concentrate	Weeds	Diseases
2.8 fl. oz./ yd² or 2.5 ga/1000 ft²	12.6 fl. oz/ 1000 ft <sup>2</sup>	Rate for easy-to-control weeds: black medic (Medicago lupulina), Persian speedwell (Veronica persica), slender speedwell (Veronica filiformis), wild geranium (Geranium robertianum), and moss. This rate can also be used for weed seedlings of: false dandelion (Hypochaeris radicata), English daisy (Bellis perennis), dandelion (Taraxacum officinale) or white clover (Trifolium repens).	Rate for control or suppression of diseases such as dollar spot, snow mold, and rust. Use the higher rate when disease pressure is high.
5.7 fl. oz./ yd² or 4.9 ga/1000 ft²	25.2 fl. oz/ 1000 ft <sup>2</sup>	Standard rate for control of most weed species: dandelion ( <i>Taraxacum officinale</i> ), English daisy ( <i>Bellis perennis</i> ), false dandelion ( <i>Hypochaeris radicata</i> ), Creeping Charlie/ Ground ivy ( <i>Glechoma hederacea</i> ), Prostrate knotweed ( <i>Polygonum aviculare</i> ), white clover ( <i>Trifolium repens</i> ), bull thistle ( <i>Cirsium vulgare</i> ), Canada thistle ( <i>Cirsium arvense</i> ), common chickweed ( <i>Stellaria media</i> ), narrow-leaved plantain ( <i>Plantago lanceolata</i> ), dove's foot geranium ( <i>Geranium molle</i> ), oxalis ( <i>Oxalis spp.</i> ), lawn burweed ( <i>Soliva ptero-sperma</i> ), and algae, and for the SUPPRESSION of broad-leaved plantain ( <i>Plantago major</i> )	
11.3 fl oz./ yd² or 9.8 ga/1000 ft²	50 fl. oz./ 1000 ft²	Rate for tough perennial weeds such as: creeping buttercup (Ranunculus repens) and broad-leaved plantain (Plantago major)	

Shake well before using. Mix one part FIESTA TURF WEED KILLER with 24 parts water (5 oz. in 1 gallon of water). Apply the mixed solution at a rate of 2.5-10 gallons/1000 ft $^2$  (2.8-11.5 fluid oz/yard $^2$ ), using the chart to determine how much diluted spray to use per area. Any standard hand-held or backpack sprayer can be used. Thoroughly spray weeds. Use a coarse nozzle setting to reduce drift. Uniform coverage is important. Repeat treatment in 3 to 4 weeks for best results.

**For Spot Treatment:** Shake well before using. Mix one part FIESTA TURF WEED KILLER with 24 parts water (5 oz. in 1 gallon of water). Thoroughly spray weeds to point of runoff using any standard hand-held or backpack sprayer. Use a coarse nozzle setting to reduce drift. Repeat treatment in 3 to 4 weeks for best results.

What will I see? Works quickly; dying weeds will be seen within hours of application. Treated weeds, algae and moss will turn brown or black. Occasionally a darkening of the grass leaf blades can occur after treatment; however the grass will recover within a few days to weeks. May stain thatch (dead grass under turf); this does not affect the health of the grass.

Use Precautions: Do not apply to drought stressed grass; ensure turf is well watered prior to application. Do not apply when the daytime temperature will exceed 85°F. Do not apply to bentgrass. If unsure of grass sensitivity, test spray a small area of the turf. Do not apply to newly seeded areas prior to grass seed germination. Avoid spraying desirable plants. Any desirable plants or structures accidentally contacted should be rinsed with water. To avoid staining, wait until treated area dries before re-entry. Do not apply if rainfall is expected within 3 hours of application. Do not irrigate within 3 hours of application. Do not apply to the same area more than 4 times a year.

### **NOTICE TO BUYER**

Seller warrants that this product conforms to the chemical description on this label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions. To the extent consistent with applicable law, this warranty does not extend to use of this product contrary to label directions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller. To the extent consistent with applicable law, the Seller shall not be liable for any such use. To the extent consistent with applicable law, the Seller makes no other warranties, either expressed or implied.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a secure place, away from open fire or flame. Keep container closed and reseal after use. If spilled, use absorbent materials and dispose of in an approved manner. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available.

### PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS. CAUTION: Causes slight eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

### FIRST AID

IF IN **EYES** 

-Hold eye open and rinse slowly and gently with water for 15-20 minutes. -Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. -Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. During other times, call the poison control center at 1-800-222-1222.

## Registrant:

# W. Neudorff GmbH KG

An der Mühle 3 · 31860 Emmerthal · Germany Phone: 250-652-5888 · www.neudorffpro.com

Patent #6,972,273

EPA Reg. No. 67702-26 EPA Est. No. 48498-CA-I

**NET CONTENTS: 2.5 GALLON** 

SEE ATTACHED BOOKLET FOR **DIRECTIONS FOR USE** 



**Active Ingredient:** 

Iron HEDTA (FeHEDTA) Other Ingredients: Total:

26.52% 73.48%

**KEEP OUT OF REACH OF CHILDREN** 

CAUTION