### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool, dry place in original container.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: (Paper Bag or Plastic Bag)

Nonrefillable container. Do not reuse or refill this container.

Completely empty bag into application equipment by shaking and tapping sides and bottom to loosen clinging particles. When completely empty, offer for recycling if available, or dispose of bag in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

91411-12-70051\_Kocide HCu\_20180329\_48\_70051\_.pdf



### **ACTIVE INGREDIENT:**

Other Ingredients: 23.0% Total:

\*Metallic Copper (Cu<sup>2+</sup>) Equivalent. 50.1% by weight † CAS No. 20427-59-2

You may also contact CHEMTEL (800) 255-3924 (24 hours) for emergency medical treatment information

See side/back panels for additional precautionary statements

## **KEEP OUT OF REACH OF CHILDREN DANGER - PELIGRO**

**FIRST AID** IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. IF SWALLOWED: Call poison control center or doctor for treatment advice Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor • Do not give anything by mouth to an unconscious person. • Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice **IF INHALED:** • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call poison control center or doctor for treatment advice. **HOT LINE NUMBER** Have the product container or label with you when calling a poison control center or doctor or going for treatment.

EPA Reg. No. 91411-12-70051 EPA Est. No. 91411-TX-1

Manufactured for: Certis USA LLC 9145 Guilford Road, Suite 175 Columbia, MD 21046

Net: 20 lb. (9.07 kg) Nonrefillable Container

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER-PELIGRO

Corrosive. Causes irreversible eye damage. Do not get in eyes, skin, or clothing. Harmful if swallowed or absorbed through skin. Harmful if inhaled. Avoid contact with skin. Avoid breathing

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

- Mixers, loaders, applicators, and other handlers must wear:
- Chemical resistant gloves made of any waterproof material

See engineering controls for additional requirements. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with the product's concentrate. Do not

### **ENGINEERING CONTROLS**

When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR Part 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIROMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may contaminate water through runoff. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Kocide LLC, 9145 Guilford Road, Suite 175, Columbia MD, 21046

ESL 20170621

eated by: 09-14-17 jw : 10-24-17 jw ze: 12" (w) x 24" (h) d size: 3" (w) x 4" (h)

Kocide HCu Active Ingredient: Copper Hydroxide\* Other Ingredients: \*Metallic Copper (Cu2+) Equivalent, 50.1% by weight †CAS No. 20427-59-2

#### KEEP OUT OF REACH OF CHILDREN **DANGER - PELIGRO**

EPA Est. No. 91411-TX-1

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted detalle. (If you do not understand the label, find someone to explain it to you in detail.)

EPA Reg. No. 91411-12-70051 Nonrefillable Container

Net: 20 lb. (9.07 kg)

A20170209

# Kocide **CERTIS**

	FIRST AID						
If In Eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes.     Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.     Call a poison control center or doctor for further treatment advice.						
If Swallowed	Call poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.						
If On Skin Or Clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.						
If Inhaled	Move person to fresh air.     If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.     Call noise pontrol center or doctor for treatment advice.						

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact CHEMTEL (800) 255-3924 (24 hours) for emergency medical treatment information Note to Physician: Probable mucosal damage may contraindicate the use of gastric

HOT LINE NUMBER

See side/back panels for additional precautionary statements PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS **DANGER-PELIGRO**Corrosive. Causes irreversible eye damage. Do not get in eyes, skin, or clothing. Harmful

is wallowed or absorbed through skin. Harmful if inhaled. Avoid contact with skin. Avoid breathing vapor or spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

- Long-sleeved shirt
- Long pants
   Shoes and socks
   Chemical resistant gloves made of any waterproof material Protective eyewear

See engineering controls for additional requirements. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been denoted or heavily contaminated with the products expended. contaminated with the product's concentrate. Do not reuse them. ENGINEERING CONTROLS:

When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR Part 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as pecified in the WPS.

#### USER SAFETY RECOMMENDATIONS

Users Should:
• Wash hands before eating, drinking, chewing gum, using tobacco, or using the wash hands before eating, difficility, chewing guilt, using lobacco, or using the toilet.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly

and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean

#### **ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and aquatic invertebrates and may contaminate water through runoff. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For requirements specific to your State or Tribe, consult the State or Tribe agency responsible for pesticide regulations.

### AGRICULTURAL USE REQUIREMENTS

AGRICULTURAL USE REQUIREMENTS
Use this product only in accordance with its labeling and with the Worker Protection
Standard, 40 CFR Part 170. This Standard contains requirements for the protection of
agricultural workers on farms, forests, nurseries, greenhouses and handlers of
agricultural insecticides. It contains requirements for training, decontamination,
notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry

- PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil or water is:
- Chemical resistant gloves made of any waterproof material
   Shoes plus socks
   Protective eyewear

interval (REI) of 48 hours without required PPE.

\*Not registered for use in California

(continued)

### AGRICULTURAL USE REQUIREMENTS (continued)

For Greenhouse Uses ONLY:
The 48 hour restricted entry interval (REI) may be reduced to 24 hours, provided that the following conditions are met:

For at least seven days following the application of copper-containing products in

- At least one container or station designed specifically for flushing eyes is available in operating condition with the WPS-required decontamination supplies for

workers entering the area treated with copper-containing products
- Workers are informed orally, in a manner they can understand:

• that residues in the treated area may be highly irritating to their eyes

• that they should take precautions, such as refraining from rubbing their eyes, to keep the residues out of their eyes

• that if they do get residues in their eyes, they should immediately flush their eyes with the overline orange of the following the containing or use flush etailing that is legated with the

eyes with the eye flush container or eye flush station that is located with the how to operate the eye flush container or eye flush station

#### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides 40 CFR part 170. The WPS applies when this product is used to produce agricultural plants on farms, forests,

Do not enter or allow others to enter until sprays have dried. PRODUCT INSTRUCTIONS KOCIDE® HCu may be applied as an aerial, ground dilute or ground concentrate spray unless specifically directed otherwise in the specific crop use directions.

The per acre use rate of KOCIDE® HCu is applicable for both dilute and concentrate The per acre use rate of KOCIDLE\* HCu is applicable for both dilute and concentrate spraying. Depending upon the equipment used and the specific rop, the spray volume applied per acre will differ. Refer to Minimum Spray Volume Table. Complete spray coverage is essential to assure optimum performance from KOCIDLE\* HCu. When treating by aerial application or with low volume application equipment, unless you have had specific previous experience, it is advisable to test for compatibility and tolerance to crop injury prior to full scale commercial utilization.

Consult the KOCIDE® HCu label for specific rates and timing of application by crop. Where application rates and intervals are provided in a range (e.g.,  $\frac{4}{10}$  to 12 pounds and 7 to 10 days), use the higher rates and shorter spray intervals when rainfall is heavy and/or disease pressure is high. Use the higher rates for large mature tree crops.

The Pre-Harvest Interval (PHI) for KOCIDE® HCu is 0-days unless noted.

#### **PRECAUTIONS** $\bullet$ If KOCIDE $^{\rm @}$ HCu is applied in a spray solution having a pH of less than 6.5, phytotoxicity

may occur.

Environmental conditions such as extended periods of wet weather, acid rain, etc. which alter the pH of the leaf surface may affect the performance of KOCIDE® HCu resulting in possible phytotoxicity or loss of effectiveness.

in possible pryloxicity or loss of effectiveness.

Agricultural chemicals may perform in an unpredictable manner when tank mixed, especially where several products are involved. Reduced effect on pests or crop injury may occur. Unless recommended on this label or by a statel/local expert, it is advisable to test for compatibility and potential crop injury prior to commercial use of a new tank mix.

While volume is important in obtaining full spray coverage, often factors such as foliage death to include the such as the control of the contro density, environmental conditions and sprayer calibration have a greater impact. Always be sure that sprayers are calibrated to spray equipment manufacturer's specifications and environmental conditions are within those recommended by State and local

and environmental conditions are within those recommended by State and local regulatory authorities.

When mixing, fill the spray tank one-half full with water. Add KOCIDE® HCu slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. Spreaders, stickers, insecticides, nutrients, etc. should be added last. If compatibility is in question, use the Compatibility Jar Test before mixing a whole tank or contact your chemical supplier. Observe all precautions and limitations on the labels of all products used in mixtures.

PRODUCT RESTRICTIONS

• Do not tank mix KOCIDE® HCu with any product containing aluminum tris (O-ethyl phosphonate) fungicide for use on any registered crops unless appropriate precautions have been taken to buffer the spray solution because severe phytotoxicity may result. Use in accordance with the most restrictive of label limitations and precautions. Do not exceed label dosage rates. This product cannot be mixed with any product containing a label prohibition against such mixing.

a label prohibition against such mixing.

This product may be reactive on masonry and metal surfaces such as galvanized roofing. Inis product may be reactive on masonry and metal surfaces such as galvanzed rooting. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.
 It must be determined if proper application equipment is available and if waste associated with its use can be properly handled. Agricultural chemicals are often reactive with the materials used in the construction of application equipment, such as aluminum, rubber and some synthetic materials. This factor should be taken into auminum, rubber and some synthetic materials. Inits factor should be taken into consideration when selecting proper application equipment. It is necessary that all application equipment be thoroughly flushed with clean water after each day's use. Apply this product only through one or more of the following types of systems: sprinkler, including center pivot, lateral move, traveler, big gun, or plastic pipe solid set system(s). Do not apply this product through any other type of irrigation system. In California, do not apply in systems which contain aluminum parts or components.

CROP CLASSIFICATION CITRUS: Grapefruit, Kumquat, Lemon, Lime, Orange, Pummelo, Tangelo and Tangerine. CONIFERS: Douglas Fir, Fir, Juniper, Leyland Cypress, Pine and Spruce. FIELD CROPS: Alfalfa, Barley, Corn, Oats, Peanut, Potato, Soybean\*, Sugar Beet and

SMALL FRUITS: Blackberry, Blueberry, Cranberry, Currant, Gooseberry, Raspberry and

TREE CROPS: Almond, Apple, Apricot, Avocado, Banana/Plantain, Cacao, Cherry, Coffee, Filbert, Mango, Nectarine, Olive, Peach, Pear, Pecan, Pistachio, Plum, Prune, Quince and Walnut.

VEGETABLES: Bean, Beet, Beet Greens, Broccoli, Brussels Sprout, Cabbage, Chinese Cabbage, Cantaloupe, Carrot, Cauliflower, Celeriac, Celery, Cucumber, Eggplant, Greens (Collard, Mustard and Turnip), Honeydew, Kale, Kohlrabi, Lettuce, Muskmelon, Okra, Onion/Garlic/Leek, Pea, Pepper, Pumpkin, Spinach, Squash, Tomato, Watercress and

MISCELLANEOUS: Atemoya, Carambola, Chives, Dill, Ginseng, Guava, Litchi, Live Oak\*, Macadamia, Mamey Sapote, Papaya, Parsley, Passion Fruit, Sugar Apple and Sycamore. GREENHOUSE AND SHADEHOUSE CROPS: KOCIDE® HCu may be used in GREENHOUSE AND SHADEHOUSE CROPS: ROCIDE® HCU may be used in greenhouses and shadehouses to control diseases on any crop on this label where physiology allows greenhouse or shadehouse culture. While specific directions are presented for Citrus, Cucumber, Eggplant, Pepper and Tomato; general use may occur for any crop on this label where physiology allows greenhouse or shadehouse culture. Consequently, injuries arising from the use of KOCIDE® HCu on these types of greenhouse and shadehouse crops are the responsibility of the user.

### TURF (non-residential)

\*Not registered for use in California.

Minimum Recommended Spray Volume (Gallons Per Acre) When Applying  $\mathsf{KOCIDE}^\otimes\mathsf{HCu}$ 

	Aerial	Dilute	Concentrate
Citrus	10	800	100**
Conifers	10	100	30
Field Crops	3	20	3
Ornamentals	10	100	50
Small Fruits	5	150	50
Tree Crops	10	400	50
Vegetables	3	20	3
Vines	5	150	50
Miscellaneous	10	150	50

\*Pesticide application equipment such as "Curtec" or other similar sprayers which are capable of obtaining thorough coverage at low volumes may be used at as low as 20 gallons per acre of spray volume.

The following specific instructions are based on general applications. The recommendations of State Agricultural Extension Services should be closely followed as to timing, frequency and number of sprays per year.

#### FROST INJURY PROTECTION **BACTERIAL ICE NUCLEATION INHIBITOR**

Application of KOCIDE® HCu made to all crops listed on this label at rates and stages of growth indicated on this label, at least 24 hours prior to anticipated frost conditions, will afford control of ice nucleating bacteria (Pseudomonas syringae, Enwinia herbicola, and Pseudomonas fluorescens) and may therefore provide some protection against light frost. Do not use KOCIDE® HCu for those geographical areas where weather conditions favor

CITRUS: Grapefruit, Kumquat, Lemon, Lime, Orange, Pummelo, Tangelo and Tangerine

KOCIDE® HCu may be mixed with dry foliar nutritionals (micronutrients) to create "Shot Bag" mixes to meet the various nutritional requirements of citrus and provide disease protection as described on this label. KOCIDE® HCu per acre rates in these mixes must not exceed the maximum labeled rates for disease control. Adding foliar nutritionals or other products to spray mixtures containing KOCIDE® HCu and applying to citrus during the post bloom period when young fruit are present may resu

Disease	Application Rate/Acre	Maximum Annual Rate/Acre	Use Instructions			
Algal Spot, Melanose, Scab	3 – 6.3 lb.	25.1 lb.	Apply as pre-bloom and post-bloom sprays. Use the higher rates when conditions favor disease.  Minimum retreatment interval is 7 days.			
Greasy Spot, Pink Pitting	3 – 6.3 lb.	25.1 lb.	Apply in summer on expanded new flush. Repeat on subsequent flushes where disease pressure is severe. Use the higher rates when conditions favor disease.  Minimum retreatment interval is 7 days.			
Alternaria Brown Spot	4 – 6.3 lb.	25.1 lb.	On susceptible varieties apply when the first spring flush appears and each flush thereafter. Application to fruit shoul after two thirds of the petals have fallen and be repeated on a 7 to 21 day schedule if needed. Use the higher rates conditions favor disease.  Minimum retreatment interval is 7 days.			
Phytophthora Brown Rot, Septoria Spot	4 – 6.3 lb.	25.1 lb.	Begin application in fall before or just after the first rain and continue if needed. For Brown Rot only, apply to skirts of trees to a height of at least 4 feet. For control of Septoria Spot or where fruit have already been infected with Brown Rot, apply to entire tree. Apply also to bare ground one foot beyond skirt.  Use the higher rates when conditions favor disease.  Minimum retreatment interval is 7 days.  NOTE: In California, in areas subject to cooper injury, add 0.25 to 0.5 pound of high quality lime per pound of KOCIDE® HCU.			
Phytophthora Foot Rot	1 lb.	25.1 lb.	Mix at a 1 pound to 0.5 to 1 gallon of water ratio, "Tre-Hold" or latex paint. Paint trunks of trees from the soil surface to the lowest scaffold limbs. Apply in May prior to summer rains and/or in the fall prior to wrapping trees for freeze protection. Treatment serves as protection for up to 1 year, but does not cure existing infections.  NOTE: Areas where microjet or low volume irrigation hit the tree trunk may require retreatment due to wash off.			
Citrus Canker (suppression)	6.3 lb.	25.1 lb.	Spray flushes 7 to 14 days after shoots begin to grow. Young fruit may require an additional application. Number and timing of applications will be dependent upon disease pressure. Under heavy pressure, spray each flush of new growth.  Minimum retreatment interval is 7 days.			
Black Spot*	3.5 - 6.3 lb.	25.1 lb.	Begin treatment prior to or when disease first appears and repeat every 7 to 21 days if needed. Use the higher rates and shorter spray intervals when conditions favor disease.  Minimum retreatment interval is 7 days.			

NOTE: Phytotoxicity may occur on young tender flush when KOCIDE® HCu is applied to citrus seedlings grown in greenhouses or shadehouses. \*Not registered for use in California.

Field Nursery Grown

To control Melanose, Scab, Pink Pitting, Greasy Spot, Brown Rot and for suppression of Citrus Canker, apply 4 to 6.3 pounds of KQCIDE® HCu per acre. Apply KQCIDE® HCu a 28 day intervals if needed depending on disease severity. Minimum retreatment interval is 7 days. Maximum Annual Rate/Acre is 25.1 lbs.

	FIELD CROPS							
Crop	Disease	Application Rate/Acre	Maximum Annual Rate/Acre	Use Instructions				
Alfalfa	Cercospora Leaf Spot, Leptosphaerulina Leaf Spot	1 lb.	2.24 lb.	Apply 10 to 14 days before each harvest or earlier if disease threatens.  Minimum retreatment interval is 30 days.  NOTE: Spray injury may occur with sensitive varieties such as Lahontan.				
Corn (Field Corn, Popcorn, Seed Corn, Sweet Corn)	Bacterial Stalk Rot	1 – 2 lb.	8.4 lb. Begin treatment when disease first appears and repeat every 7 to 10 days if needed. Use the frates and shorter spray intervals when conditions favor disease.  Minimum retreatment interval is 7 days.					
Peanut	Cercospora Leaf Spot	1.5 lb.	9.46 lb.	Begin spraying at 35 to 40 days after planting or when disease symptoms first appear and repeat at 7 to 14 day intervals if needed. Reduce sprays to 7 day intervals during humid weather. Flowable sulfur may be added.  Minimum retreatment interval is 7 days.				
Potato	Early Blight, Late Blight	1 – 4 lb.	50 lb.	Apply 1 to 2 pounds at 5 to 10 day intervals if needed starting when plants are 2 to 6 inches high in locations where disease is light. Apply up to 4 pounds per acre when disease is more severe. Under conditions of severe disease, control with KOCIDE® HCu will be improved by tank mixing with other compatible fungicides registered for use on potatoes. Read and follow all label instructions of tank mix partners.  Minimum retreatment interval is 5 days.				
Soybean*	Bacterial Blight, Downy Mildew	1 – 1.5 lb.	9.48 lb.	For protective sprays, make first application when plants are 6 inches high; repeat on a 7 to 14 day schedule if needed depending on environmental conditions. Use the higher rates for more severe disease.  Minimum retreatment interval is 7 days.				
Sugar Beet	Cercospora Leaf Spot	2 – 2.5 lb.	15.7 lb.	Begin applications when conditions first favor disease development and repeat at 10 to 14 day intervals if needed. Use the higher rates when conditions favor disease.  Minimum retreatment interval is 10 days.				
Wheat, Barley, Oats	Fusarium Head Blight Suppression*, Helminthosporium Spot Blotch, Powdery Mildew Suppression	1 lb.	2.12 lb.	Make applications for early season disease control through heading. Use higher rates when conditions favor disease.  Add an adjuvant.  Minimum retreatment interval is 10 days.				

	SMALL FRUITS								
Crop	Disease	Application Rate/Acre	Maximum Annual Rate/Acre	Use Instructions					
Blackberry (Aurora, Boysen, Cascade, Chehalem, Logan, Marion, Santiam, Thornless Evergreen)	Anthracnose, Cane Spot, Leaf Spot, Pseudomonas Blight, Purple Blotch, Yellow Rust	4 lb.	20 lb.	Make fall application after harvest. Apply delayed dormant spray after pruning/training in the spring. If needed, agricultural-type spray oil may be added.					
	Anthracnose, Cane Spot, Leaf Spot, Purple Blotch, Yellow Rust	2 lb.	20 lb.	Apply when leaf buds begin to open and repeat when flower buds show white. Repeat on a 7 day interval if needed. If needed, agricultural-type spray oil may be added.  NOTE: Crop injury may occur if applied to foliage under certain environmental conditions such as hot or prolonged moist periods. Discontinue applications if signs of crop injury appear.  Minimum retreatment interval is 7 days.					
Blueberry	Bacterial Canker	3 – 4 lb.	16.8 lb.	Make first application before fall rains and a second application 4 weeks later. Use the higher rates when conditions favor disease.					
	Fruit Rot, Phomopsis Twig Blight	3 – 4.2 lb.	16.8 lb.	Dormant Application: Begin applications when bloom buds begin to swell. Make additional applications at 7 to 14 day intervals if needed before blooms open.  Minimum retreatment interval is 7 days.					

SMALL FRUITS (continued)									
Crop	Disease	Application Rate/Acre	Maximum Annual Rate/Acre	Use Instructions					
Cranberry	Fruit Rot	4.2 lb.	25.1 lb.	Make first application in late bloom. Apply one or two additional applications at 7 to 14 day intervals if needed depending on disease severity.  Minimum retreatment interval is 7 days.					
	Rose Bloom	4.2 lb.	25.1 lb.	Apply three sprays on 7 to 14 day schedule if needed as soon as symptoms are observed. Minimum retreatment interval is 7 days.					
	Bacterial Stem Canker	4.2 lb.	25.1 lb.	Apply post-harvest and again in spring at bud swell. Apply one or two additional applications at 7 to 14 day intervals if needed depending on disease severity. Minimum retreatment interval is 7 days.					
	Leaf Blight, Red Leaf Spot, Stem Blight, Tip Blight (Monilinia)	4.2 lb.	25.1 lb.	Apply delayed dormant spray in the spring. Repeat at 7 to 14 day intervals if needed through pre-bloom.  Minimum retreatment interval is 7 days.					
Currant, Gooseberry	Anthracnose, Leaf Spot	5 lb.	20 lb.	Make initial application after first leaves have expanded. Continue on a 10 to 14 day schedule if needed during wet conditions in the spring. Make an additional application after harvest. Minimum retreatment interval is 10 days.					
Raspberry	Anthracnose, Cane Spot, Leaf Spot, Pseudomonas Blight, Purple Blotch, Yellow Rust	4 lb.	20 lb.	Make fall application after harvest. Apply delayed dormant spray after training in the spring. If needed, agricultural-type spray oil may be added.					
	Anthracnose, Cane Spot, Leaf Spot, Purple	2 lb.	20 lb.	Apply when leaf buds begin to open and repeat when flower buds show white. Repeat on a 7 day interval if needed. If needed, agricultural-type spray oil may be added.					
	Blotch, Yellow Rust			<b>NOTE:</b> Crop injury may occur if applied to foliage under certain environmental conditions such as hot or prolonged moist periods. Discontinue applications if signs of crop injury appear. Minimum retreatment interval is 7 days.					
Strawberry	Angular Leaf Spot (Xanthomonas), Leaf Blight, Leaf Scorch, Leaf Spot	2 - 3 lb.	16.3 lb.	Begin application when plants are established and continue on a weekly schedule throughout the season. Apply in at least 20 gallons of water. Use the higher rates when conditions favor disease. NOTE: Discontinue applications if signs of crop injury appear. Minimum retreatment interval is 7 days.					

	Leaf Spot			Minimum retreatment interval is 7 days.
Crop	Disease	Application Rate/Acre	Maximum Annual Rate/Acre	TREE CROPS  Use Instructions
Almond only	Bacterial Blast	1 – 3 lb.	35.9 lb.	For bacterial blast control in sprinkler irrigated orchards or where disease is severe, apply 1.0 poun per acre post-bloom at 2 week intervals if needed or just before sprinkling.
Almond only	Bacterial Spot (Xanthomonas	8 – 16 lb. 0.5 – 2 lb.	35.9 lb. 35.9 lb.	Dormant: Make first application at late dormant. Use the higher rates when conditions favor disease Pink through full bloom: Maximum use rate is 2.0 pounds.
	arboricola pv. Pruni)			Petal Fall: Maximum use rate is 1.0 pounds.  Post Petal Fall: Maximum use rate is 0.5 pounds.  Time sprays around rain events and temperature. Make a minimum of one application to prevenew infections.  Note: Copper applied after bloom can be potentially phytotoxic. Leaf spotting and premature lefall can occur if rates are extended.
Almond, Apricot,	Bacterial Blast	8 – 16 lb.	35.9 lb.	Minimum retreatment interval is 5 days.  Make first application before fall rains and a second at late dormant. Use the higher rates wh
Cherry, Plum, Prune	(Pseudomonas), Bacterial Canker, Coryneum Blight (Shot Hole)			conditions favor disease. If needed, agricultural-type spray oil may be added. Minimum retreatment interval is 7 days. For Cherries: Where disease is severe, an additional application shortly after harvest may be require NOTE: Foliar injury may occur from post-bloom sprays on almonds, especially on NePlus varieties
	Blossom Brown Rot, Coryneum Blight (Shot Hole)	3 lb. (Almond) (All Others) 2 – 3 lb.	35.9 lb.	Apply during early bloom. Do not apply after full bloom or injury may occur. Use the higher rat when rainfall is heavy and disease pressure is high.  Minimum retreatment interval is 5 days.
	Black Knot (Plum)	2 – 3 lb.	35.9 lb.	Make an application at bud swell up to early bloom for early season disease suppression. App before full bloom. Minimum retreatment interval is 5 days. Use the higher rates when rainfall is hea and disease pressure is high. Minimum retreatment interval is 5 days.  NOTE: To avoid plant injury, do not use after full bloom.
	Cherry Leaf Spot (Sour Cherries Only)	3 lb.	35.9 lb.	Apply at petal fall as well as 1 to 2 times after petal fall.  Do not apply to sweet cherry or the English Morello variety as severe injury will result. The addit of 1 to 3 pounds of hydrated lime per pound of KOCIDE® HCu may reduce crop injury. Minimum retreatment interval is 5 days.  NOTE: Moderate to severe injury such as leaf spotting and defoliation may occur from post-blo applications.
Apple	Anthracnose, Blossom Blast, European Canker (Nectria), Shoot Blast (Pseudomonas)	12 – 16 lb.	31.9 lb.	Apply before fall rains. Use the higher rates when conditions favor disease. <b>NOTE:</b> Use on yellow varieties may cause discoloration. To avoid discoloration, pick before sprayi Only one dormant application allowed per year.
	Apple Scab, Fire Blight	4 – 12 lb.	31.9 lb.	Make application between silver-tip and green-tip. Apply as a full cover spray for early season diseasuppression.  NOTE: Moderate to severe crop injury may occur from late application; discontinue use when greet tip reaches 1/2 inch.
	Apple Scab	1 lb.	31.9 lb.	Only one application allowed per year.  Extended spray schedule where fruit finish is not a concern: Continued applications may be made 5 to 7 day intervals if needed between 1/2 inch green-tip and first cover spray.  NOTE: Moderate to severe crop injury may result from this extended spray schedule. It is not intend
	Fire Blight	1.3 lb.		for fresh market apples or for apples where fruit finish is a concern as it is likely to cause f russetting. The addition of 1 to 3 pounds of hydrated lime per pound of KOCIDE® HCu may reduce frop injury.  Minimum retreatment interval is 5 days.
Avocado	Collar Rot, Crown Rot  Anthracnose, Blotch,	4 lb. 4 – 6.3 lb.	31.9 lb.	Mix in 100 gallons of water. Apply 4 gallons of suspension as a drench on the lower trunk area each tree. Apply in early spring or in fall after harvest for best results. Do not apply to foliage or fr NOTE: Do not use if soil pH is below 5.5 since copper toxicity may result.  Apply when bloom buds begin to swell and continue application at 14 to 30 day intervals for five
	Scab			six applications. Use the higher rates when conditions favor disease.  Minimum retreatment interval is 14 days.
Banana, Plantain	Sigatoka (Black and Yellow)	2.1 lb.	37.7 lb.	Apply by air in 3 gallons of water. Apply at 7 to 14 day intervals if needed. If needed, agricultu type spray oil may be added. Apply at 21 day intervals during dry periods.  Minimum retreatment interval is 7 days.
Cacao	Black Pitting  Black Pod	2.1 lb. 2 - 4.5 lb.	37.7 lb.	Mix in 100 gallons of water. Apply to the fruit stem and the basal portion of the leaf crown. At during the first and second weeks after fruit emergence.  Minimum retreatment interval is 7 days.  Begin applications at the start of the rainy season and continue while infection conditions per
				Apply at 14 to 21 day intervals if needed depending on disease severity. For drier areas, make to four applications using 2 to 4 pounds per acre according to disease incidence and planting densiminum retreatment interval is 14 days.
Coffee	Coffee Berry Disease (Colletotrichum coffeanum)	4.2 lb.	25.1 lb.	Apply first spray after flowering and before onset of long rains and then at 14 to 28 day interval needed until picking.
	Bacterial Blight (Pseudomonas syringae)	4.2 lb.	25.1 lb.	Begin spray program before the onset of long rainy periods and continue throughout the rainy seat at 14 to 21 day intervals if needed. The critical time for spraying to control this disease is just before during and after flowering(s), especially when coinciding with wet weather.
	Leaf Rust (Hemileia vastatrix)	3 – 4.2 lb.	25.1 lb.	Apply before the onset of rain and then at 14 to 21 day intervals if needed while the rains contin Use the higher rates when rainfall is heavy and disease pressure is high. Minimum retreatment interval is 14 days.
	Iron Spot (Cercospora coffeicola), Pink Disease (Corticium salmonicolor)	2 lb.	25.1 lb.	Use concentrate or dilute spray. Begin treatment at the start of wet season and continue at mon intervals for three applications.
Filbert (only for use in Washington & Oregon)	Bacterial Blight	8 – 12 lb.	47.9 lb.	Apply as a post-harvest spray. In seasons of heavy rainfall, apply a second spray when three-four of the leaves have dropped. Use the higher rates when rainfall is heavy and disease pressure is hill fineded, agricultural-type spray oil may be added.  Minimum retreatment interval is 14 days.
	Eastern Filbert Blight	8 – 12 lb.	47.9 lb.	Apply as a dilute spray in adequate water for thorough coverage. Make applications starting at I swell to bud break and continue at 14 day intervals if needed until early May. Thorough coverage essential. Use the higher rates when rainfall is heavy and disease pressure is high. If need agricultural-type spray oil or sticking agent may be added.  Minimum retreatment interval is 14 days.
Mango	Anthracnose	4 – 6.4 lb.	95.8 lb.	Apply at 7 to 30 day intervals after fruit set until harvest. Use the higher rates when rainfall is he and disease pressure is high.  Minimum retreatment interval is 7 days.
Olive	Olive Knot, Peacock Spot	5 – 8 lb.	35.9 lb.	Make first application before winter rains begin. A second application in early spring should be main disease is severe. Apply the higher rates for heavy disease pressure or when conditions far disease development.  Minimum retreatment interval is 30 days.
Peach, Nectarine	Bacterial Blast (Pseudomonas), Bacterial Canker, Bacterial Spot (Xanthomonas), Coryneum Blight (Shot Hole), Leaf Curl	8 – 16 lb.	35.9 lb.	Make first application before fall rains and a second at late dormant. For peach leaf curl, late dorm application must be made before leaf buds swell. Use the higher rates when rainfall is heavy a disease pressure is high. If needed, agricultural-type spray oil may be added.  Minimum retreatment interval is 7 days.
	Blossom Brown Rot, Coryneum Blight (Shot Hole), Leaf Curl	3 – 6 lb.	35.9 lb.	Full cover spray at pink bud. Use the higher rates when conditions favor disease.  Minimum retreatment interval is 5 days.
	Bacterial Spot	1 – 3 lb.	*35.9 lb.	Apply as a post bloom cover spray. Repeat at 5 day intervals if needed. Minimum retreatment interval is 5 days.  NOTE: Do not spray three weeks prior to harvest. Spotting of leaves and defoliation may occur from use in cover sprays. Discontinue us injury occurs.
Pear	Fire Blight	1 lb.	31.9 lb.	*Maximum single growing season application is 3.0 pounds per acre.  Apply at 5 day intervals if needed throughout the bloom period.  Minimum retreatment interval is 5 days.  NOTE: Russetting may occur in copper sensitive varieties. Excessive dosages may cause fruit rus on any variety.
	Blossom Blast (Pseudomonas)	9 – 16 lb.	31.9 lb.	Apply before fall rains and again during dormancy before spring growth starts. Use the higher rawhen disease pressure is high or when conditions favor disease development.
Pecan	Kernel Rot, Shuck Rot (Phytophthora cactorum), Zonate Leaf Spot (Cristulariella pyramidalis)	2 – 4.2 lb.	16.8 lb.	For suppression, apply in sufficient water to ensure complete spray coverage at 2 to 4 week inten- if needed, starting at kernel growth and continue until shucks open. Use the higher rates and sho spray intervals if frequent rainfall occurs. Minimum retreatment interval is 14 days.
	Ball Moss, Spanish Moss	2 – 4.2 lb.	16.8 lb.	Apply in 100 gallons of water in the spring when ball moss is actively growing, using 1.5 gallon spray per foot of tree height. Make sure to wet ball moss tufts thoroughly. The addition of a non-ic surfactant will improve control. A second application may be required after 12 months.
Pistachio	Botryosphaeria Panicle and Shoot Blight, Botrytis Blight, Late Blight (Alternaria alternata), Septoria Leaf Blight	3 – 4.2 lb.	16.8 lb.	Make initial application at bud swell and repeat on a 14 to 28 day schedule if needed. If dise conditions are severe, use the higher rates and shorter spray intervals.  Minimum retreatment interval is 14 days.
Quince	Fire Blight	1 lb.	31.9 lb.	Apply at 5 day intervals if needed throughout the bloom period. Apply in adequate water for thoror coverage. Minimum retreatment interval is 5 days.
Walnut	Walnut Blight	5 – 8 lb.	63.9 lb.	Apply first spray at early pre-bloom prior to or when catkins are partially expanded. Make addition applications during bloom and early nutlet stage on a 7 day interval if needed when frequent rais or extended periods of moisture occur. Thorough coverage of catkins, leaves and nutlets is esset for effective control.

Crop	Disease	Application Rate/Acre	Maximum Annual Rate/Acre	Use Instructions	
Bean (Dry, Green)	Brown Spot, Common Blight, Halo Blight	1 – 1.5 lb.	9.46 lb.	For protective sprays, make first application when plants are 6 inches high; repeat on a 7 to 14 day schedule if needed depending on environmental conditions. Use the higher rates for more severe disease.  Minimum retreatment interval is 7 days.	
Beet (Table Beet, Beet Greens)	Cercospora Leaf Spot	2 – 2.5 lb.	15 lb.	Begin applications when conditions first favor disease development and repeat at 10 to 14 day intervals if needed. Use the higher rates when conditions favor disease.  Minimum retreatment interval is 10 days.	
Carrot	Alternaria Leaf Spot, Cercospora Leaf Spot	2 lb.	9.98 lb.	Begin applications when disease first threatens and repeat at 7 to 14 day intervals if needed depending on disease severity.  Minimum retreatment interval is 7 days.	
Celery, Celeriac	Bacterial Blight, Cercospora Early Blight, Septoria Late Blight	2 lb.	10.6 lb.	Begin applications as soon as plants are first established in the field, repeating at 7day intervals if needed depending on disease severity and environmental conditions.  Minimum retreatment interval is 7 days.	
Crucifers (Broccoli; Brussels Sprout; Cabbage; Cabbage, Chinese; Cauliflower; Greens, Collard; Greens, Mustard; Greens, Turnip; Kale; Kohlrabi)	Black Leaf Spot (Alternaria), Black Rot (Xanthomonas), Downy Mildew	1lb.	5.29 lb.	Begin application after transplants are set in the field, or shortly after emergence of field seeded crops or when conditions favor disease development. Apply at 7 to 10 day intervals if needed. Minimum retreatment interval is 7 days.  NOTE: Reddening of older leaves may occur on broccoli and a flecking of wrapper leaves may occur on cabbage.	
Cucurbits (Cantaloupe, Cucumber, Honeydew, Muskmelon, Pumpkin, Squash, Watermelon)	Alternaria Leaf Spot, Angular Leaf Spot, Anthracnose, Downy Mildew, Gummy Stem Blight, Powdery Mildew, Watermelon Bacterial Fruit Blotch (suppression)	1.5 – 2 lb.	10.5 lb.	Begin applications prior to disease development and continue while conditions are favorable for disease development. Repeat at 5 to 7 day intervals if needed. Use the higher rates when conditions favor disease.  Minimum retreatment interval is 5 days.  NOTE: Crop injury may occur from application at higher rates and shorter intervals. Discontinue use if injury occurs.	
Eggplant	Alternaria Blight, Anthracnose, Phomopsis	1.5 lb.	15 lb.	Begin applications prior to development of disease symptoms. Repeat sprays at 7 to 10 day intervals if needed depending on disease severity.  Minimum retreatment interval is 7 days.	
Lettuce including Endive, Escarole	Downy Mildew	1 – 2 lb.	16 lb.	Begin applications when disease symptoms first appear or when conditions favor diseast development. Repeat at 5 to 10 day intervals if needed depending on disease severity. Minimum retreatment interval is 5 days.  NOTE: Determine if there is varietal sensitivity prior to use. Injury may occur to sensitive lettu varieties and under adverse weather conditions. Discontinue use if injury occurs.	
Okra	Anthracnose, Bacterial Leaf Spot, Leaf Spots, Pod Spot, Powdery Mildew	1 – 2 lb.	10.5 lb.	Begin treatment when disease first threatens and repeat every 5 to 10 days if needed depending on disease severity. Use the higher rates and shorter spray intervals when conditions favor disease. Minimum retreatment interval is 5 days.	
Onion, Garlic, Leek	Bacterial Blight,	1 – 1.5 lb.	12 lb.	Begin when plants are 4 to 6 inches high and repeat at 7 to 10 day intervals if needed depending on	
	Downy Mildew, Purple Blotch	2 lb.		disease severity. Can cause phytotoxicity to leaves. Minimum retreatment interval is 7 days.	
Pea	Powdery Mildew	1.5 lb.	7.9 lb.	Begin applications when disease symptoms first appear and repeat at weekly intervals if needed.  Minimum retreatment interval is 7 days.	
Pepper	Anthracnose, Bacterial Spot, Cercospora Leaf Spot	1.5 lb.	23.6 lb.	Begin applications when conditions first favor disease development and repeat at 3 to 10 day intervals if needed depending on disease severity.  Minimum retreatment interval is 3 days.	

are present.

VEGETABLES

Minimum retreatment interval is 7 days.

NOTE: Adequate control may not be obtained when copper tolerant species of Xanthomonas bac

(continued)

	VEGETABLES (continued)							
Crop	Disease	Application Rate/Acre	Maximum Annual Rate/Acre	Use Instructions				
Spinach	Anthracnose, Blue Mold, Cercospora Leaf Spot, Downy Mildew*, White Rust disease	1.5 lb.	7.9 lb.	Begin application when disease first appears or when conditions favor disease development. Repeat at 7 to 10 day intervals if needed. <b>NOTE</b> : Flecking may occur on spinach leaves.  Minimum retreatment interval is 7 days.				
Tomato Anthracnose, Bacterial Speck, Bacterial Spot, Early Blight, Gray Leaf Mold, Lafe Blight, Septoria Leaf Spot	1 lb.	34.7 lb. (processing)	Begin applications when disease first threatens and repeat at 3 to 10 day intervals if needed depending on disease severity. Use the higher rates when conditions favor disease.					
	Mold, Late Blight,	2 – 3 lb.	16 lb. (fresh market)	Minimum retreatment interval is 3 days.				
Watercress	Cercospora Leaf Spot	1 lb.	4 lb.	For applications made to watercress, production fields must be drained of water at least 24 hours prior to each application and water must not be reapplied to the field for a minimum of 24 hours following each application.				
				Begin applications when plants are first established in the field, repeating at 7 to 14 day intervals if needed depending on disease severity.				
				Apply using ground spray equipment at no less than 50 gallons of spray solution per acre.  Do not exceed four applications per crop.  Minimum retreatment interval is 7 days.				

	VINES							
Crop	Disease	Application Rate/Acre	Maximum Annual Rate/Acre	Use Instructions				
Grape	Black Rot, Downy Mildew, Phomopsis, Powdery Mildew	2 – 6 lb.	39.9 lb.	Begin applications at bud break with subsequent applications throughout the season depending on disease severity. Repeat at 3 day intervals if needed. Use the higher rates when conditions favor disease.  Minimum retreatment interval is 3 days.  NOTE: Foliage injury may occur on copper sensitive varieties such as Concord, Delaware, Niagara and Rosette. Either test for sensitivity or add 1 to 3 pounds of hydrated lime per pound of KOCIDE® HCu.				
Hops	Downy Mildew	1 lb.	5.3 lb.	Make crown treatment after pruning, but before training. After training, apply at 10 day intervals if needed.  Minimum retreatment interval is 10 days.  NOTE: Discontinue use two weeks before harvest.				
Kiwi	Erwinia herbicola, Pseudomonas fluorescens,	4.2 lb.	12.6 lb.	Apply in 200 gallons of water per acre. Make applications on a monthly basis.  Do not exceed three applications per crop.  Minimum retreatment interval is 30 days.				

				MISCELLANEOUS	
Crop	Disease	Application Rate/Acre	Maximum Annual Rate/Acre	Use Instructions	
Atemoya	Anthracnose	4 – 6.3 lb.	25.1 lb.	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage. Use the higher rates for severe disease. Minimum retreatment interval is 7 days.	
Carambola	Anthracnose	4.2 lb.	21 lb.	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage. Minimum retreatment interval is 7 days.	
Chives	Downy Mildew	1 lb.	5.3 lb.	Begin applications when plants are established in the field. Repeat applications every 7 to 10 days if needed depending on disease conditions.  Minimum retreatment interval is 7 days.	
Dill	Phoma Leaf Spot, Rhizoctonia Foliage Blight	1.5 lb.	7.9 lb.	Begin applications when plants are first established in the field and repeat at 7 to 10 day intervals in needed depending upon disease severity and environmental conditions. Minimum retreatment intervals 7 days.	
Ginseng	Alternaria Leaf Blight, Stem Blight	2.1 lb.	10.5 lb.	Use as a tank mix with the appropriate amount of a product containing the active ingredient iprodic in 100 gallons of water. Use in accordance with the most restrictive of label limitations a precautions. No label dosage rates are to be exceeded. This product cannot be mixed with a product containing a label prohibition against such mixing. Begin KOCIDE® HCu "iprodio applications as soon as plants have emerged in spring. Applications can be repeated every 7 ds if needed until plants become dormant in fall. Apply fungicides at least 8 hours before rein. Use of spreader-sticker or sticker is advised.  NOTE: Alternaria Leaf and Stem Blight is most severe in humid conditions such as those found the dense canopies of 2 to 4 year old Ginseng. It is very important that the stems be thoroug covered with fungicide; therefore, use a spray apparatus which distributes the fungicide through the canopy.  Minimum retreatment interval is 7 days.	
Guava	Anthracnose, Red Algae	2.4 lb.	9.84 lb.	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage. Minimum retreatment interval is 7 days.	
Litchi	Anthracnose	2.4 lb.	9.84 lb.	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage. Minimum retreatment interval is 7 days.	
Live Oak*	Ball Moss, Spanish Moss	4 lb.	40 lb.	Apply in 100 gallons of water in the spring when ball moss is actively growing, using 1.5 gallons of spray per foot of tree height. Make sure to wet ball moss tufts thoroughly. The addition of a non-ionic surfactant will improve control. A second application may be required after 12 months.  Minimum retreatment interval is 12 months.  NOTE: This product may be injurious to ornamentals grown under Live Oaks. This product may be reactive on metal and masonry surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.	
Macadamia	Anthracnose	4.7 lb.	18.9 lb.	Initiate sprays at first sign of flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage. Minimum retreatment interval is 7 days.	
	Phytophthora Blight (P. capsici), Raceme Blight (Botrytis cinerea)	3 – 4 lb.	18.9 lb.	Apply during raceme development and bloom periods. Apply in sufficient water for thorough coverage. Use the higher rates when conditions favor disease.  Minimum retreatment interval is 7 days.	
Mamey Sapote	Algal Leaf Spot, Anthracnose	3 – 4.2 lb.	16.8 lb.	Apply when conditions favor disease development. Repeat on 14 to 30 day schedule if needed as disease severity and environmental conditions dictate. Use the higher rates when conditions favor disease. Minimum retreatment interval is 14 days.	
Papaya	Anthracnose	4 – 5 lb.	42.4 lb.	Apply before disease appears. Apply at 7 day intervals if needed. The addition of an approved spreader is desirable. Use the higher rates when conditions favor disease. Minimum retreatment interval is 7 days.	
Parsley	Bacterial Blight (Pseudomonas sp.)	2 lb.	4 lb.	Begin applications when plants are first established in the field and repeat at 10 day intervals it needed depending on disease seventy and environmental conditions.  Minimum retreatment interval is 10 days.	
Passion Fruit	Anthracnose	4.7 lb.	18.8 lb.	Make initial application just before flowering and repeat on a weekly schedule until just before harvest Apply in sufficient water for thorough coverage. Minimum retreatment interval is 7 days.	
Sugar Apple (Annona)	Anthracnose	6.3 lb.	25.2 lb.	Make initial application just before flowering and repeat on a weekly schedule until just before harvest Apply in sufficient water for thorough coverage. Minimum retreatment interval is 7 days.	
Sycamore	Anthracnose	2 – 4 lb.	40 lb.	Apply as a full cover spray in 100 gallons of water or sufficient volume for thorough coverage. Make first application at bud crack and second application 7 to 10 days later at 10% leaf expansion. Use the higher rates when conditions favor disease.  Minimum retreatment interval is 7 days.	

**CONIFERS** 

\*Not registered for use in California

For use on conifers, including Douglas Fir, Fir, Juniper, Leyland Cypress, Pine and Spruce, in Christmas tree plantings and silviculture nurseries. For control of foliar diseases, apply KOCIDE® HCu as a thorough cover spray at rates ranging from 1.5 to 4 pounds per acre. Begin applications in the spring at the initiation of new growth and repeat at 7 to 30 day intervals if needed. Use the higher rates when disease pressure is severe or when environmental conditions favor disease development. Maximum annual rate per acre is 40 pounds of product.

OCIDE® HCu is registered for use on the liste Scientific Name Disease Crop Douglas Fi Pseudotsuga menziesi Abies spp. Needlecasts Anthracnose, Phomopsis Twig Dieback Juniper Juniperus spp. Leyland Cypress X Cupressocyparis leylandii Cercospora Needle Blight

Pine Pinus spp. Needlecasts Picea spp. Needlecasts Spruce Lichens: To control lichens on any of the conifers above, apply 3 to 4 pounds of KOCIDE® HCu per acre as a dormant application before new growth emerges in the spring. The addition of a non-ionic surfactant will improve control. A second application may be required after 12 months. NOTE: Do not buffer or combine with emulsifiable concentrate insecticides.

**GREENHOUSE AND SHADEHOUSE CROPS** 

Notice to User: KOCIDE® HCu may be used in greenhouses and shadehouses to control diseases on crops which appear on this label, and specific instructions have been developed for the crops listed. The grower should bear in mind that the sensitivity of crops grown in greenhouses and shadehouses differs greatly from crops grown under field conditions Neither the manufacturer nor seller has determined whether or not KOCIDE® HCu can be used safely on all greenhouse and shadehouse grown crops. The user must determine if KOCIDE® HCu can be used safely prior to commercial use. In a small area, apply the specified rates to the plants in question, e.g. foliage, fruit, etc., and observe for 7 to 10 days for symptoms of phytotoxicity prior to commercial use. Consequently, injuries arising from the use of KOCIDE® HCu on these types of greenhouse and shadehouse crops are the responsibility of the user.

Apply KOCIDE® HCu according to specific rates given for those crops in pounds per acre. Two level tablespoons of KOCIDE® HCu per 1,000 square feet is equivalent to 1.0 pound of product per acre. Apply KOCIDE® HCu in adequate water for thorough coverage of plant parts. Begin application at first sign of disease and repeat if needed; use shorter spray intervals during periods when severe disease conditions persist. For maximum annual rates per acre, refer to the crop specific directions.

NOTE: Phytotoxicity may occur on young tender flush when KOCIDE® HCu is applied to citrus seedlings grown in greenhouses or shadehouses.

Crop	Disease	Rate per 1,000 Sq. Ft.	Use instructions
Citrus (Non-Bearing Nursery)	Brown Rot, Citrus Canker, Greasy Spot, Melanose, Pink Pitting, Scab	8 TBSP.	Begin applications when disease first threatens. Repeat at 7 to 30 day intervals if needed depending on disease severity.
Cucumber	Angular Leaf Spot, Downy Mildew	3 - 4 TBSP.	Apply at 5 to 7 day intervals when plants begin to vine. Use the higher rates when conditions favor disease.
Eggplant	Alternaria Blight, Anthracnose, Phomopsis	2 - 3 TBSP.	Begin applications prior to development of disease symptoms. Repeat sprays at 7 to 10 day intervals if needed depending on disease severity.
Pepper	Bacterial Spot	2 - 3 TBSP.	Begin applications when conditions first favor disease development and repeat at 3 to 10 day intervals if needed depending on disease severity. Use the higher rates when conditions favor disease.
Tomato	Anthracnose, Bacterial Speck, Bacterial Spot, Early Blight, Gray Leaf Mold, Late Blight, Septoria Leaf Spot	4 - 6 TBSP.	Begin applications when disease first threatens and repeat at 3 to 10 day intervals if needed depending on disease severity. Use the higher rates when conditions favor disease.

### **ORNAMENTALS**

Use KOCIDE® HCu for control of bacterial and fungal diseases of foliage, flow itals in greenhouses, shade houses, outdoor nurseries and outdoor landscape

For omamental crops in dormancy, apply as a thorough cover spray at rates ranging from 1.0 to 4.0 pounds per acre of KOCIDE® HCu. When new growth is present, apply as a thorough cover spray at rates ranging from 1.0 to 2.0 pounds per acre of KOCIDE® HCu. Two level tablespoons of KOCIDE® HCu per 1,000 square feet is equivalent to 1.0 pounds of product per acre. Begin application at first sign of disease and repeat at 7 to 14 day intervals if needed; use the higher rates and shorter spray intervals during periods of frequent rains or when severe disease conditions persist. Maximum annual rate per acre is 40 pounds.

KOCIDE® HCu may be used alone or in combination with other fungicides registered for use on ornamentals as a maintenance spray. Use in accordance with the most restrictive of label limitations and precautions. Do not exceed label dosage rates. This product cannot be mixed with any product containing a label prohibition against such mixing.

Notice to User: Plant sensitivities to KOCIDE® HCu have been found to be acceptable for the specific genera and species listed on this label under the conditions tested. However phytotoxicity may occur. Due to the large number of species and varieties of ornamental and nursery plants, and the wide range of growing conditions, it is impossible to test every one for sensitivity to KOCIDE® HCu. Neither the manufacturer nor seller has determined whether or not KOCIDE® HCu can be safely used on ornamental or nursery plants not isted on this label. The user must determine if KOCIDE® HCu can be used safely prior to commercial use. In a small area, apply the specified rates to the plants in question, i.e bedding plants, foliage, etc., and observe for 7 to 10 days for symptoms of phytotoxicity prior to commercial use.

NOTE: This product may be reactive on masonry and metal surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc. Scientific Name Disease

Сгор	Scientific Name	Disease
Aglaonema*	Aglaonema spp.	Bacterial Leaf Spot
Althea (Rose of Sharon)	Hibiscus syriacus	Bacterial Leaf Spot
Andromeda, Japanese*	Pieris japonica	Leaf Spots, Twig Blight
Aralia	Dizygotheca elegantissima	Alternaria, Cercospora Leaf Spot, Xanthomonas Leaf Spot
Arborvitae	Thuja spp.	Alternaria Twig Blight, Cercospora Leaf Blight
Aster*	Aster spp.	Downy Mildew, Leaf Spots
Azalea <sup>1</sup>	Rhododendron spp.	Botrytis Blight, Cercospora Leaf Spot, Phytophthora Dieback, Powdery Mildew
Beech*	Fagus spp.	Leaf Spots
Begonia	Begonia semperflorens	Bacterial Leaf Spot (Erwinia spp., Pseudomonas spp., Xanthomonas spp.)
Bougainvillea	Bougainvillea spectabilis	Anthracnose, Bacterial Leaf Spot
Boxwood*	Buxus spp.	Leaf Spots
Camellia	Camellia japonica, C. sasanqua	Anthracnose, Bacterial Leaf Spot
Camphor Tree	Cinnamomum camphora	Pseudomonas Leaf Spot
Canna	Canna spp.	Pseudomonas Leaf Spot
Carnation <sup>1</sup>	Dianthus spp.	Alternaria Blight, Botrytis Blight, Pseudomonas Leaf Spot
Cedar*	Cedrus spp.	Tip Blight
Cherry, Nanking*	Prunus tomentosa	Bacterial Leaf Spot
Chinese Tallow Tree	Sapium sebiferum	Bacterial Leaf Spot (Pseudomonas spp., Xanthomonas spp.)
Chrysanthemum <sup>1</sup>	Chrysanthemum morifolium	Botrytis Blight, Pseudomonas Leaf Spot, Septoria Leaf Spot
Cotoneaster	Cotoneaster spp.	Botrytis Blight
Crabapple*	Malus spp.	Fire Blight
Cypress*	Cupressus spp.	Twig Blight
Dahlia	Dahlia pinnata	Alternaria Leaf Spot, Botrytis Gray Mold, Cercospora Leaf Spot
Delphinium*	Delphinium spp.	Leaf Spots
Dianthus	Dianthus spp.	Bacterial Soft Rot, Bacterial Spot
Dogwood, Flowering	Cornus florida	Anthracnose
Dogwood, Kousa*	Cornus kousa	Fungal Leaf Spots
Douglas Fir	Pseudotsuga menziesii	Rhabdocline Needlecast
Dracaena*	Dracaena marginata	Bacterial Leaf Spot
Dumb Cane*	Dieffenbachia spp.	Bacterial Leaf Spot
Dusty Miller	Senecio cineraria	Bacterial Leaf Spot (Pseudomonas cichorii)
Echinacea Echinacea	Echinacea spp.	Bacterial Leaf Spot (Pseudomonas cichorii)
Elm, Chinese	Ulmus parvifolia	Xanthomonas Leaf Spot
Euonymus		Anthracnose, Botrytis Blight
Fern Boston*	Euonymus spp.	
	Nephrolepis exaltata	Bacterial Leaf Spot
Fern, Holly	Cyrtomium falcatum	Pseudomonas Leaf Spot
Fig, Weeping*	Ficus benjamina	Bacterial Leaf Spot
Filbert (Ornamental)*	Corylus spp.	Filbert Blight
Fir*	Abies spp.	Needlecasts  Alternation Leaf Coats Retation Bard Ret. Companyon Leaf Coats
Gardenia	Gardenia jasminoides	Alternaria Leaf Spot, Botrytis Bud Rot, Cercospora Leaf Spot
Geranium	Pelargonium spp.	Alternaria Leaf Spot, Botrytis Gray Mold, Cercospora Leaf Spot
Gladiola	Gladiolus spp.	Alternaria Leaf Spot, Anthracnose, Bacterial Leaf Blight, Botrytis Gray Mold
Golden Rain Tree	Koelreuteria paniculata	Bacterial Leaf Spot
Grape Ivy*	Cissus spp.	Bacterial Leaf Spot
Hawthorn*	Crataegus spp.	Fire Blight
Hibiscus <sup>2</sup>	Hibiscus spp.	Bacterial Leaf Spot

**ORNAMENTALS** (continued) Crop Scientific Name Holly\* llex spp. Bacterial Blight, Leaf Spots Honeylocust\* Gleditsia triacanthos Bacterial Leaf Spot Honeysuckle, Tatarian' Lonicera tatarica Bacterial Leaf Spot Impatiens sallerana Bacterial Leaf Spot Indian Hawthorn<sup>3</sup> Raphiolepis indica Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Iris spp Hedera helix, H. canariensis Ivy (English, Algerian)<sup>1</sup> Xanthomonas Leaf Spot xora Ixora coccinea Xanthomonas Leaf Spot Juniper Juniperus spp. Anthracnose, Phomopsis Twig Dieback\* antana Lantana camera Bacterial Leaf Spot Cercospora Needle Blight eyland Cypress X Cupressocyparis leylandii Syringa spp. Cercospora Leaf Spot, Pseudomonas Blight\* Lilium longifloru Lily, Easter Botrytis Blight Tilia spp. Anthracnose, Leaf Blight Loblolly Bay Gordonia lasianthus Eriobotrya japonica Colletotrichum spp., Entomosporium maculata oquat Magnolia (Southern) Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot Magnolia grandiflora Magnolia (Sweet Bay) Magnolia virginiana Anthracnose Magnolia (Oriental) Bacterial Leaf Spot Magnolia soulangiana Mandevilla Mandevilla spp. Anthracnose Pseudomonas Leaf Blight Maple\* Acer spp. Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Marigold Tagetes spp. Mountain-Ash\* Sorbus spp. Fire Blight Mulberry, Contorted\* Morus bombycis Bacterial Leaf Spot Mulberry, Weeping Morus alba Bacterial Leaf Spot Narcissus spp Narcissus\* Leaf Blight Nephthytis\* Syngonium podophyllum Bacterial Leaf Spot Oak\* Quercus spp. Leaf Spots Algal Leaf Spot (Cephaleuros virescens) Oak, Laurel Quercus laurifolia Nerium oleander Bacterial Leaf Spot, Fungal Leaf Spot Oleander Oregon Grapeholly\* Mahonia aquifolium Volutella Leaf Blight Pachysandra procumben Palm, Date Phoenix canariensi Pestalotia Leaf Spot Palm, European Fan Chamaerops humilis Pestalotia Leaf Spot Chamaedorea elegans Bacterial Leaf Spor Exosporium Leaf Spot, Phytophthora Bud Rot alm, Washingtonia Pestalotia Leaf Spot Washingtonia robusta Peach (Flowering)65 Prunus spp. Bacterial Blast, Brown Rot, Fire Blight Pear (Flowering) Pyrus calleryana Fire Blight, Leaf Spots Bacterial Leaf Spot (Pseudomonas spp.\*, Xanthomonas spp.) Pentas (Egyptian Star) Pentas spp. Peony Paeonia spp. Botrytis Blight Periwinkle Catharanthus roseus, Vinca spp. Phomopsis Stem Blight hilodendron Philodendron selloum Bacterial Leaf Spot Phlox spp. Alternaria Leaf Spot hlox Photinia (Red Tip) Photinia x fraseri, P. glabra Anthracnose, Entomosporium Leaf Spot Pine\* Pinus spp. Needlecasts Pistachio Pistacia chinensis Anthracnose Hosta spp Plantain Lilv<sup>4</sup> Bacterial Leaf Spot Plum (Flowering)6\* Prunus spp. Bacterial Blast, Brown Rot, Fire Blight othos Scindapsus spp Bacterial Leaf Spot Calliandra spp. Powder Puff Plant Bacterial Leaf Spot Pyracantha spp yracantha Fire Blight, Scab Rhododendror Rhododendron spp Alternaria Flower Spot Rosa spp Black Spot, Powdery Milder Snapdragon Antirrhinum majus Anthracnose, Dieback, Downy Mildew Spathe Flower Spathiphyllum spp Bacterial Leaf Spot Spirea Spiraea spp. Fire Blight Picea spp. pruce Needlecasts Anthracnose, Leaf Spots\* Platanus spp. Sycamore Tulipa spp. Anthracnose, Botrytis Blight ulip Schefflera spp. Umbrella Tree\* Bacterial Leaf Spot Verbena Verbena spp. Xanthomonas Leaf Spot Viburnum Viburnum odoratissimum, V. Anthracnose plicatum, V. suspensum Viola (Pansy, Violet) Viola spp. Downy Mildew Willow Salix spp. Anthracnose Taxus spp Needle Bliaht

Zinnia spp Zinnia\* Leaf Spots Discoloration of foliage and/or blooms have been noted on some varieties. To prevent residues on commercial plants, do not spray immediately before selling season. Hibiscus - Do not apply to plants in flower pee

Some cultivars may be sensitive to KOCIDE® HCu

Yucca spp

Yucca (Adam's Needle)

5Apply KOCIDE® HĆu at 3.0 – 5.0 pounds per acre. Maximum annual rate per acre is 150 pounds. Do not apply any additional copper pesticide to this land for 36 months. Minimu etreatment interval is 7 days Apply dormant through bloom only.

Cercospora Leaf Spot, Septoria Leaf Spot

NOTE: Phytotoxicity may depend on varietal differences. If unfamiliar with the use of KOCIDE® HCu, apply the specified rate to a few plants and observe after 7 to 10 days for Control of Ball Moss\*, Spanish Moss\* and Lichens\* on Ornamental and Shade Trees: Apply KOCIDE® HCu in early spring when the trees are dormant. Apply 3 to 4 pounds of KOCIDE® HCu in 100 gallons of water, using 1.5 gallons of spray per foot of tree height. Be sure to thoroughly wet ball moss tufts, Spanish moss or lichens. The addition of a non-ionic surfactant will improve control. A second application may be required after 12 months.

NOTE: KOCIDE® HCu may be injurious to some ornamental plants growing beneath the trees. This product may be reactive on masonry and metal surfaces such as galvan roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc. Cold Storage Protection for Dormant Rootstock\*: To protect bare-root nursery trees from Phytophthora Crown Rot and Botrvtis. use 1.5 to 2 pounds of KOCIDE® HCu per 100 gallons of water. Apply as a dip or spray to the roots and lower stems of dormant rootstock prior to placing in cold storage. Do not apply to rootstock less than 2 years old. \*Not registered for use in California.

TURF

For control of algae in turfgrasses on sod farms, golf courses, cemeteries, and industrial turf areas. Apply 3 to 6 pounds per acre (1.1 to 2.2 oz. per 1000 square feet). Apply in sufficient water to provide adequate coverage. KOCIDE® HCu may be used alone or in combination with other registered turf fungicides as a maintenance spray. Observe all precautions and limitations on the label of each product used in tank mixes.

Minimum retreatment interval is 10 days. Maximum single application rate is 6 pounds per acre (3 pounds metallic copper equivalent). Maximum annual application rate is 42 pounds per acre (21 pounds metallic copper equivalent). NOTE: Phytotoxicity may occur depending on varietal differences. Apply the recommended rate to a small area and observe for 7 to 10 days for signs of injury. If phytotoxicity occurs, discontinue use. Do not apply in spray solutions with a pH of less than 6.5.

CHEMIGATION INSTRUCTIONS

Do not apply this product through any irrigation (chemigation) system using aluminum regardless of whether the irrigation system is flushed with water after use of this product age to the system may occur. Such application is prohibited

Apply this product only through one or more of the following types of systems: sprinkler, including center pivot, lateral move, traveler, big gun, or plastic pipe solid set system(s) which contain no aluminum parts or components. Crop injury, lack of effectiveness or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise

Shut off injection equipment after treatment and continue to operate irrigation system until KOCIDE® HCu has been cleared from the last sprinkler head.

CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS Public water system means a system for the provision to the public of pi average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply International systems contacted by places water systems must contact a universal extension processor of the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into the reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system

interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

NOTE: It must be determined if proper application equipment is available and if waste associated with its use can be properly handled. Agricultural chemicals are often reactive with the materials used in the construction of application equipment, such as aluminum, rubber and some synthetic materials. This factor should be taken into consideration when selecting proper application equipment. It is necessary that all application equipment be thoroughly flushed with clean water after each day's use.

When mixing, fill the nurse tank half full with water. Add KOCIDE® HCu slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. Stickers, spreaders, insecticides, nutrients, etc. should be added last. If compatibility is in question, use the Compatibility Jar Test before mixing a whole tank. Because of the wide variety of possible combinations which can be encountered, observe all precautions and limitations on the labels of all products used in mixtures. Agitate the mixture in the nurse tank.

KOCIDE® HCu should be added through a traveling irrigation system continuously or at the last 30 minutes of solid set irrigation systems. Shut off injection equipment after treatment and continue to operate irrigation system until KOCIDE® HCu has been cleared from the last sprinkler head.

### SPRINKLER CHEMIGATION

The system must contain a functional check valve, vacuum relief valve located on the irrigation pipeline to prevent water source contamination The pesticide injection pipeline must also contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

NOTE: It must be determined if proper application equipment is available and if waste associated with its use can be properly handled. Agricultural chemicals are often reactive with the materials used in the construction of application equipment, such as aluminum, rubber and some synthetic materials. This factor should be taken into consideration when selecting proper application equipment. It is necessary that all application equipment be thoroughly flushed with clean water after each day's use. When mixing, fill the nurse tank half full with water. Add KOCIDE® HCu slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. Stickers, spreaders, insecticides, nutrients, etc. should be added last. If compatibility is in question, use the Compatibility Jar Test before mixing a whole tank. Because of the wide variety of possible combinations which can be encountered, observe all precautions and limitations on the labels of all products used in mixtures.

Agitate the mixture in the nurse tank. NOCIDE® HCu should be added through a traveling irrigation system continuously or at the last 30 minutes of solid set irrigation systems. Shut off injection equipment after treatment and continue to operate irrigation system until KOCIDE® HCu has been cleared from the last sprinkler head.

> SPRAY DRIFT MANAGEMENT and the method of application (e.g., ground, aerial, airblast, chemigation)

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and the method of can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this pro **Droplet Size** 

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles Wind Speed Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition (approximately 3 to 10 mph), and there are no sensitive

areas within 250 feet downwind Temperature Inversions If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Other State and Local Requirements Applicators must follow all state and local pesticide drift requirements regarding application of copper compounds. Where states have more stringent regulations, they must be observed.

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates

Additional Requirements for aerial applications:

- The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter

- Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height The leases spray at the lowest religin consistent with efficacy and fight safety. But not release spray at a reight greater than 10 feet above the drup campy unless a greater neight is required for aircraft safety.

When applications are made with a crosswind, the swath must be displaced downwind. The applicator must compensate for this displacement at the up and downwind edge of the application area by adjusting the path of the aircraft upwind.

Additional requirements for Ground Boom Application: Do not apply with a nozzle height greater than 4 feet above the crop canopy.

### STORAGE AND DISPOSAL

Oo not contaminate water, food or feed by storage or disposa

PESTICIDE STORAGE: Store in a cool, dry place in original container.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. Container Disposal: (Paper Bag or Plastic Bag)

Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment by shaking and tapping sides and bottom to loosen clinging particles. When completely empty, offer for recycling if available, or dispose of bag in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States

KOCIDE® HCu is a registered trademark of Kocide LLC.
"Curtec" is a registered trademark of Bei Incorporated.
"Tre-Hold" is a registered trademark of Amvac Chemical Corporation

LIMITATION OF WARRANTY AND LIABILITY NOTICE: Read this Limitation of Warranty and Liability Before Buying or Using This Product, If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

t is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of KOCIDE. These risks can cause ineffectiveness of the product, crop injury, or injury to non-target crops or plants. WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS. KOCIDE warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the nherent risks described above, when used in accordance with the Directions for Use under normal conditions.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, KOCIDE MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL KOCIDE OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF KOCIDE OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF KOCIDE OF SELLER, THE REPLACEMENT OF THE PRODUCT.

To the extent consistent with applicable law that allows such requirement, KOCIDE or your Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify KOCIDE or your Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise, or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

(continued)