

# Jump Shot™

- INTENDED FOR COMMERCIAL USE ONLY
- A READY-TO-USE STABLE EMULSION INSECTICIDE
- FOR USE OUTDOORS AND NON-FOOD AREAS OF FOOD/FEED PROCESSING ESTABLISHMENTS, KENNELS, COMMERCIAL BUILDINGS, WAREHOUSES, THEATERS, OFFICE BUILDINGS, SCHOOLS, MOTELS, AND HOTELS.

- FOR CONTROL OF FLEAS, BROWN DOG TICKS AND LICE ON PREMISES
- FOR CONTROL OF TICKS WHICH MAY CAUSE LYME DISEASE ON PREMISES
- FOR CONTROL OF PESTS ON LIVESTOCK AND LIVESTOCK PREMISES
- PROVIDES 60-DAY (Two Months) CONTROL OF COCKROACHES
- PROVIDES 60-DAY (Two Months) CONTROL OF FLEAS
- PROVIDES 60-DAY (Two Months) CONTROL OF STORED PRODUCT PESTS

**Insecticida residual PRECAUCION AL CONSUMIDOR: Si usted no lee ingles no use este producto hasta que la etiqueta le haya sido explicada ampliamente. (TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)**

#### ACTIVE INGREDIENT:

\*Permethrin ..... **0.50%**

OTHER INGREDIENTS: ..... **99.50%**  
**100.00%**

\*(3-phenoxyphenyl) methyl (+/-) cis/trans 3-(2,2-dichloroethenyl) 2,2-dimethylcyclopropanecarboxylate. Cis/trans ratio: min. 35% (+/-) cis and max. 65% (+/-) trans.

## KEEP OUT OF REACH OF CHILDREN CAUTION

#### PRECAUTIONARY STATEMENTS

#### Hazards To Humans & Domestic Animals

**CAUTION:** Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove contaminated clothing and wash before reuse.

Sold by:

# MOMAR®

INCORPORATED

1830 Ellsworth Industrial Drive • Atlanta, GA 30318

Phone: 1-800-556-3967

*Solutions For The World Since 1947.*

EPA Reg. No. 3862-132-1553  
EPA Est. No. 1553-GA-1

**NET CONTENTS:  
(check one)**



1-GALLON



4X1  
GALLON BOX



12X1  
QUART BOX



5-GALLON



7-GALLON



20-GALLON



35-GALLON



55-GALLON



275-GALLON

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### GENERAL PRECAUTIONS AND RESTRICTIONS

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

#### SHAKE WELL BEFORE USING

Do not apply this product in a way that will contact workers or other persons, either directly or through drift; only protected handlers may be in the area during application. Do not breathe dusts, vapors, or spray mist. Except when applying to live stock, do not enter or allow others to enter until sprays have dried. When applied as a space spray or fog, do not enter or allow others to enter until vapors, mists, and aerosols have dispersed, and the treated area has been thoroughly ventilated.

Do not apply when food is present. Do not use in food areas of food handling establishments, restaurants, or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served, such as dining rooms, but excluding areas where foods may be prepared or held. All food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

#### INDOOR APPLICATION

Nonfood/feed areas are areas such as garbage rooms, laboratories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

Vacate treated areas and ventilate before reoccupying. For maximum effectiveness, a combination of spot, surface treatment and fogging is recommended.

Do not apply to classrooms while in use.

**SURFACE SPRAYING:** TO CONTROL COCKROACHES, WATERBUGS, PALMETTO BUGS, SCORPIONS, MILLIPEDES, CENTIPEDES, SOWBUGS, LUBBUGS, ANTS, SILVERFISH, FIREBRATS, SPIDERS, CRICKETS, CLOVER MITES, CHEESE MITES, GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, RUST RED FLOUR BEETLES, DRUGSTORE BEETLES, MEAL WORMS, GRAIN MITES AND CADELLES: Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry.

**TO CONTROL CARPET BEETLES:** Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving.

**TO CONTROL BEDBUGS:** Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Allow articles to dry thoroughly before use.

**TO CONTROL FLEAS (ADULTS AND LARVAE) AND TICKS, INCLUDING TICKS WHICH MAY CARRY LYME DISEASE (ADULTS AND LARVAE):** Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Allow to dry thoroughly before allowing dogs to contact treated areas.

IF REINFESTATION OCCURS REPEAT TREATMENT BUT DO NOT REPEAT MORE FREQUENTLY THAN EVERY 14 DAYS.

**TO CONTROL CLOTHES MOTHS:** Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

**TO KILL LICE AND LOUSE EGGS:** Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

**SPACE SPRAYING:** Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of House Flies, Fruit Flies, Gnats, Mosquitoes, Skipper Flies, Wasps, Hornets, Bees, Black Flies, Angoumois Grain Moths, Mediterranean Flour Moths, Indian Meal Moths and Tobacco Moths: Direct spray at an upward angle, distributing it uniformly throughout the entire area at a rate of 1 ounce per 1000 cubic feet of space. Keep area closed for 2 hours. Ventilate before reoccupying.

**FOGGING:** To kill exposed or accessible stages of the above named crawling insects, use a rate of 1 gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly when possible. Keep area closed for 2 hours. Ventilate before reoccupying.

**ON LIVESTOCK:** To protect cattle (beef and dairy), goats, sheep, hogs and horses from Horn Flies, Face Flies, House Flies, Mosquitoes and Gnats: Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

**TO CONTROL STABLE FLIES, HORSE FLIES AND DEER FLIES:** Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

**TO CONTROL BLOOD SUCKING LICE:** Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

**TO CONTROL POULTRY LICE:** Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

**TO CONTROL BEDBUGS AND MITES IN POULTRY HOUSES:** Spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

FIRST AID:	
<b>IF ON SKIN OR CLOTHING</b>	- Take off contaminated clothing. - Rinse skin immediately with plenty of water for 15-20 minutes. - Call a poison control center or doctor for treatment advice.
<b>IF IN EYES</b>	- Hold eye open and rinse slowly and gently with water for 15-20 minutes. - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. - Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-222-1222 for emergency medical treatment information.	

#### ENVIRONMENTAL HAZARDS for TERRESTRIAL APPLICATIONS

This pesticide is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage And Spill Procedures:** Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

**Pesticide Disposal:** Pesticide or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

**Container Disposal - (For containers greater than 5 gallons):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth. Ensure at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinseate into application equipment or a mix tank or store rinseate for later use or disposal. Repeat this procedure two more times.

*(For containers equal to or less than 5 gallons):* Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinseate into application equipment or a mix tank or store rinseate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.