FIRST AID Hold eye open and rinse slowly and gently with water for 15-20 Remove contact lenses, if present, after the first 5 minutes, then Call a poison control center or doctor for treatment advice. IF SWALLOWED: | • Call a poison control center or doctor immediately for treatment Have person sip a glass of water if able to swallow.
Do not induce vomiting unless told to by a poison control center or • Do not give anything by mouth to an unconscious person. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact **1-800-255-3924** for emergency medical treatment information.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate use of gastric

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS **WARNING – AVISO**

Causes substantial but temporary eve injury. Harmful if swallowed, absorbed through the skin or inhaled. Do not get in eyes, on skin, or on clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the bathroom. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- · Chemical-resistant gloves made of any waterproof material, such as polyvinyl chloride, nitrile rubber, or butyl rubber

IF IN EYES:

IF INHALED:

 Protective eyewear such as safety glasses, goggles, or face shield
 Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from

PHYSICAL AND CHEMICAL

HAZARDS

Do not mix or allow to come in contact with oxidizing agents or flame retardants. Hazardous chemical reaction may occur.

USERS SHOULD: Wash hands before eating, drinking, cnewing gum, using tobacco or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may contaminate water

through runoff. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash-waters or rinsate.

Drift and runoff may be hazardous to aquatic organisms in waters adjacent to treated areas.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries

outside of the United States. KOCIDE® is registered trademark of Kocide LLC 91411-10-70051_Kocide 2000_20180828_49_70051_.pdf



FUNGICIDE/BACTERICIDE+

Dry Flowable

ACTIVE INGREDIENT: BY WT.

*Non-public health bactericide

*Metallic Copper Equivalent 35%



FOR ORGANIC PRODUCTION

KEEP OUT OF REACH OF CHILDREN WARNING

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

EPA Reg. No. 91411-10-70051 EPA Est. No. 91411-TX-001

Manufacture For: Certis USA LLC 9145 Guilford Road, Suite 175 Columbia, MD 21046



Net: 15 lbs. (Nonrefillable Container)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer for recycling if available, or dispose of empty bag in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact ChemTel at **1-800-255-3924**, day or night.



ESL20171011

Made In U.S.A. A20170169

←.625"**→ | ←**.125" ----3.125" DFU Area ------**→**|**←**.5"**→**|**←**.5"**→**

FUNGICIDE/ BACTERICIDE .	
Dry Flowable By W Active Ingredient: By W Copper Hydroxide* (CAS No. 20427-59-2) 9 Inert Ingredients: 2 TOTAL 11	46.2%

† Non-public health bactericide * Metallic Copper Equivalent 35%

KEEP OUT OF REACH OF CHILDREN WARNING
Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

EPA Reg. No. 91411-10-70051

Kocide

CERTIS

FOR ORGANIC PRODUCTION

Columbia, MD 210 20180503	40 L S T E D For Organic Use
ESL20171011	FOR ORGANIC PRODUCTION
	FIRST AID
IF IN EYES:	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF INHALED:	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
	HOT LINE NUMBER

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-255-3924 for emergency medical treatment information. NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate use of gastric

PRECAUTIONARY STATEMENTS **HAZARDS TO HUMANS & DOMESTIC ANIMALS** WARNING - AVISO

Causes substantial but temporary eye injury. Harmful if swallowed, absorbed through the skin or inhaled. Do not get in eyes, on skin, or on clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the bathroom. Remove and wash contaminated clothing before

PERSONAL PROTECTIVE EQUIPMENT (PPE)
Mixers, loaders, applicators and other handlers must wear:

• Long-sleeved shirt and long pants
• Chemical-resistant gloves made of any waterproof material, such as polyvinyl chloride, nitrile rubber, or butyl rubber

 Shoes plus socks
 Protective eyewear such as safety glasses, goggles, or face shield
 Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. PHYSICAL AND CHEMICAL HAZARDS

Do not mix or allow to come in contact with oxidizing agents or flame retardants. Haz-

ardous chemical reaction may occur.

USER SAFETY RECOMMENDATIONS
USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the tollet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

through runoff.

This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash-waters or rinsate.

Drift and runoff may be hazardous to aquatic organisms in waters adjacent to treated

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Les provisions for presurder regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours without required PPE.

(REI) of 48 hours without required PPE.
PPE required for early entry to treated areas that is permitted under the Worker Protection
Standard and that involves contact with anything that has been treated, such as plants,
soil or water, is:
• Coveralls
• Chemical-resistant gloves made of any waterproof material, such as polyvinyl chloride, nitrile rubber, or butyl rubber
• Shoes plus socks
• Protective eyewear such as safety glasses, goggles, or face shield

AGRICULTURAL USE REQUIREMENTS (continued)

For Greenhouse Uses ONLY:

The 48-hour restricted entry interval (REI) may be reduced to 24-hour REI, provided that the following conditions are met:

For at least seven days following the application of copper-containing products in

For at least seven days following the application of copper-containing products in greenhouses:

at least one container or station designed specifically for flushing eyes is available in operating condition with the WPS- required decontamination supplies for workers entering the area treated with copper-containing products,

workers are informed orally, in a manner they can understand:
that residues in the treated area may be highly irritating to the eyes,
that they should take precautions, such as refraining from rubbing their eyes, to keep the residues out of their eyes,
that if they do get residues in their eyes, they should immediately flush their eyes with the eye flush container or eye flush station that is located with the decontamination supplies, and

nation supplies, and how to operate the eye flush container or eye flush station.

NON-AGRICULTURAL USE REQUIREMENTS The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides 40 CFR Part 170. The WPS applies when this product is used to produce agricultural plants on farms, forests,

Do not enter or allow others to enter until sprays have dried. GENERAL INSTRUCTIONS

GENERAL INSTRUCTIONS

Kocide 2000-O may be applied as an aerial, ground dilute or ground concentrate spray unless specifically directed otherwise in the specific crop use directions.

The per acre use rate of Kocide 2000-O is applicable for both dilute and concentrate spraying. Depending upon the equipment used and the specific crop, the spray volume applied per acre will differ. Refer to Minimum Recommended Spray Volume Table. Complete spray coverage is essential to assure optimum performance from Kocide 2000-O. When treating by aerial application or with low volume application or with Kocyde 2000-O. On the volume and the volume application or with Kocyde 2000-O. On the lor specific rates and timing of application by crop.

Consult the Kocide 2000-O label for specific rates and timing of application by crop. Where application rates and intervals are provided in a range (e.g. 4 to 12 ounds and 7 to 10 days), the higher rates and shorter spray intervals are recommended when rainfall is heavy and/or disease pressure is high. Use the higher rates for large mature tree crops.

SPECIAL PRECAUTIONS
The Pre-Harvest Interval (PHI) for Kocide 2000-O is 0-days unless noted.

If Kocide 2000-O is applied in a spray solution having a pH of less than 6.5, phytotoxicity

may occur.

Do not tank mix Kocide 2000-O with fungicides containing the active ingredient aluminum tris (O-ethyl phosphonate) for use on any registered crops unless appropriate precautions have been taken to buffer the spray solution because severe phytotoxicity may result. Use in accordance with the most restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing. This product may be reactive on masonry and metal surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc. Environmental conditions such as extended periods of wet weather, acid rain, etc. which alter the pH of the leaf surface may affect the performance of Kocide 2000-O resulting in possible phytotoxicity or loss of effectiveness.

Agricultural chemicals may perform in an unpredictable manner when tank mixed, especially where several products are involved. Reduced effect on pests or crop injury may occur. Unless recommended on this label or by a statellocal expert, it is advisable to test for compatibility and potential crop injury prior to commercial use of a new tank mix; otherwise, tank mixing should not be undertaken.

It must be determined if proper application equipment is available and if waste associated with its use can be properly handled. Agricultural chemicals are often reactive with the materials used in the construction of application equipment, such as aluminum, rubber and some synthetic materials. This factor should be taken into consideration when selecting proper application equipment after each day's use.

Apply this product only through one or more of the following types of systems: sprinkler, including center pivot, lateral move, traveler, big gun, or plastic pipe solid set system(s). Do not apply this product through any other type of irrigation system. In California, do not apply in systems which contain aluminum parts or components. may occur.

• Do not tank mix Kocide 2000-O with fungicides containing the active ingredient aluminum

When mixing, fill the spray tank one-half full with water. Add Kocide 2000-O slowly to tank

When mixing, fill the spray tank one-half full with water. Add Kocide 2000-O slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. Spreaders, stickers, insecticides, nutrients, etc. should be added last. If compatibility is in question, use the Compatibility Jar Test before mixing a whole tank or contact your chemical supplier. Observe all precautions and limitations on the labels of all products used in mixtures.
 It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

CROP CLASSIFICATION Grapefruit, Kumquat, Lemon, Lime, Orange, Pummelo, Tangelo, and Douglas Fir, Fir*, Juniper, Leyland Cypress*, Pine*, and Spruce*.
Alfalfa, Barley, Com*, Oats, Peanut, Potato, Sugar Beet, and Wheat.
Blackberry, Blueberry*, Cranberry, Currant, Gooseberry, Raspberry,
and Strawberry. SMALL FRUITS: TREE CROPS:

and Strawberry.

Almond, Apple, Apricot, Avocado, Banana/Plantain, Cacao, Cherry,
Coffee, Filbert, Mango, Nectarine, Olive, Peach, Pear, Pecan, Pistachio, Plum, Prune, Quince, and Walnut.
Bean, Beet, Beet Greens, Broccoli, Brussels Sprout, Cabbage, Chinese Cabbage, Cantaloupe, Carrot, Cauliflower, Celeriac, Celery,
Cumber, Egghant, Greens (Collard, Mustard and Turnip), Honeydew, Kale, Kohlrabi, Muskmelon, Okra*, Onion/Garlio/Leek, Pea,
Pepper, Pumpkin, Spinach, Squash, Tomato, Watercress, and Watermelon. VEGETABLES:

VINES: Grape, Hops, and Kiwi.

MISCELLANEOUS: Atemoya, Carambola, Chives, Dill, Ginseng, Guava, Litchi,
Macadamia, Mamey Sapote, Papaya, Parsley, Passion Fruit, Pecan,
Sugar Apple, and Sycamore.

GREENHOUSE AND SHADEHOUSE CROPS: Kocide 2000-0 may be used in greenhouses and shadehouses to control diseases on any crop on this label where physiology allows greenhouse or shadehouse culture. While specific directions are presented for Cucumber, Eggplant, Pepper and Tomato; general use may occur for any crop

on this label where physiology allows greenhouse or shadehouse culture. *Not registered for use in California. Minimum Recommended Spray Volume (Gallons Per Acre)

	•	,
When Applying	Kocide	2000-O

	Ground						
	Aerial	Dilute	Concentrate				
trus	10	800	100**				
onifers	10	100	30				
eld Crops	3	20	3				
mall Fruits	5	150	50				
ee Crops	10	400	50				
egetables	3	20	3				
nes	5	150	50				
iscellaneous	10	150	50				

**Pesticide application equipment such as "Curtec" or other similar sprayers which are capable of obtaining thorough coverage at low volumes may be used at as low as 20 gallons per acre of spray volume.

The following specific instructions are based on general application procedures. The recommendations of the State Agricultural Extension Service should be closely followed as to timing, frequency and number of sprays per season. FROST INJURY PROTECTION
BACTERIAL ICE NUCLEATION INHIBITOR

Application of Kocide 2000-O made to all crops listed on this label at rates and stages of growth indicated on this label, at least 24 hours prior to anticipated frost conditions, will afford control of ice nucleating bacteria (*Pseudomonas syringae*, *Erwinia herbicola*, and *Pseudomonas fluorescens*) and may therefore provide some protection against light frost.

Not recommended for those geographical areas where weather conditions favor sever frost. **CITRUS**

Kocide 2000-O may be mixed with dry foliar nutritionals (micronutrients) to create "Shot Bag" mixes to meet the various nutritional requirements of citrus and provide disease protection as described on this label. Kocide 2000-O per acre rates in these mixes must not exceed the maximum recommended labeled rates for disease control. Adding foliar nutritionals or other products to spray mixtures containing Kocide 2000-O and applying to citrus during the post bloom period when young fruit are present may result in sorav burn.

result in spray burn.						
Disease	Application Rate/Acre	Maximum Annual Rate/Acre	Use Instructions			
Algal Spot, Melanose, Scab	3-9 lbs.	36 lbs.	Apply as pre-bloom and post-bloom sprays. Use the higher rates when conditions favor disease. Minimum retreatment interval is 7 days.			
Greasy Spot, Pink Pitting	1.5-4.5 lbs.	36 lbs.	Apply in summer on expanded new flush. Repeat on subsequent flushes where disease pressure is severe. Use the higher rates when conditions favor disease. Minimum retreatment interval is 7 days.			
Alternaria Brown Spot	3-6 lbs.	36 lbs.	On susceptible varieties apply when the first spring flush appears and each flush thereafter. Application to fruit should start after two thirds of the petals have fallen and be repeated on a 7- to 21-day schedule if needed. Use the higher rates when conditions favor disease.			
Phytophthora Brown Rot, Septoria Spot	3-6 lbs.	36 lbs.	Begin application in fall before or just after the first rain and continue if needed. For Brown Rot only, apply to skirts of trees to a height of at least 4 feet. For control of Septoria Spot or where fruit have already been infected with Brown Rot, apply to entire tree. Apply also to bare ground one foot beyond skirt. Use the higher rates when conditions favor disease. Minimum retreatment interval is 7 days. NOTE: In California, in areas subject to copper injury, add 1/3 to 1 pound of high quality lime per pound of Kocide 2000-O.			
Phytophthora Foot Rot	0.75 lb.	36 lbs.	Mix with 1 quart of water, "Tre-Hold" or latex paint. Paint trunks of trees from the soil surface to the lowest scaffold lim Apply in May prior to summer rains and/or in the fall prior to wrapping trees for freeze protection. Treatment serves protection for up to 1 year, but does not cure existing infections. NOTE: Areas where microjet or low volume irrigation hit the tree trunk may require retreatment due to wash off.			
Citrus Canker (suppression)	2-4 lbs.	36 lbs.	Spray flushes 7 to 14 days after shoots begin to grow. Young fruit may require an additional application. Number and timing of applications will be dependent upon disease pressure. Under heavy pressure, each flush of new growth should be sprayed. Minimum retreatment interval is 7 days.			
Black Spot*	2-4 lbs.	36 lbs.	Begin treatment prior to or when disease first appears and repeat every 7 to 21 days if needed. Use the higher rates and shorter spray intervals when conditions favor disease. Minimum retreatment interval is 7 days.			

NOTE: Phytotoxicity may occur on young tender flush when Kocide 2000-O is applied to citrus seedlings grown in greenhouses or shadehouses.

CITRUS - Field Nursery Grown

apply 3 to 6 pounds of Kocide 2000-O per acre. Apply Kocide 2000-O at 28 To control Melanose, Scab, Pink Pitting, Greasy Spot, Brown Rot and for

FIELD CROPS						
Crop	Disease	Application Rate/Acre	Maximum Annual Rate/Acre	Use Instructions		
Alfalfa	Cercospora Leaf Spot, Leptosphaerulina Leaf Spot	1.5 lbs.	3.2 lbs.	Apply 10 to 14 days before each harvest or earlier if disease threatens. Repeat every 30 days if needed. NOTE : Spray injury may occur with sensitive varieties such as Lahontan.		
Corn* (Field Corn, Popcorn, Seed Corn, Sweet Corn)	Bacterial Stalk Rot	1-3 lbs.	12 lbs.	Begin treatment when disease first appears and repeat every 7- to 10-days if needed. Use the higher rates and shorter spray intervals when conditions favor disease.		
Peanut	Cercospora Leaf Spot	1-2.25 lbs.	13.5 lbs.	Begin spraying at 35 to 40 days after planting or when disease symptoms first appear and repeat at 7- to 14-day intervals if needed. Reduce sprays to 7-day intervals during humid weather. Use the higher rates when conditions favor disease. Flowable suffur may be added.		
Potato	Early Blight, Late Blight	0.75-3 lbs.	71.4 lbs.	Apply 0.75 to 1.25 lbs. at 5- to 10-day intervals if needed starting when plants are 2 to 6 inches high in locations where disease is light. Apply up to 3 pounds per acre when disease is more severe. Under conditions of severe disease, control with Kocide 2000-O will be improved by tank mixing with other compatible fungicides registered for use on potatoes. Read and follow all label instructions of tank mix partners.		
Sugar Beet	Cercospora Leaf Spot	1.5-3.74 lbs.	22.5 lbs.	Begin applications when conditions first favor disease development and repeat at 10- to 14-day intervals if needed. Use the higher rates when conditions favor disease. Addition of a spreader/sticker is recommended.		
Wheat, Barley, Oats	Fusarium Head Blight Suppression*, Helminthosporium Spot Blotch, Powdery Mildew Suppression,	1-1.5 lbs.	3.0 lbs.	Make applications for early season disease control through heading. Minimum retreatment interval is 10-days. Use higher rates when conditions favor disease. Addition of adjuvants is recommended.		

	Cidillo L
*Not registered for use in C	alifornia.

	Giullie Diotoli, Otelli Must	1	1	
*Not registered for use in (California.	•		
			SMALL FRU	ITS
Crop	Disease	Application Rate/Acre	Maximum Annual Rate/Acre	Use Instructions
Blackberry (Aurora, Boysen, Cascade, Chehalem,	Anthracnose, Cane Spot, Leaf Spot, Pseudomonas Blight, Purple Blotch, Yellow Rust	3 lbs.	28.6 lbs.	Make fall application after harvest. Apply delayed dormant spray after pruning/training in the spring. If needed, agricultural-type spray oil may be added.
Logan, Marion, Santiam, Thornless Evergreen)	Anthracnose, Cane Spot, Leaf Spot, Purple Blotch, Yellow Rust	1.5 lbs.	28.6 lbs.	Apply when leaf buds begin to open and repeat when flower buds show white. Repeat on a 7-day interval if needed. If needed, agricultural-type spray oil may be added. NOTE: Crop injury may occur if applied to foliage under certain environmental conditions such as hot or prolonged moist periods. Discontinue applications if signs of crop injury appear.
Blueberry*	Bacterial Canker	3-6 lbs.	24 lbs.	Make first application before fall rains and a second application 4 weeks later. Use the higher rates when conditions favor disease.
	Fruit Rot, Phomopsis Twig Blight	2-4 lbs.	24 lbs.	Dormant Application: Begin applications when bloom buds begin to swell. Make additional applications at 7- to 14-day intervals if needed before blooms open.
Cranberry	Fruit Rot	6 lbs.	36 lbs.	Make first application in late bloom. Apply one or two additional applications at 7- to 14-day intervals if needed depending on disease severity.
	Rose Bloom	6 lbs.	36 lbs.	Apply three sprays on 7- to 14-day schedule if needed as soon as symptoms are observed.
	Bacterial Stem Canker	6 lbs.	36 lbs.	Apply post-harvest and again in spring at bud swell. Apply one or two additional applications at 7 to 14 day intervals if needed depending on disease severity.
	Leaf Blight, Red Leaf Spot, Stem Blight, Tip Blight (<i>Monilinia</i>)	6 lbs.	36 lbs.	Apply delayed dormant spray in the spring. Repeat at 7 to 14-day intervals if needed through pre-bloom.
Currant, Gooseberry	Anthracnose, Leaf Spot	7.5 lbs.	45.7 lbs.	Make initial application after first leaves have expanded. Continue on a 10- to 14-day schedule if needed during wet conditions in the spring. Make an additional application after harvest.
Raspberry	Anthracnose, Cane Spot, Leaf Spot, Pseudomonas Blight, Purple Blotch, Yellow Rust	3 lbs.	28.6 lbs.	Make fall application after harvest. Apply delayed dormant spray after training in the spring. If needed, agricultural-type spray oil may be added.
	Anthracnose, Cane Spot, Leaf Spot, Purple Blotch, Yellow Rust	1.5 lbs.	28.6 lbs.	Apply when leaf buds begin to open and repeat when flower buds show white. Repeat on a 7-day interval if needed. If needed, agricultural-type spray oil may be added. NOTE: Crop injury may occur if applied to foliage under certain environmental conditions such as hot or prolonged moist periods. Discontinue applications if signs of crop injury appear.
Strawberry	Angular Leaf Spot (Xanthomonas), Leaf Blight, Leaf Scorch, Leaf Spot	1.5-2.25 lbs.	23.4 lbs.	Begin application when plants are established and continue on a 7-day schedule throughout the season. Apply in at least 20 gallons of water. Use the higher rates when conditions favor disease. NOTE: Discontinue applications if signs of crop injury appear.

Crop	Disease	Application	Maximum	Use Instructions
Almond only	Destarial Diagram	Rate/Acre	Annual Rate/Acre	Abroad Only Fasherhaid bleet andrel is spiritle injected authority with
Almond only	Bacterial Blast	0.57 lb.	51.4 lbs.	Almond Only: For bacterial blast control in sprinkler irrigated orchards or whe disease is severe, apply 0.75 pound per acre post-bloom at 5- to 14-day interval if needed or just before sprinkling.
Almond, Apricot, Cherry, Plum, Prune	Bacterial Blast (<i>Pseudomonas</i>), Bacterial Canker, Coryneum Blight (Shot Hole)	6-12 lbs.	51.4 lbs.	Make first application before fall rains and a second at late dormant. Use thigher rates when conditions favor disease. Minimum retreatment interval is days. If needed, agricultural-type spray oil may be added. For Cherries: Where disease is severe, an additional application shortly aft harvest may be required. NOTE: Foliar injury may occur from post-bloom sprays on almonds, especia on NePlus varieties.
	Blossom Brown Rot, Coryneum Blight (Shot Hole)	4.3 lbs.	51.4 lbs.	Apply during early bloom. Do not apply after full bloom or injury may occur. U the higher rates when rainfall is heavy and disease pressure is high. Minimu
	Black Knot* (Plum)	3-4.3 lbs.	51.4 lbs.	retreatment interval is 5 days. Make an application at bud swell up to early bloom for early season disea suppression. Apply before full bloom. Minimum retreatment interval is 5 day Use the higher rates when rainfall is heavy and disease pressure is high. NOTE: To avoid plant injury, do not use after full bloom.
	Cherry Leaf Spot* (Sour Cherries Only)	4.3 lbs.	51.4 lbs.	Apply at petal fall as well as 1 to 2 times after petal fall. Use the lower rat where disease infection is light and use the higher rates for a dormant application or where diseases infection is moderate to heavy. Minimum retreatment inten is 5 days. Do not apply to sweet cherry or the English Morello variety as sever injury will result. The addition of 1 to 3 pounds of hydrated lime per pound Kocide 2000-O may reduce crop injury. NOTE: Moderate to severe injury such as leaf spotting and defoliation may occ from post-bloom applications.
Apple	Anthracnose, Blossom Blast, European Canker (<i>Nectria</i>),	9-12 lbs.	45.7 lbs.	Apply before fall rains. Use the higher rates when conditions favor disease. NOTE: Use on yellow varieties may cause discoloration. To avoid discoloratio
	Shoot Blast (Pseudomonas) Apple Scab*, Fire Blight	6-12 lbs.	45.7 lbs.	pick before spraying. Make application between silver-tip and green-tip. Apply as a full cover spray tearly season disease suppression. NOTE: Moderate to severe crop injury may occur from late application.
	Apple Scab*	0.75-1.75 lbs.	45.7 lbs.	discontinue use when green-tip reaches 1/2 inch. Extended spray schedule where fruit finish is not a concern: Continue
		0.5.0.75.11		applications may be made at 5- to 7-day intervals if needed between 1/2 in green-tip and first cover spray. NOTE: Moderate to severe crop injury may result from this extended spr. schedule. It is not intended for fresh market apples or for apples where fruit finis is a concern as it is likely to cause fruit russetting. The addition of 1 to 3 poun
	Fire Blight* Collar Rot, Crown Rot	0.5-0.75 lb. 3 lbs.	45.7 lbs.	of hydrated lime per pound of Kocide 2000-O may reduce crop injury. Mix in 100 gallons of water. Apply 4 gallons of suspension as a drench on the lower trunk area of each tree. Apply in early spring or in fall after harvest for be results. Do not apply to foliage or fruit. NOTE: Do not use if soil pH is below 5.5 since copper toxicity may result.
Avocado	Anthracnose, Blotch, Scab	6-9 lbs.	54 lbs.	Apply when bloom buds begin to swell and continue application at 14- to 30-di intervals for five to six applications. Use the higher rates when conditions fav disease.
Banana, Plantain	Sigatoka (Black and Yellow)	1.5 lbs.	54 lbs.	Apply at 7- to 14-day intervals if needed.
	Black Pitting	3 lbs.	54 lbs.	Mix in 100 gallons of water. Apply to the fruit stem and the basal portion of the leaf crown. Apply during the first and second weeks after fruit emergence.
Cacao	Black Pod	1.5-6.4 lbs.	45 lbs.	Begin applications at the start of the rainy season and continue while infectic conditions persist. Apply 1.5 to 3.5 lbs. at 14-to 21-day intervals if need depending on disease severity. For drier areas, make two to four application using 4.5 to 6.4 pounds per acre according to disease incidence and plantifiednsity.
Coffee	Coffee Berry Disease (Colletotrichum coffeanum)	4.5-6 lbs.	36 lbs.	Apply first spray after flowering and before onset of long rains and then at 14- 28-day intervals if needed until picking. Use the higher rates when condition favor disease.
	Bacterial Blight (Pseudomonas syringae)	4.5-6 lbs.	36 lbs.	Begin spray program before the onset of long rainy periods and continum throughout the rainy season at 14- to 21-day intervals if needed. The critical time for spraying to control this disease is just before, during and after flowering(sespecially when coinciding with wet weather. Use the higher rates when rainfing is heavy and disease pressure is high.
	Leaf Rust (Hemileia vastatrix)	1.5-3 lbs.	36 lbs.	Apply before the onset of rain and then at 14- to 21-day intervals if needed whithe rains continue. Use the higher rates when rainfall is heavy and diseast pressure is high.
	Iron Spot (Cercospora coffeicola), Pink Disease (Corticium salmonicolor)	1.5 lbs.	36 lbs.	Use concentrate or dilute spray. Begin treatment at the start of wet season ar continue at monthly intervals for three applications.
Filbert (WA, OR only)	Bacterial Blight	12-17 lbs.	68.6 lbs.	Apply as a post-harvest spray. In seasons of heavy rainfall, apply a second sprawhen three-fourths of the leaves have dropped. Wait at least 14 days betwee applications. Use the higher rates when rainfall is heavy and disease pressure high. If needed, agricultural-type spray oil may be added.
	Eastern Filbert Blight	12-17 lbs.	68.6 lbs.	Apply as a dilute spray in adequate water for thorough coverage. Mal applications starting at bud swell to bud break and continue at 14-day intervals needed until early May. Thorough coverage is essential. Use the higher rate when rainfall is heavy and disease pressure is high. If needed, agricultural-type spray oil or sticking agent may be added.
Mango	Anthracnose	6-9 lbs.	137 lbs.	Apply at 7-day intervals after fruit set until harvest. Use the higher rates whe rainfall is heavy and disease pressure is high.
Olive	Olive Knot, Peacock Spot	6-9 lbs.	51 lbs.	Make first application before winter rains begin. A second application in ear spring should be made if disease is severe. Apply the higher rates for hear disease pressure or when conditions favor disease development. Minimu retreatment interval is 30 days.
Peach, Nectarine	Bacterial Blast (<i>Pseudomonas</i>), Bacterial Canker, Bacterial Spot (<i>Xanthomonas</i>), Coryneum Blight (Shot Hole), Leaf Curl	6-12 lbs.	51.4 lbs.	Make first application before fall rains and a second at late dormant. For pear leaf curl, late dormant application must be made before leaf buds swell. Use th higher rates when rainfall is heavy and disease pressure is high. If neede agricultural-type spray oil may be added. Minimum retreatment interval is 7-day
	Blossom Brown Rot, Coryneum Blight (Shot Hole), Leaf Curl	4.3 lbs.	51.4 lbs.	Full cover spray at pink bud. Minimum retreatment interval is 5 days.
	Bacterial Spot	0.75 lb.	51.4 lbs.	Post-bloom application applied at first and second cover sprays. Minimu retreatment interval is 5 days. NOTE: Do not spray 3 weeks prior to harvest. Use only recommended rate Spotting of leaves and defoliation may occur from use in cover sprays.
Pear	Fire Blight	0.75 lb.	45.7 lbs.	Apply at 5-day intervals if needed throughout the bloom period. NOTE: Russetting may occur in copper sensitive varieties. Excessive dosage may cause fruit russet on any variety.
	Blossom Blast (Pseudomonas)	9-12 lbs.	45.7 lbs.	Apply before fall rains and again during dormancy before spring growth start Use the higher rates when disease pressure is high or when conditions fav disease development.
Pecan	Kernel Rot, Shuck Rot (Phytophthora cactorum), Zonate Leaf Spot (Cristulariella pyramidalis)	1.5-3 lbs.	24 lbs.	For suppression, apply in sufficient water to ensure complete spray coverage 14- to 28-day intervals if needed, starting at kernel growth, and continue un shucks open. Use the higher rates and shorter spray intervals if frequent rainfroccurs.
	Ball Moss*, Spanish Moss*	4.5-6 lbs.	24 lbs.	Apply in 100 gallons of water in the spring when ball moss is actively growin using 1 1/2 gallons of spray per foot of tree height. Make sure to wet ball most utifs thoroughly. The addition of a non-ionic surfactant will improve control. second application may be required after 12 months.
Pistachio	Botryosphaeria Panicle and Shoot Blight, Botrytis Blight, Late Blight (<i>Alternaria</i> <i>alternata</i>), Septoria Leaf Blight	3-6 lbs.	24 lbs.	Make initial application at bud swell and repeat on a 14- to 28-day schedule needed. If disease conditions are severe, use the higher rates and shorter spraintervals.
Quince	Fire Blight	0.75 lb.	45.7 lbs.	Apply at 5-day intervals if needed throughout the bloom period. Apply in adequa water for thorough coverage.
Walnut	Walnut Blight	6-9 lbs.	91 lbs	Apply first spray at early pre-bloom prior to or when catkins are partial expanded. Make additional applications during bloom and early nutlet stage or a 7-day interval if needed when frequent rainfall or extended periods of moistum occur. Thorough coverage of catkins, leaves and nutlets is essential for effective control.

VEGLIABLES					
Crop	Disease	Application Rate/Acre	Maximum Annual Rate/Acre	Use Instructions	
Bean (Dry, Green)	Brown Spot, Common Blight, Downy Mildew*, Halo Blight	0.75-2.25 lbs.	13.5 lbs.	For protective sprays, make first application when plants are 6 inches high; repeal on a 7- to 14-day schedule if needed depending on environmental conditions. Use the higher rates for more severe disease.	
Beet (Table Beet, Beet Greens)	Cercospora Leaf Spot	1.5-3.74 lbs.	22.5 lbs.	Begin applications when conditions first favor disease development and repea at 10- to 14-day intervals if needed. Use the higher rates when conditions favor disease.	
Carrot	Alternaria Leaf Spot, Cercospora Leaf Spot	1.5-2.75 lbs.	14.3 lbs.	Begin applications when disease first threatens and repeat at 7- to 14-day intervals if needed depending on disease severity.	
Celery, Celeriac	Bacterial Blight, Cercospora Early Blight, Septoria Late Blight	1.5 lbs.	15.1 lbs.	Begin applications as soon as plants are first established in the field, repeating at 7-day intervals if needed depending on disease severity and environmental conditions.	
Crucifers (Broccoli; Brussels Sprout; Cabbage; Cabbage, Chinese; Cauliflower; Greens, Collard; Greens, Mustard; Greens, Turnip; Kale; Kohlrabi)	Black Leaf Spot (Alternaria), Black Rot (Xanthomonas), Downy Mildew	0.75-1.5 lbs.	7.57 lbs.	Begin application after transplants are set in the field, or shortly after emergence of field seeded crops or when conditions favor disease development. Apply a 7- to 10-day intervals if needed. Use the higher rates when conditions favor disease NOTE: Reddening of older leaves may occur on broccoli and a flecking of wrapper leaves may occur on cabbage.	
Cucurbits (Cantaloupe, Cucumber, Honeydew, Muskmelon, Pumpkin, Squash, Watermelon)	Alternaria Leaf Spot, Angular Leaf Spot, Anthracnose, Downy Mildew, Gummy Stem Blight, Powdery Mildew, Watermelon Bacterial Fruit Blotch (suppression)	1-2.25 lbs.	15 lbs.	Begin applications prior to disease development and continue while conditions are favorable for disease development. Repeat at 5- to 7-day intervals if needed Use the higher rates when conditions favor disease. NOTE: Crop injury may occur from application at higher rates and shorte intervals. Discontinue use if injury occurs.	
Eggplant	Alternaria Blight, Anthracnose, Phomopsis	1.5 lbs.	22.6 lbs.	Begin applications prior to development of disease symptoms. Repeat sprays at 7- to 10-day intervals if needed depending on disease severity.	
Okra*	Anthracnose, Bacterial Leaf Spot, Leaf Spots, Pod Spot, Powdery Mildew	1.5-3 lbs.	15 lbs.	Begin treatment when disease first threatens and repeat every 5- to-10 days in needed depending on disease severity. Use the higher rates and shorter spray intervals when conditions favor disease.	
Onion, Garlic, Leek	Bacterial Blight, Downy Mildew, Purple Blotch	0.75 - 1.5 lbs.	17.1 lbs.	Begin when plants are 4 to 6 inches high and repeat at 7- to 10-day intervals in needed depending on disease severity. Can cause phytotoxicity to leaves.	
Pea	Powdery Mildew	1-2.25 lbs.	11.3 lbs.	Begin applications when disease symptoms first appear and repeat at 7-day intervals if needed. Use the higher rates when conditions favor disease.	
Pepper	Anthracnose, Bacterial Spot, Cercospora Leaf Spot	1.5-2.25 lbs.	33.9 lbs.	Begin applications when conditions first favor disease development and repeal at 3- to 10-day intervals if needed depending on disease severity. Use the higher rates when conditions favor disease.	
Spinach	Anthracnose, Blue Mold, Cercospora Leaf Spot, White Rust disease	1.5-2.25 lbs.	11.3 lbs.	Begin application when disease first appears or when conditions favor disease development. Repeat at 7- to 10-day intervals if needed. Use the higher rates when conditions favor disease. NOTE: Flecking may occur on spinach leaves.	
Tomato	Anthracnose, Bacterial Speck, Bacterial Spot, Early Blight,	1.5 lbs. (processing)	49.7 lbs. (processing)	Begin applications when disease first threatens and repeat at 3- to 10-d intervals if needed depending on disease severity. Where applicable, use to	
	Gray Leaf Mold, Late Blight, Septoria Leaf Spot	1.5-3 lbs. (fresh market)	22.8 lbs. (fresh market)	higher rates when conditions favor disease.	
Watercress	Cercospora Leaf Spot	1.5 lbs.	6.06 lbs.	Begin applications when plants are first established in the field, repeating at 7-to 14-day intervals if needed depending on disease severity. Do not exceed four applications per crop. Apply using ground spray equipment at no less than 50 gallons of spray solution per acre.	

VEGETABLES

NOTE: Adequate control may not be obtained when copper tolerant species of


anthomonas bacteria are present.

*Not registered for use in California

*Not registered for use in California

	VINES							
Crop	Disease	Application Rate/Acre	Maximum Annual Rate/Acre	Use Instructions				
Grape	Black Rot, Downy Mildew, Phomopsis, Powdery Mildew	1.5-3 lbs.	57.1 lbs.	Begin applications at bud break with subsequent applications throughout the season depending on disease severity. Repeat at 3-day intervals if needed. Use the higher rates when conditions favor disease. NOTE: Foliage in				
Hops	Downy Mildew	1.5 lbs.	7.57 lbs.	Make crown treatment after pruning, but before training. After training, apply at 10-day intervals if needed. NOTE: Discontinue use two weeks before harvest.				
Kiwi	Erwinia herbicola, Pseudomonas fluorescens, Pseudomonas syringae	6 lbs.	18 lbs.	Apply in 200 gallons of water per acre. Make applications on a 30-day basis. A maximum of three applications may be made.				

	MISCELLANEOUS						
Crop	Disease	Application Rate/Acre	Maximum Annual Rate/Acre	Use Instructions			
Atemoya	Anthracnose	2.25-3.5 lbs.	36 lbs.	Make initial application just before flowering and repeat on a 7-day schedule unl just before harvest. Apply in sufficient water for thorough coverage. Use th higher rates for severe disease.			
Carambola	Anthracnose	4.5-6 lbs.	30 lbs.	Make initial application just before flowering and repeat on a 7-day schedule unl just before harvest. Apply in sufficient water for thorough coverage. Use th higher rates for severe disease.			
Chives	Downy Mildew	1.5 lbs.	7.57 lbs.	Begin applications when plants are established in the field. Repeat application every 7 to 10 days if needed depending on disease conditions.			
Dill	Phoma Leaf Spot, Rhizoctonia Foliage Blight	1.5-2.25 lbs.	11.3 lbs.	Begin applications when plants are first established in the field and repeat at 7 to 10-day intervals if needed depending upon disease severity and environments conditions. Use the higher rates when conditions favor disease.			
Ginseng	Altemaria Leaf Blight, Stem Blight	2-3 lbs.	15 lbs.	Use as a tank mix with the appropriate amount of a product containing the activing redient iprodione. Use label rate of iprodione in 100 gallons of water. Use i accordance with the most restrictive of label limitations and precautions. No lab dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing. Begin Kocide 2000-C "iprodione" applications as soon as plants have emerged in spring. Application should be repeated every 7 days if needed until plants become dommant in Apply tungicides at least 8 hours before rain. Use of a spreader-sticker or stick is advised. NOTE: Alternaria Leaf and Stem Blight is most severe in humid conditions suc as those found in the dense canopies of 2- to 4-year old Ginseng. It is ver important that the stems be thoroughly covered with fungicide; therefore, use spray apparatus which distributes the fungicide throughout the canopy.			
Guava	Anthracnose, Red Algae	2.25-3.5 lbs.	14.1 lbs.	Make initial application just before flowering and repeat every 7 days until jubefore harvest. Apply in sufficient water for thorough coverage. Use the higher rates for severe disease.			
Litchi	Anthracnose	2.25-3.5 lbs.	14.1 lbs.	Make initial application just before flowering and repeat every 7 days until just before harvest. Apply in sufficient water for thorough coverage. Use the higher rates for severe disease.			



CONIFERS

For use on conifers, including Douglas Fir, Fir*, Juniper, Leyland Cypress*, Pine* and Spruce*, in Christmas tree plantings, forest stands and silviculture nurseries.

For control of foliar diseases, apply Kocide 2000-O as a thorough cover spray at rates ranging from 1.5 to 3 pounds per acre. Begin applications in the spring at the initiation of new growth and repeat at 7 to 30 day intervals if needed. Use the higher rates when disease pressure is severe or when environmental conditions favor disease development. Maximum annual rate per acre is 57.1 lbs. ended for use on the listed conifers for control of the following diseases

Crop	Scientific Name	Disease	
Douglas Fir	Pseudotsuga menziesii	Rhabdocline Needlecast	
Fir*	Abies spp.	Needlecasts	
Juniper	Juniperus spp.	Anthracnose, Phomopsis Twig Dieback*	
Leyland Cypress*	X Cupressocyparis leylandii	Cercospora Needle Blight	
Pine*	Pinus spp.	Needlecasts	
Spruce*	Picea spp.	Needlecasts	
Lichens*: To control lichens on any of the conifers above, apply 6 to 10 pounds of Kocide 2000-O per acre as a dormant application before new growth emerges in the spring. The addition of a non-ionic surfactant will improve control. A second application may be required after 12 months.			

NOTE: Do not buffer or combine with emulsifiable concentrate insecticides. *Not registered for use in California.

2

GREENHOUSE AND SHADEHOUSE CROPS

Notice to User: Kocide 2000-O may be used in greenhouses and shadehouses to control diseases on crops which appear on this label, and specific instructions have been developed for the crops listed. The grower should bear in mind that the sensitivity of crops grown in greenhouses and shadehouses differs greatly from crops grown under field conditions. conditions.

Neither the manufacturer nor seller has determined whether or not Kocide 2000-O can be used safely on all greenhouse and shadehouse grown crops. Consequently; injury arising from the use of Kocide 2000-O on these types of greenhouse and shadehouse crops is the responsibility of the user. The user should determine if Kocide 2000-O can be used safely prior to commercial use. In a small area, apply the recommended rates to the plants in question, i.e., foliage, fruit, etc., and observe for 7 to 10 days for symptoms of

phytotoxicity prior to commercial use.

Apply Kocide 2000-O according to specific rates given for those crops in pounds per acre. One level tablespoon of Kocide 2000-O per 1,000 square feet is equivalent to 1.5 pounds of product per acre. Kocide 2000-O should be applied in adequate water for thorough coverage of plant parts. Begin application at first sign of disease and repeat if needed; use shorter spray intervals during periods when severe disease conditions persist. For maximum annual rates per acre, refer to the crop specific directions.

NOTE: Phytotoxicity may occur on young tender flush when Kocide 2000-O is applied to citrus seedlings grown in greenhouses or shadehouses.

Crop	Disease	Rate per 1,000 Sq Ft	Use Instructions
Citrus (Non-Bearing Nursery)	Brown Rot, Citrus Canker, Greasy Spot, Melanose, Pink Pitting, Scab	3 TBSP	Begin applications when disease first threatens. Repeat at 7- to 30-day intervals if needed depending on disease severity.
Cucumber	Angular Leaf Spot, Downy Mildew	1 - 2 1/2 TBSP	Apply at 5- to 7-day intervals when plants begin to vine. Use the higher rates when conditions favor disease.
Eggplant	Alternaria Blight, Anthracnose, Phomopsis	1 1/2 TBSP	Begin applications prior to development of disease symptoms. Repeat sprays at 7- to 10-day intervals if needed depending on disease severity.
Pepper	Bacterial Spot	1 1/2 - 2 1/2 TBSP	Begin applications when conditions first favor disease development and repeat at 3- to 10-day intervals if needed depending on disease severity. Use the higher rates when conditions favor disease.
Tomato	Anthracnose, Bacterial Speck, Bacterial Spot, Early Blight, Gray Leaf Mold, Late Blight, Septoria Leaf Spot	1 1/2 - 3 TBSP	Begin applications when disease first threatens and repeat at 3- to 10-day intervals if needed depending on disease severity. Use the higher rates when conditions favor disease.

ORNAMENTALS

Use Kocide 2000-O for control of bacterial and fungal diseases of foliage, flower entals in greenhouses, shade houses, outdoor nurseries and outdoor landscape

Use Kocide 2000-O for control of bacterial and fungal diseases of foliage, flowers and stems on ornamentals in greenhouses, shade houses, outdoor nurseries and outdoor landscape plantings.

For ornamental crops in dormancy, apply as a thorough cover spray at rates ranging from 0.75 to 2 pounds per acre of Kocide 2000-O. When new growth is present, apply as a thorough cover spray at rates ranging from 0.75 to 2 pounds per acre of Kocide 2000-O. One level tablespoon of Kocide 2000-O per 1.000 square feet is equivalent to 1.5 pounds of product per acre. Begin application at first sign of disease and repeat at 7- to 14-day intervals if needed; use the higher rates and shorter spray intervals during periods of frequent rains or when severe disease conditions persist. Maximum seasonal rate per acre is 57.1 lbs.

Kocide 2000-O may be used alone or in combination with other fungicides registered for use on ornamentals as a maintenance spray. Use in accordance with the most restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

Notice to User: Plant sensitivities to Kocide 2000-O have been found to be acceptable for the specific genera and species listed on this label under the conditions tested. However, phytotoxicity may occur. Due to the large number of species and varieties of ornamental and nursery plants, and the wide range of growing conditions, it is impossible to test every one for sensitivity to Kocide 2000-O. Neither the manufacturer nor seller has determined whether or not Kocide 2000-O can be safely used on ornamental or nursery plants not listed on this label. The user should determine if Kocide 2000-O can be used safely prior to commercial use.

NOTE: This product may be reactive on masonry and metal surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.

_	lasonry and metal surfaces such as ga	alvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.	
Crop	Scientific Name	Disease	
Aglaonema*	Aglaonema spp.	Bacterial Leaf Spot	
Althea (Rose of Sharon)	Hibiscus syriacus	Bacterial Leaf Spot	
Andromeda, Japanese*	Pieris japonica	Leaf Spots, Twig Blight	
Aralia	Dizygotheca elegantissima	Alternaria, Cercospora Leaf Spot, Xanthomonas Leaf Spot	
Arborvitae	Thuja spp.	Alternaria Twig Blight, Cercospora Leaf Blight	
Aster*	Aster spp.	Downy Mildew, Leaf Spots	
Azalea 1	Rhododendron spp.	Botrytis Blight, Cercospora Leaf Spot, Phytophthora Dieback, Powdery Mildew	
Beech*	Fagus spp.	Leaf Spots	
Begonia	Begonia semperflorens	Bacterial Leaf Spot (Erwinia spp., Pseudomonas spp., Xanthomonas spp.)	
Bougainvillea	Bougainvillea spectabilis	Anthracnose, Bacterial Leaf Spot	
Boxwood*	Buxus spp.	Leaf Spots	
Camellia	Camellia japonica, C. sasangua	Anthracnose, Bacterial Leaf Spot	
Camphor Tree	Cinnamomum camphora	Pseudomonas Leaf Spot	
Canna	Canna spp.	Pseudomonas Leaf Spot	
Carnation ¹	Dianthus spp.	Alternaria Blight, Botrytis Blight, Pseudomonas Leaf Spot	
Cedar*	Cedrus spp.	Tip Blight	
Cherry, Nanking*	Prunus tomentosa	Bacterial Leaf Spot	
Chinese Tallow Tree	Sapium sebiferum	·	
	'	Bacterial Leaf Spot (Pseudomonas spp., Xanthomonas spp.)	
Chrysanthemum ¹	Chrysanthemum morifolium	Botrytis Blight, Pseudomonas Leaf Spot, Septoria Leaf Spot	
Cotoneaster	Cotoneaster spp.	Botrytis Blight	
Crabapple*	Malus spp.	Fire Blight	
Cypress*	Cupressus spp.	Twig Blight	
Dahlia	Dahlia pinnata	Alternaria Leaf Spot, Botrytis Gray Mold, Cercospora Leaf Spot	
Delphinium*	Delphinium spp.	Leaf Spots	
Dianthus	Dianthus spp.	Bacterial Soft Rot, Bacterial Spot	
Dogwood, Flowering	Cornus florida	Anthracnose	
Dogwood, Kousa*	Cornus kousa	Fungal Leaf Spots	
Douglas Fir	Pseudotsuga menziesii	Rhabdocline Needlecast	
Dracaena*	Dracaena marginata	Bacterial Leaf Spot	
Dumb Cane*	Dieffenbachia spp.	Bacterial Leaf Spot	
Dusty Miller	Senecio cineraria	Bacterial Leaf Spot (Pseudomonas cichorii)	
Echinacea	Echinacea spp.	Bacterial Leaf Spot (Pseudomonas cichorii)	
Elm, Chinese	Ulmus parvifolia	Xanthomonas Leaf Spot	
Euonymus	Euonymus spp.	Anthracnose, Botrytis Blight	
Fern Boston*	Nephrolepis exaltata	Bacterial Leaf Spot	
Fern, Holly	Cyrtomium falcatum	Pseudomonas Leaf Spot	
Fig, Weeping*	Ficus benjamina	Bacterial Leaf Spot	
Filbert (Ornamental)*	Corylus spp.	Filbert Blight	
Fir*	Abies spp.	Needlecasts	
Gardenia	Gardenia jasminoides	Alternaria Leaf Spot, Botrytis Bud Rot, Cercospora Leaf Spot	
Geranium	Pelargonium spp.	Alternaria Leaf Spot, Botrytis Gray Mold, Cercospora Leaf Spot	
Gladiola			
	Gladiolus spp.	Alternaria Leaf Spot, Anthracnose, Bacterial Leaf Blight, Botrytis Gray Mold	
Golden Rain Tree	Koelreuteria paniculata	Bacterial Leaf Spot	
Grape Ivy*	Cissus spp.	Bacterial Leaf Spot	
Hawthorn*	Crataegus spp.	Fire Blight	
Hibiscus ⁴	Hibiscus spp.	Bacterial Leaf Spot	
Holly*			
	llex spp.	Bacterial Blight, Leaf Spots	
Honeylocust*	Gleditsia triacanthos	Bacterial Leaf Spot	
Honeysuckle, Tatarian*	Gleditsia triacanthos Lonicera tatarica	Bacterial Leaf Spot Bacterial Leaf Spot	
Honeysuckle, Tatarian*	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot	
Honeysuckle, Tatarian* Impatiens Indian Hawthom ⁵	Gleditsia triacanthos Lonicera tatarica	Bacterial Leaf Spot Bacterial Leaf Spot	
Honeysuckle, Tatarian* Impatiens Indian Hawthom ⁵ Iris ⁶ *	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp.	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot	
Honeysuckle, Tatarian* Impatiens Indian Hawthom ⁵ Iris ^{6*} Ivy (English, Algerian) ¹	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot	
Honeysuckle, Tatarian* Impatiens Indian Hawthom ⁵ Iris ⁶ *	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp.	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot	
Honeysuckle, Tatarian* Impatiens Indian Hawthom ⁵ Iris ^{6*} Ivy (English, Algerian) ¹	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn ⁵ Iris ^{6*} Ivy (English, Algerian) ¹ Ixora	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn ⁵ Iris ^{6*} Ivy (English, Algerian) ¹ Ixora Juniper	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp.	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback*	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn ⁵ Iris ^{6*} Ivy (English, Algerian) ¹ Ixora Juniper Lantana	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn ⁵ Iris ^{6*} Ivy (English, Algerian) ¹ Ixora Juniper Lantana Leyland Cypress*	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot Cercospora Needle Blight	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn 5 Iris 6* Ivy (English, Algerian) 1 Ixora Juniper Lantana Leyland Cypress* Lilac	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii Syringa spp.	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot Cercospora Needle Blight Cercospora Leaf Spot, Pseudomonas Blight*	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn 5 Iris 6* Ivy (English, Algerian) 1 Ixora Juniper Lantana Leyland Cypress* Lilac Lily, Easter 2	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii Syringa spp. Lilium longiflorum	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot Cercospora Needle Blight Cercospora Leaf Spot, Pseudomonas Blight* Bottytis Blight	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn 5 Iris 6* Iry (English, Algerian) 1 Ixora Juniper Lantana Leyland Cypress* Lilac Lily, Easter 2 Linden*	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii Syringa spp. Lilium longiflorum Tilia spp.	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Canthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot Cercospora Needle Blight Cercospora Leaf Spot, Pseudomonas Blight* Bottytis Blight Anthracnose, Leaf Blight	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn ⁵ Iris ^{6*} Iry (English, Algerian) ¹ Ixora Juniper Lantana Leyland Cypress* Lilac Lily, Easter ² Linden* Loblolly Bay	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii Syringa spp. Lilium longiflorum Tilia spp. Gordonia lasianthus	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot Cercospora Needle Blight Cercospora Leaf Spot, Pseudomonas Blight* Botrytis Blight Anthracnose, Leaf Blight Anthracnose	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn ⁵ Iris ^{6*} Iry (English, Algerian) ¹ Ixora Juniper Lantana Leyland Cypress* Lilac Lily, Easter ² Linden* Loblolly Bay Loquat (ornamental)	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii Syringa spp. Lilium longiflorum Tilia spp. Gordonia lasianthus Eriobotrya japonica	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot Cercospora Needle Blight Cercospora Leaf Spot, Pseudomonas Blight* Botrytis Blight Anthracnose, Leaf Blight Anthracnose, Leaf Blight Colletotrichum spp., Entomosporium maculata	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn 5 Iris 6* Ivy (English, Algerian) 1 Ixora Juniper Lantana Leyland Cypress* Liilac Liily, Easter 2 Linden* Loblolly Bay Loquat (ornamental) Magnolia (Southern)	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii Syringa spp. Lillum longiflorum Tilia spp. Gordonia lasianthus Eriobotrya japonica Magnolia grandiflora	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot Cercospora Needle Blight Cercospora Leaf Spot, Pseudomonas Blight* Bottyiis Blight Anthracnose, Leaf Blight Anthracnose, Leaf Blight Anthracnose Colletotrichum spp., Entomosporium maculata Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn 5 Iris 6* Ivy (English, Algerian) 1 Ixora Juniper Lantana Leyland Cypress* Lilac Lily, Easter 2 Linden* Loblolly Bay Loquat (ornamental) Magnolia (Southern) Magnolia (Sweet Bay)	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii Syringa spp. Lilium longiflorum Tilia spp. Gordonia lasianthus Eriobotrya japonica Magnolia grandiflora Magnolia virginiana Magnolia soulangiana	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot Cercospora Needle Blight Cercospora Needle Blight Cercospora Leaf Spot, Pseudomonas Blight* Bottytis Blight Anthracnose, Leaf Blight Anthracnose, Leaf Blight Anthracnose Colletotrichum spp., Entomosporium maculata Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot Anthracnose	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn 5 Iris 6* Ivy (English, Algerian) 1 Ixora Juniper Lantana Leyland Cypress* Lilac Lily, Easter 2 Linden* Loblolly Bay Loquat (ornamental) Magnolia (Southern) Magnolia (Southern) Magnolia (Oriental) Mandevilla	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii Syringa spp. Lillum longiflorum Tilia spp. Gordonia lasianthus Eriobotrya japonica Magnolia virginiana Magnolia soulangiana Mandevilla spp.	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot Cercospora Needle Blight Cercospora Needle Blight Cercospora Leaf Spot, Pseudomonas Blight* Bottytis Blight Anthracnose, Leaf Blight Anthracnose, Leaf Blight Anthracnose Colletotrichum spp., Entomosporium maculata Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot Anthracnose Bacterial Leaf Spot Anthracnose	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn 5 Iris 6* Ivy (English, Algerian) 1 Ixora Juniper Lantana Leyland Cypress* Lilac Lily, Easter 2 Linden* Loblolly Bay Loquat (ornamental) Magnolia (Southern) Magnolia (Southern) Magnolia (Oriental) Mandevilla Maple*	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii Syringa spp. Lillum longiflorum Tilla spp. Gordonia lasianthus Eriobotrya japonica Magnolia grandiflora Magnolia virginiana Magnolia soulangiana Mandevilla spp. Acer spp.	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot Cercospora Needle Blight Cercospora Needle Blight Cercospora Leaf Spot, Pseudomonas Blight* Botrytis Blight Anthracnose, Leaf Blight Anthracnose, Leaf Blight Anthracnose Colletotrichum spp., Entomosporium maculata Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot Anthracnose Bacterial Leaf Spot Anthracnose Bacterial Leaf Spot Anthracnose Pseudomonas Leaf Blight	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn 5 Iris 6* Ivy (English, Algerian) 1 Ixora Juniper Lantana Leyland Cypress* Lilac Lily, Easter 2 Linden* Loblolly Bay Loquat (ornamental) Magnolia (Southern) Magnolia (Sweet Bay) Magnolia (Oriental) Mandevilla Maple* Marigold	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii Syringa spp. Lilium longiflorum Tilia spp. Gordonia lasianthus Eriobotrya japonica Magnolia grandiflora Magnolia virginiana Magnolia soulangiana Mandevilla spp. Acer spp. Tagetes spp.	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot Cercospora Needle Blight Cercospora Needle Blight Cercospora Leaf Spot, Pseudomonas Blight* Botrytis Blight Anthracnose, Leaf Blight Anthracnose, Leaf Blight Anthracnose Colletotrichum spp., Entomosporium maculata Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot Anthracnose Bacterial Leaf Spot Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn 5 Iris 6* Iry (English, Algerian) 1 Ixora Juniper Lantana Leyland Cypress* Lilac Lily, Easter 2 Linden* Loblolly Bay Loquat (ornamental) Magnolia (Southern) Magnolia (Sweet Bay) Magnolia (Oriental) Mandevilla Maple* Marigold Mountain-Ash*	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii Syringa spp. Lilium longiflorum Tilla spp. Gordonia lasianthus Eriobotrya japonica Magnolia grandiflora Magnolia virginiana Magnolia soulangiana Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp.	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot Cercospora Needle Blight Cercospora Needle Blight Cercospora Leaf Spot, Pseudomonas Blight* Botrytis Blight Anthracnose, Leaf Blight Anthracnose, Leaf Blight Anthracnose Collelotrichum spp., Entomosporium maculata Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot Anthracnose Bacterial Leaf Spot Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn 5 Iris 6* Ivy (English, Algerian) 1 Ixora Juniper Lantana Leyland Cypress* Lilac Lily, Easter 2 Linden* Loblolly Bay Loquat (ornamental) Magnolia (Southern) Magnolia (Southern) Magnolia (Sweet Bay) Magnolia (Oriental) Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted*	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii Syringa spp. Lilium longiflorum Tilia spp. Gordonia lasianthus Eriobotrya japonica Magnolia grandiflora Magnolia virginiana Magnolia soulangiana Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot Cercospora Needle Blight Cercospora Needle Blight Cercospora Leaf Spot, Pseudomonas Blight* Botrytis Blight Anthracnose, Leaf Blight Anthracnose, Leaf Blight Anthracnose, Entomosporium maculata Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot Anthracnose Bacterial Leaf Spot Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn 5 Iris 6* Ivy (English, Algerian) 1 Ixora Juniper Lantana Leyland Cypress* Lilac Lily, Easter 2 Linden* Loblolly Bay Loquat (ornamental) Magnolia (Southern) Magnolia (Sweet Bay) Magnolia (Oriental) Mandevilla Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted* Mulberry, Weeping	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii Syringa spp. Lilium longiflorum Tilla spp. Gordonia lasianthus Eriobotrya japonica Magnolia grandiflora Magnolia virginiana Magnolia virginiana Magnolia soulangiana Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis Morus alba	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot Cercospora Needle Blight Cercospora Needle Blight Cercospora Leaf Spot, Pseudomonas Blight* Botrytis Blight Anthracnose, Leaf Blight Anthracnose, Leaf Blight Anthracnose Colletotrichum spp., Entomosporium maculata Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot Anthracnose Bacterial Leaf Spot Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn 5 Iris 6* Iry (English, Algerian) 1 Ixora Juniper Lantana Leyland Cypress* Lilac Lily, Easter 2 Linden* Loblolly Bay Loquat (ornamental) Magnolia (Southern) Magnolia (Southern) Magnolia (Oriental) Mandevilla Mandevilla Mandevilla Mandel* Marigold Mountain-Ash* Mulberry, Contorted* Mulberry, Weeping Narcissus*	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii Syringa spp. Lilium longiflorum Tilia spp. Gordonia lasianthus Eriobotrya japonica Magnolia virginiana Magnolia virginiana Magnolia soulangiana Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis Morus alba Narcissus spp.	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot Cercospora Needle Blight Cercospora Needle Blight Cercospora Leaf Spot, Pseudomonas Blight* Botrytis Blight Anthracnose, Leaf Blight Anthracnose, Leaf Blight Anthracnose Colletotrichum spp., Entomosporium maculata Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot Anthracnose Bacterial Leaf Spot Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn 5 Iris 6* Ivy (English, Algerian) 1 Ixora Juniper Lantana Leyland Cypress* Lilac Lily, Easter 2 Linden* Loblolly Bay Loquat (ornamental) Magnolia (Southern) Magnolia (Southern) Magnolia (Oriental) Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted* Mulberry, Weeping Narcissus* Nephthytis*	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii Syringa spp. Lilium longiflorum Tilia spp. Gordonia lasianthus Eriobotrya japonica Magnolia grandiflora Magnolia virginiana Magnolia soulangiana Mandeivilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis Morus alba Narcissus spp. Syngonium podophyllum	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot Cercospora Needle Blight Cercospora Needle Blight Cercospora Leaf Spot, Pseudomonas Blight* Botrytis Blight Anthracnose, Leaf Blight Anthracnose, Leaf Blight Anthracnose Colletotrichum spp., Entomosporium maculata Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot Anthracnose Bacterial Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot	
Honeysuckle, Tatarian* Impatiens Indian Hawthom 5 Iris 6** Ivy (English, Algerian) 1 Ixora Juniper Lantana Leyland Cypress* Lilac Lily, Easter 2 Linden* Loblolly Bay Loquat (ornamental) Magnolia (Southern) Magnolia (Southern) Magnolia (Southern) Magnolia (Sweet Bay) Magnolia (Oriental) Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted* Mulberry, Weeping Narcissus* Nephthytis*	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii Syringa spp. Lillim longiflorum Tilia spp. Gordonia lasianthus Eriobotrya japonica Magnolia grandiflora Magnolia grandiflora Magnolia soulangiana Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis Morus alba Narcissus spp. Syngonium podophyllum Quercus spp.	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot Cercospora Needle Blight Cercospora Needle Blight Cercospora Leaf Spot, Pseudomonas Blight* Botrytis Blight Anthracnose, Leaf Blight Anthracnose, Leaf Blight Anthracnose Colletotrichum spp., Entomosporium maculata Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot Anthracnose Bacterial Leaf Spot, Anthracnose, Bacterial Leaf Spot Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot Leaf Spot Leaf Spot Leaf Spot Leaf Spot	
Honeysuckle, Tatarian* Impatiens Indian Hawthom 5 Iris 6* Ivy (English, Algerian) 1 Ixora Juniper Lantana Leyland Cypress* Lilac Lily, Easter 2 Linden* Loblolly Bay Loquat (ornamental) Magnolia (Southern) Magnolia (Southern) Magnolia (Sweet Bay) Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted* Mulberry, Weeping Narcissus* Nephthytis* Oak* Oak, Laurel	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii Syringa spp. Lillim longiflorum Tilia spp. Gordonia lasianthus Eriobotrya japonica Magnolia grandiflora Magnolia virginiana Magnolia virginiana Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis Morus alba Narcissus spp. Syngonium podophyllum Quercus spp. Quercus laurifolia	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot Cercospora Needle Blight Cercospora Needle Blight Cercospora Leaf Spot, Pseudomonas Blight* Botrytis Blight Anthracnose, Leaf Blight Anthracnose, Leaf Blight Anthracnose Colletotrichum spp., Entomosporium maculata Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot Anthracnose Bacterial Leaf Spot Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot Leaf Spot Leaf Spot Leaf Spot Leaf Spot Leaf Spot Leaf Spots Algal Leaf Spot Leaf Spots Algal Leaf Spot Leaf Spots Algal Leaf Spot (Cephaleuros virescens)	
Honeysuckle, Tatarian* Impatiens Indian Hawthom 5 Iris 6* Iry (English, Algerian) 1 Ixora Juniper Lantana Leyland Cypress* Lilac Lily, Easter 2 Linden* Loblolly Bay Loquat (ornamental) Magnolia (Southern) Magnolia (Southern) Magnolia (Sweet Bay) Magnolia (Oriental) Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted* Mulberry, Weeping Narcissus* Nephthytis* Oak* Oak, Laurel Oleander	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii Syringa spp. Lilium longiflorum Tilia spp. Gordonia lasianthus Eriobotrya japonica Magnolia grandiflora Magnolia virginiana Magnolia soulangiana Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis Morus alba Narcissus spp. Syrgonium podophyllum Quercus spp. Quercus laurifolia Nerium oleander	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot Cercospora Needle Blight Cercospora Needle Blight Cercospora Leaf Spot, Pseudomonas Blight* Botrytis Blight Anthracnose, Leaf Blight Anthracnose, Leaf Blight Anthracnose, Entomosporium maculata Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot Anthracnose Bacterial Leaf Spot Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot Bacterial Leaf Spot Leaf Blight Bacterial Leaf Spot Leaf Spots Algal Leaf Spot Leaf Spots Algal Leaf Spot (Cephaleuros virescens) Bacterial Leaf Spot, Fungal Leaf Spot	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn 5 Iris 6* Iry (English, Algerian) 1 Ixora Juniper Lantana Leyland Cypress* Lilac Lily, Easter 2 Linden* Loblolly Bay Loquat (ornamental) Magnolia (Southern) Magnolia (Southern) Magnolia (Oriental) Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted* Mulberry, Weeping Narcissus* Nephthytis* Oak* Oak, Laurel Oleander Oregon Grape Holly*	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii Syringa spp. Lilium longiflorum Tilia spp. Gordonia lasianthus Eriobotrya japonica Magnolia grandiflora Magnolia grandiflora Magnolia soulangiana Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis Morus alba Narcissus spp. Syngonium podophyllum Quercus spp. Quercus laurifolia Nerium oleander Mahonia aquifolium	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot Cercospora Needle Blight Cercospora Needle Blight Cercospora Leaf Spot, Pseudomonas Blight* Botrytis Blight Anthracnose, Leaf Blight Anthracnose, Leaf Blight Anthracnose Collectrichum spp., Entomosporium maculata Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot Anthracnose Bacterial Leaf Spot Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot Leaf Blight Bacterial Leaf Spot Leaf Spots Algal Leaf Spot, Fungal Leaf Spot Leaf Spots	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn 5 Iris 6* Iry (English, Algerian) 1 Ixora Juniper Lantana Leyland Cypress* Lilac Lily, Easter 2 Linden* Loblolly Bay Loquat (ornamental) Magnolia (Southern) Magnolia (Southern) Magnolia (Southern) Magnolia (Oriental) Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted* Mulberry, Weeping Narcissus* Nephthytis* Oak* Oak, Laurel Oleander Oregon Grape Holly* Pachysandra	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii Syringa spp. Lilium longiflorum Tilia spp. Gordonia lasianthus Eriobotrya japonica Magnolia grandiflora Magnolia virginiana Magnolia soulangiana Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis Morus alba Narcissus spp. Syrgonium podophyllum Quercus spp. Quercus laurifolia Nerium oleander	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot Cercospora Needle Blight Cercospora Needle Blight Cercospora Leaf Spot, Pseudomonas Blight* Botrytis Blight Anthracnose, Leaf Blight Anthracnose, Leaf Blight Anthracnose, Entomosporium maculata Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot Anthracnose Bacterial Leaf Spot Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot Bacterial Leaf Spot Leaf Blight Bacterial Leaf Spot Leaf Spots Algal Leaf Spot Leaf Spots Algal Leaf Spot (Cephaleuros virescens) Bacterial Leaf Spot, Fungal Leaf Spot	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn 5 Iris 6* Iry (English, Algerian) 1 Ixora Juniper Lantana Leyland Cypress* Lilac Lily, Easter 2 Linden* Loblolly Bay Loquat (ornamental) Magnolia (Southern) Magnolia (Southern) Magnolia (Oriental) Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted* Mulberry, Weeping Narcissus* Nephthytis* Oak* Oak, Laurel Oleander Oregon Grape Holly*	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii Syringa spp. Lilium longiflorum Tilia spp. Gordonia lasianthus Eriobotrya japonica Magnolia grandiflora Magnolia grandiflora Magnolia soulangiana Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis Morus alba Narcissus spp. Syngonium podophyllum Quercus spp. Quercus laurifolia Nerium oleander Mahonia aquifolium	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot Cercospora Needle Blight Cercospora Needle Blight Cercospora Leaf Spot, Pseudomonas Blight* Botrytis Blight Anthracnose, Leaf Blight Anthracnose, Leaf Blight Anthracnose Collectorichum spp., Entomosporium maculata Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot Anthracnose Bacterial Leaf Spot Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot Bacterial Leaf Spot Leaf Blight Bacterial Leaf Spot Leaf Spots Algal Leaf Spot (Cephaleuros virescens) Bacterial Leaf Spot, Fungal Leaf Spot Leaf Spots	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn 5 Iris 6* Iry (English, Algerian) 1 Ixora Juniper Lantana Leyland Cypress* Lilac Lily, Easter 2 Linden* Loblolly Bay Loquat (ornamental) Magnolia (Southern) Magnolia (Southern) Magnolia (Southern) Magnolia (Oriental) Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted* Mulberry, Weeping Narcissus* Nephthytis* Oak* Oak, Laurel Oleander Oregon Grape Holly* Pachysandra Palm, Date (ornamental) Palm, European Fan	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii Syringa spp. Lilium longiflorum Tilia spp. Gordonia lasianthus Eriobotrya japonica Magnolia grandiflora Magnolia virginiana Magnolia soulangiana Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis Morus alba Narcissus spp. Syngonium podophyllum Quercus spp. Quercus faurifolia Nerium oleander Mahonia aquifolium Pachysandra procumbens	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Anthracnose, Phomopsis Twig Dieback* Bacterial Leaf Spot Cercospora Needle Blight Cercospora Needle Blight Cercospora Leaf Spot, Pseudomonas Blight* Bottytis Blight Anthracnose, Leaf Blight Anthracnose, Leaf Blight Anthracnose Collectorichum spp., Entomosporium maculata Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot Anthracnose Bacterial Leaf Spot Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Leaf Blight Bacterial Leaf Spot Leaf Spots Algal Leaf Spot, Fungal Leaf Spot Leaf Spots Volutella Leaf Blight	
Honeysuckle, Tatarian* Impatiens Indian Hawthorn 5 Iris 6* Iry (English, Algerian) 1 Ixora Juniper Lantana Leyland Cypress* Lilac Lily, Easter 2 Linden* Loblolly Bay Loquat (ornamental) Magnolia (Southern) Magnolia (Southern) Magnolia (Southern) Magnolia (Oriental) Magnolia (Oriental) Magnolia (Oriental) Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted* Mulberry, Weeping Narcissus* Nephthytis* Oak* Oak, Laurel Oleander Oregon Grape Holly* Pachysandra	Gleditsia triacanthos Lonicera tatarica Impatiens wallerana Raphiolepis indica Iris spp. Hedera helix, H. canariensis Ixora coccinea Juniperus spp. Lantana camera X Cupressocyparis leylandii Syringa spp. Lilium longiflorum Tilia spp. Gordonia lasianthus Eriobotrya japonica Magnolia grandiflora Magnolia virginiana Magnolia virginiana Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis Morus alba Narcissus spp. Syngonium podophyllum Quercus spp. Quercus laurifolia Nerium oleander Mahonia aquifolium Pachysandra procumbens Phoenix canariensis	Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Bacterial Leaf Spot Anthracnose, Entomosporium Leaf Spot Bacterial Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Xanthomonas Leaf Spot Cercospora Needle Blight Cercospora Needle Blight Cercospora Leaf Spot, Pseudomonas Blight* Botrytis Blight Anthracnose, Leaf Blight Anthracnose, Leaf Blight Anthracnose, Leaf Blight Anthracnose Collectorichum spp., Entomosporium maculata Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot Leaf Blight Bacterial Leaf Spot Leaf Spots Algal Leaf Spot (Cephaleuros virescens) Bacterial Leaf Spots Volutella Leaf Blight Pestalotia Leaf Blight Pestalotia Leaf Spot	

ORNAMENTALS (continued) Jse Kocide 2000-O for control of bacterial and fungal diseases of foliage, flowers and stems on ornamentals in greenhouses, shade houses, outdoor nurseries and outdoor landscape

Use Kocide 2000-O for control of bacterial and fungal diseases of foliage, flowers and stems on ornamentals in greenhouses, shade houses, outdoor nurseries and outdoor landscape plantings.
For ornamental crops in dormancy, apply as a thorough cover spray at rates ranging from 0.75 to 2 pounds per acre of Kocide 2000-O. When new growth is present, apply as a thorough cover spray at rates ranging from 0.75 to 2 pounds per acre of Kocide 2000-O. One level tablespoon of Kocide 2000-O per 1,000 square feet is equivalent to 1.5 pounds of product per acre. Begin application at first sign of disease and repeat at 7- to 14-day intervals if needed; use the higher rates and shorter spray intervals during periods of frequent rains or when severe disease conditions persist. Maximum seasonal rate per acre is 57.1 lbs.
Kocide 2000-O may be used alone or in combination with other fungicides registered for use on ornamentals as a maintenance spray. Use in accordance with the most restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

Notice to User: Plant sensitivities to Kocide 2000-O have been found to be acceptable for the specific genera and species listed on this label under the conditions tested. However, phytotoxicity may occur. Due to the large number of species and varieties of ornamental and nursery plants, and the wide range of growing conditions, it is impossible to test every one for sensitivity to Kocide 2000-O. Neither the manufacturer nor seller has determined whether or not Kocide 2000-O can be safely used on ornamental or nursery plants not itseld on this label. The user should determine if Kocide 2000-O per present and sensitivity to commercial use. In a small area, apply the recommended rates to the plants in question, i.e., bedding plants, foliage, etc., and observe for 7- to 10-days for symptoms of phytotoxicity prior to commercial use.

NOTE: This product may be reactive

Crop	Scientific Name	Disease	
Palm, Washingtonia	Washingtonia robusta	Pestalotia Leaf Spot	
Peach (Flowering) (ornamental) 3*	Prunus spp.	Bacterial Blast, Brown Rot, Fire Blight	
Pear (Flowering) (ornamental)	Pyrus calleryana	Fire Blight, Leaf Spots	
Pentas (Egyptian Star)	Pentas spp.	Bacterial Leaf Spot (Pseudomonas spp.*, Xanthomonas spp.)	
Peony	Paeonia spp.	Botrytis Blight	
Periwinkle	Catharanthus roseus, Vinca spp.	Phomopsis Stem Blight	
Philodendron	Philodendron selloum	Bacterial Leaf Spot	
Phlox	Phlox spp.	Alternaria Leaf Spot	
Photinia (Red Tip)	Photinia x fraseri, P. glabra	Anthracnose, Entomosporium Leaf Spot	
Pine*	Pinus spp.	Needlecasts	
Pistachio (ornamental)	Pistacia chinensis	Anthracnose	
Plantain Lily 6	Hosta spp.	Bacterial Leaf Spot	
Plum (Flowering) (ornamental) 3*	Prunus spp.	Bacterial Blast, Brown Rot, Fire Blight	
Pothos*	Scindapsus spp.	Bacterial Leaf Spot	
Powder Puff Plant	Calliandra spp.	Bacterial Leaf Spot	
Pyracantha	Pyracantha spp.	Fire Blight, Scab	
Rhododendron	Rhododendron spp.	Alternaria Flower Spot	
Rose 1	Rosa spp.	Black Spot, Powdery Mildew	
Snapdragon	Antirrhinum majus	Anthracnose, Dieback, Downy Mildew	
Spathe Flower*	Spathiphyllum spp.	Bacterial Leaf Spot	
Spirea*	Spiraea spp.	Fire Blight	
Spruce*	Picea spp.	Needlecasts	
Sycamore	Platanus spp.	Anthracnose, Leaf Spots*	
Tulip	Tulipa spp.	Anthracnose, Botrytis Blight	
Umbrella Tree*	Schefflera spp.	Bacterial Leaf Spot	
Verbena	Verbena spp.	Xanthomonas Leaf Spot	
Viburnum	Viburnum odoratissimum, V. plicatum, V. suspensum	Anthracnose	
Viola (Pansy, Violet)	Viola spp.	Downy Mildew	
Willow	Salix spp.	Anthracnose	
Yew*	Taxus spp.	Needle Blight	
Yucca (Adam's Needle)	Yucca spp.	Cercospora Leaf Spot, Septoria Leaf Spot	
Zinnia*	Zinnia spp.	Leaf Spots	

Discoloration of foliage and/or blooms has been noted on some varieties. To prevent residues on commercial plants, do not spray immediately before selling season.

Apply Kocide 2000-0 at 2.25 to 3,75 pounds per acre. Maximum seasonal rate per acre is 214 lbs. Do not apply any additional copper pesticide to this land for 36 months. Apply Kocide 2000-O at 2.25 to 3.75 pounds per acr
 Apply dormant through bloom only.
 Hibiscus - Do not apply to plants in flower.
 For Indian Hawthom use 1.5 to 3.0 pounds per acre.
 Some cultivars may be sensitive to Kocide 2000-O.
 Not registered for use in California.

NOTE: Phytotoxicity may depend on varietal differences. If unfamiliar with the use of Kocide 2000-O, apply the recommended rate to a few plants and observe after 7 to 10 days for symptoms of phytotoxicity.

Control of Ball Moss*, Spanish Moss* and Lichens* on Ornamental and Shade Trees: Apply Kocide 2000-O in early spring when the trees are dormant. Apply 4.5 to 6 pounds of Kocide 2000-O in 100 gallons of water, using 1 1/2 gallons of spray per foot of tree height. Be sure to thoroughly wet ball moss tufts, Spanish moss or lichens. The addition of a non-ionic surfactant will improve control. A second application may be required after 12 months.

NOTE: Kocide 2000-O may be injurious to some ornamental plants growing beneath the trees. This product may be reactive on masonry and metal surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.

Cold Storage Protection for Dormant Rootstock*: To protect bars-roof unsery trees from Phytophthora Crown Rot and Botrytis, use 2 to 3 pounds of Kocide 2000-O per 100 gallons of water. Apply as a dip or spray to the roots and lower stems of dormant rootstock prior to placing in cold storage. Do not apply to rootstock less than 2 years old.

*Not registered for use in California.

GENERAL CHEMIGATION INSTRUCTIONS

Apply this product only through one or more of the following types of systems: sprinkler, including center pivot, lateral move, traveler, big gun, or plastic pipe solid set system(s). Do not apply this product through any other type of irrigation systems. In California, do not apply in systems which contain aluminum parts or components.

Crop injury, lack of effectiveness or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Shut off injection equipment after treatment and continue to operate irrigation system until Kocide 2000-O has been cleared from the last sprinkler head.

CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS
Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into the reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

NOTE: It must be determined if proper application equipment is availabl

recommended. Kocide 2000-O should be added through a traveling irrigation system continuously or at the last 30 minutes of solid set irrigation systems. Shut off injection equipment after treatment and continue to operate irrigation system until Kocide 2000-O has been cleared from the last sprinkler head.

SPRINKLER CHEMIGATION
The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The postificial principle principle principle principle are contain a functional customatic quick closing check valve to prevent the flow of fluid back toward the injection pump.

In e system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must also contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The epsticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interiock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fittled with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

NOTE: It must be determined if proper application equipment is available and if waste associated with its use can be properly handled. Agricultural chemicals are often reactive with the materials used in the construction of application equipment, such as aluminum, rubber and some synthetic materials. This factor should be taken into consideration when selecting proper application equipment. It is necessary that all application equipment be thoroughly flushed with clean water after each day's use.

When mixing, fill the nurse tank half full with water. Add Kocide 2000-O slowly to tank while hydraulic or mechanical agitation is operating and continue filling with

recommended.
Kocide 2000-O should be added through a traveling irrigation system continuously or at the last 30 minutes of solid set irrigation systems. Shut off injection equipment after treatment and continue to operate irrigation system until Kocide 2000-O has been cleared from the last sprinkler head.

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet size:

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet size:

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind speed:

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition (approximately 3 to 10 mph), and there are not sensitive areas within 250 feet downwind.

Temperature Inversions:

If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Other State and Local Requirements:

Applicators must follow all state and local pesticide drift requirements regarding application of copper compounds. Where states have stringent regulations, they must be observed.

Equipment:

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Additional requirements for aerial applications:

- The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

- Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.

- When applications are made with a crosswind, the swath must be displaced downwind. The applicator must compensate for this displacement at the up and downwind edge of the applications are made with a crosswind, the swath must be displaced downwind. The applicator must compensate for this displacement at t

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest cannot be disposed of by use according to laurel instructions, without pour other research office for guidance.

EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Completely empty paper or plastic bag, fiber sack or drum liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer for recycling if available or dispose of empty paper or plastic bag, fiber sack or fiber drum and liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact ChemTel

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States. "Curtec" is a registered trademark of Bei Incorporated "Tre-Hold" is a registered trademark of Amvac Chemical Corporation.

LIMITATION OF WARRANTY AND LIABILITY

LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read this Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of Certis. These risks can cause: ineffectiveness of the product, crop injury, or injury to non-target crops or plants. WHEN YOU BUY OR USE TIPS PRODUCT, YOU ACREE TO ACCEPT THESE RISKS.

Certis warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, CERTIS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL CERTIS OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE INSULTIVE OF CERTIS OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY, WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF CERTIS OR SELLER. THE REPLACEMENT OF THE PRODUCT.

To the extent consistent with applic

Certis LLC, 9145 Guilford Road, Suite 175, Columbia, MD 21046 All rights reserved

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer for recycling if available, or dispose of empty bag in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact ChemTel at **1-800-255-3924**, day or night.



FUNGICIDE/BACTERICIDE+

Dry Flowable

ACTIVE INGREDIENT:	BY WT.
Copper Hydroxide* (CAS No. 20427-59-2)	53.8%
INERT INGREDIENTS:	46.2%
TOTAL:	100.0%

⁺Non-public health bactericide *Metallic Copper Equivalent 35%



FOR ORGANIC PRODUCTION

KEEP OUT OF REACH OF CHILDREN WARNING

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

EPA Reg. No. 91411-10-70051 EPA Est. No. 91411-TX-1

Manufactured For: Certis USA LLC 9145 Guilford Road, Suite 175 Columbia, MD 21045

NET: 20 LBS.

(nonrefillable container)

FIRST AID				
IF IN EYES:	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. 			
IF SWALLOWED:	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person. 			
IF INHALED:	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. 			

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for You may also contact **1-800-255-3924** for emergency medical treatment information.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate use of gastric lavage.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS **WARNING - AVISO**

Causes substantial but temporary eye injury. Harmful if swallowed, absorbed through the skin or inhaled. Do not get in eyes, on skin, or on clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the bathroom. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders, applicators, and other handlers must wear: Long-sleeved shirt and long pants

- Chemical-resistant gloves made of any waterproof material, such as polyvinyl chloride, nitrile

- Shoes plus socks - Protective eyewear such as safety glasses, goggles, or face shield
Discard clothing and other absorbent materials that have been drenched or heavily contaminated

with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

PHYSICAL AND CHEMICAL HAZARDS

Do not mix or allow to come in contact with oxidizing agents or flame retardants. Hazardous chemical reaction may occur.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

ENVIROMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may contaminate water through runoff. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash-waters or

Drift and runoff may be hazardous to aquatic organisms in waters adjacent to treated areas.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries

KOCIDE® is registered trademark of Kocide LLC

ESL20171011

A20170170

Made In U.S.A.

5H4/Z14/S/15

FUNGICIDE/ BACTERICIDE .	
Dry Flowable By W Active Ingredient: By W Copper Hydroxide* (CAS No. 20427-59-2) 9 Inert Ingredients: 2 TOTAL 11	46.2%

† Non-public health bactericide * Metallic Copper Equivalent 35%

KEEP OUT OF REACH OF CHILDREN WARNING
Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

EPA Reg. No. 91411-10-70051

Kocide

CERTIS

FOR ORGANIC PRODUCTION

Columbia, MD 210 20180503	40 L S T E D For Organic Use
ESL20171011	FOR ORGANIC PRODUCTION
	FIRST AID
IF IN EYES:	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF INHALED:	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
	HOT LINE NUMBER

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-255-3924 for emergency medical treatment information. NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate use of gastric

PRECAUTIONARY STATEMENTS **HAZARDS TO HUMANS & DOMESTIC ANIMALS** WARNING - AVISO

Causes substantial but temporary eye injury. Harmful if swallowed, absorbed through the skin or inhaled. Do not get in eyes, on skin, or on clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the bathroom. Remove and wash contaminated clothing before

PERSONAL PROTECTIVE EQUIPMENT (PPE)
Mixers, loaders, applicators and other handlers must wear:

• Long-sleeved shirt and long pants
• Chemical-resistant gloves made of any waterproof material, such as polyvinyl chloride, nitrile rubber, or butyl rubber

 Shoes plus socks
 Protective eyewear such as safety glasses, goggles, or face shield
 Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. PHYSICAL AND CHEMICAL HAZARDS

Do not mix or allow to come in contact with oxidizing agents or flame retardants. Haz-

ardous chemical reaction may occur.

USER SAFETY RECOMMENDATIONS
USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the tollet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

through runoff.

This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash-waters or rinsate.

Drift and runoff may be hazardous to aquatic organisms in waters adjacent to treated

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Les provisions for presurder regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours without required PPE.

(REI) of 48 hours without required PPE.
PPE required for early entry to treated areas that is permitted under the Worker Protection
Standard and that involves contact with anything that has been treated, such as plants,
soil or water, is:
• Coveralls
• Chemical-resistant gloves made of any waterproof material, such as polyvinyl chloride, nitrile rubber, or butyl rubber
• Shoes plus socks
• Protective eyewear such as safety glasses, goggles, or face shield

AGRICULTURAL USE REQUIREMENTS (continued)

For Greenhouse Uses ONLY:

The 48-hour restricted entry interval (REI) may be reduced to 24-hour REI, provided that the following conditions are met:

For at least seven days following the application of copper-containing products in

For at least seven days following the application of copper-containing products in greenhouses:

at least one container or station designed specifically for flushing eyes is available in operating condition with the WPS- required decontamination supplies for workers entering the area treated with copper-containing products,

workers are informed orally, in a manner they can understand:
that residues in the treated area may be highly irritating to the eyes,
that they should take precautions, such as refraining from rubbing their eyes, to keep the residues out of their eyes,
that if they do get residues in their eyes, they should immediately flush their eyes with the eye flush container or eye flush station that is located with the decontamination supplies, and

nation supplies, and how to operate the eye flush container or eye flush station.

NON-AGRICULTURAL USE REQUIREMENTS The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides 40 CFR Part 170. The WPS applies when this product is used to produce agricultural plants on farms, forests,

Do not enter or allow others to enter until sprays have dried. GENERAL INSTRUCTIONS

GENERAL INSTRUCTIONS

Kocide 2000-O may be applied as an aerial, ground dilute or ground concentrate spray unless specifically directed otherwise in the specific crop use directions.

The per acre use rate of Kocide 2000-O is applicable for both dilute and concentrate spraying. Depending upon the equipment used and the specific crop, the spray volume applied per acre will differ. Refer to Minimum Recommended Spray Volume Table. Complete spray coverage is essential to assure optimum performance from Kocide 2000-O. When treating by aerial application or with low volume application or with Kocyde 2000-O. On the volume and the volume application or with Kocyde 2000-O. On the lor specific rates and timing of application by crop.

Consult the Kocide 2000-O label for specific rates and timing of application by crop. Where application rates and intervals are provided in a range (e.g. 4 to 12 ounds and 7 to 10 days), the higher rates and shorter spray intervals are recommended when rainfall is heavy and/or disease pressure is high. Use the higher rates for large mature tree crops.

SPECIAL PRECAUTIONS
The Pre-Harvest Interval (PHI) for Kocide 2000-O is 0-days unless noted.

If Kocide 2000-O is applied in a spray solution having a pH of less than 6.5, phytotoxicity

may occur.

Do not tank mix Kocide 2000-O with fungicides containing the active ingredient aluminum tris (O-ethyl phosphonate) for use on any registered crops unless appropriate precautions have been taken to buffer the spray solution because severe phytotoxicity may result. Use in accordance with the most restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing. This product may be reactive on masonry and metal surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc. Environmental conditions such as extended periods of wet weather, acid rain, etc. which alter the pH of the leaf surface may affect the performance of Kocide 2000-O resulting in possible phytotoxicity or loss of effectiveness.

Agricultural chemicals may perform in an unpredictable manner when tank mixed, especially where several products are involved. Reduced effect on pests or crop injury may occur. Unless recommended on this label or by a statellocal expert, it is advisable to test for compatibility and potential crop injury prior to commercial use of a new tank mix; otherwise, tank mixing should not be undertaken.

It must be determined if proper application equipment is available and if waste associated with its use can be properly handled. Agricultural chemicals are often reactive with the materials used in the construction of application equipment, such as aluminum, rubber and some synthetic materials. This factor should be taken into consideration when selecting proper application equipment after each day's use.

Apply this product only through one or more of the following types of systems: sprinkler, including center pivot, lateral move, traveler, big gun, or plastic pipe solid set system(s). Do not apply this product through any other type of irrigation system. In California, do not apply in systems which contain aluminum parts or components. may occur.

• Do not tank mix Kocide 2000-O with fungicides containing the active ingredient aluminum

When mixing, fill the spray tank one-half full with water. Add Kocide 2000-O slowly to tank

When mixing, fill the spray tank one-half full with water. Add Kocide 2000-O slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. Spreaders, stickers, insecticides, nutrients, etc. should be added last. If compatibility is in question, use the Compatibility Jar Test before mixing a whole tank or contact your chemical supplier. Observe all precautions and limitations on the labels of all products used in mixtures.
 It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

CROP CLASSIFICATION Grapefruit, Kumquat, Lemon, Lime, Orange, Pummelo, Tangelo, and Douglas Fir, Fir*, Juniper, Leyland Cypress*, Pine*, and Spruce*.
Alfalfa, Barley, Com*, Oats, Peanut, Potato, Sugar Beet, and Wheat.
Blackberry, Blueberry*, Cranberry, Currant, Gooseberry, Raspberry,
and Strawberry. SMALL FRUITS: TREE CROPS:

and Strawberry.

Almond, Apple, Apricot, Avocado, Banana/Plantain, Cacao, Cherry,
Coffee, Filbert, Mango, Nectarine, Olive, Peach, Pear, Pecan, Pistachio, Plum, Prune, Quince, and Walnut.
Bean, Beet, Beet Greens, Broccoli, Brussels Sprout, Cabbage, Chinese Cabbage, Cantaloupe, Carrot, Cauliflower, Celeriac, Celery,
Cumber, Egghant, Greens (Collard, Mustard and Turnip), Honeydew, Kale, Kohlrabi, Muskmelon, Okra*, Onion/Garlio/Leek, Pea,
Pepper, Pumpkin, Spinach, Squash, Tomato, Watercress, and Watermelon. VEGETABLES:

VINES: Grape, Hops, and Kiwi.

MISCELLANEOUS: Atemoya, Carambola, Chives, Dill, Ginseng, Guava, Litchi,
Macadamia, Mamey Sapote, Papaya, Parsley, Passion Fruit, Pecan,
Sugar Apple, and Sycamore.

GREENHOUSE AND SHADEHOUSE CROPS: Kocide 2000-0 may be used in greenhouses and shadehouses to control diseases on any crop on this label where physiology allows greenhouse or shadehouse culture. While specific directions are presented for Cucumber, Eggplant, Pepper and Tomato; general use may occur for any crop

on this label where physiology allows greenhouse or shadehouse culture. *Not registered for use in California. Minimum Recommended Spray Volume (Gallons Per Acre)

	•	,
When Applying	Kocide	2000-O

		G	round
	Aerial	Dilute	Concentrate
trus	10	800	100**
onifers	10	100	30
eld Crops	3	20	3
mall Fruits	5	150	50
ee Crops	10	400	50
egetables	3	20	3
nes	5	150	50
iscellaneous	10	150	50

**Pesticide application equipment such as "Curtec" or other similar sprayers which are capable of obtaining thorough coverage at low volumes may be used at as low as 20 gallons per acre of spray volume.

The following specific instructions are based on general application procedures. The recommendations of the State Agricultural Extension Service should be closely followed as to timing, frequency and number of sprays per season. FROST INJURY PROTECTION
BACTERIAL ICE NUCLEATION INHIBITOR

Application of Kocide 2000-O made to all crops listed on this label at rates and stages of growth indicated on this label, at least 24 hours prior to anticipated frost conditions, will afford control of ice nucleating bacteria (*Pseudomonas syringae*, *Erwinia herbicola*, and *Pseudomonas fluorescens*) and may therefore provide some protection against light frost.

Not recommended for those geographical areas where weather conditions favor sever frost. **CITRUS**

Kocide 2000-O may be mixed with dry foliar nutritionals (micronutrients) to create "Shot Bag" mixes to meet the various nutritional requirements of citrus and provide disease protection as described on this label. Kocide 2000-O per acre rates in these mixes must not exceed the maximum recommended labeled rates for disease control. Adding foliar nutritionals or other products to spray mixtures containing Kocide 2000-O and applying to citrus during the post bloom period when young fruit are present may result in sorav burn.

result in spray burn.			
Disease	Application Rate/Acre	Maximum Annual Rate/Acre	Use Instructions
Algal Spot, Melanose, Scab	3-9 lbs.	36 lbs.	Apply as pre-bloom and post-bloom sprays. Use the higher rates when conditions favor disease. Minimum retreatment interval is 7 days.
Greasy Spot, Pink Pitting	1.5-4.5 lbs.	36 lbs.	Apply in summer on expanded new flush. Repeat on subsequent flushes where disease pressure is severe. Use the higher rates when conditions favor disease. Minimum retreatment interval is 7 days.
Alternaria Brown Spot	3-6 lbs.	36 lbs.	On susceptible varieties apply when the first spring flush appears and each flush thereafter. Application to fruit should start after two thirds of the petals have fallen and be repeated on a 7- to 21-day schedule if needed. Use the higher rates when conditions favor disease.
Phytophthora Brown Rot, Septoria Spot	3-6 lbs.	36 lbs.	Begin application in fall before or just after the first rain and continue if needed. For Brown Rot only, apply to skirts of trees to a height of at least 4 feet. For control of Septoria Spot or where fruit have already been infected with Brown Rot, apply to entire tree. Apply also to bare ground one foot beyond skirt. Use the higher rates when conditions favor disease. Minimum retreatment interval is 7 days. NOTE: In California, in areas subject to copper injury, add 1/3 to 1 pound of high quality lime per pound of Kocide 2000-O.
Phytophthora Foot Rot	0.75 lb.	36 lbs.	Mix with 1 quart of water, "Tre-Hold" or latex paint. Paint trunks of trees from the soil surface to the lowest scaffold limbs. Apply in May prior to summer rains and/or in the fall prior to wrapping trees for freeze protection. Treatment serves as protection for up to 1 year, but does not cure existing infections. NOTE: Areas where microjet or low volume irrigation hit the tree trunk may require retreatment due to wash off.
Citrus Canker (suppression)	2-4 lbs.	36 lbs.	Spray flushes 7 to 14 days after shoots begin to grow. Young fruit may require an additional application. Number and timing of applications will be dependent upon disease pressure. Under heavy pressure, each flush of new growth should be sprayed. Minimum retreatment interval is 7 days.
Black Spot*	2-4 lbs.	36 lbs.	Begin treatment prior to or when disease first appears and repeat every 7 to 21 days if needed. Use the higher rates and shorter spray intervals when conditions favor disease. Minimum retreatment interval is 7 days.

NOTE: Phytotoxicity may occur on young tender flush when Kocide 2000-O is applied to citrus seedlings grown in greenhouses or shadehouses.

CITRUS - Field Nursery Grown

apply 3 to 6 pounds of Kocide 2000-O per acre. Apply Kocide 2000-O at 28 To control Melanose, Scab, Pink Pitting, Greasy Spot, Brown Rot and for

			FIELD CRO	PS
Crop	Disease	Application Rate/Acre	Maximum Annual Rate/Acre	Use Instructions
Alfalfa	Cercospora Leaf Spot, Leptosphaerulina Leaf Spot	1.5 lbs.	3.2 lbs.	Apply 10 to 14 days before each harvest or earlier if disease threatens. Repeat every 30 days if needed. NOTE: Spray injury may occur with sensitive varieties such as Lahontan.
Corn* (Field Corn, Popcorn, Seed Corn, Sweet Corn)	Bacterial Stalk Rot	1-3 lbs.	12 lbs.	Begin treatment when disease first appears and repeat every 7- to 10-days if needed. Use the higher rates and shorter spray intervals when conditions favor disease.
Peanut	Cercospora Leaf Spot	1-2.25 lbs.	13.5 lbs.	Begin spraying at 35 to 40 days after planting or when disease symptoms first appear and repeat at 7- to 14-day intervals if needed. Reduce sprays to 7-day intervals during humid weather. Use the higher rates when conditions favor disease. Flowable suffur may be added.
Potato	Early Blight, Late Blight	0.75-3 lbs.	71.4 lbs.	Apply 0.75 to 1.25 lbs. at 5- to 10-day intervals if needed starting when plants are 2 to 6 inches high in locations where disease is light. Apply up to 3 pounds per acre when disease is more severe. Under conditions of severe disease, control with Kocide 2000-O will be improved by tank mixing with other compatible fungicides registered for use on potatoes. Read and follow all label instructions of tank mix partners.
Sugar Beet	Cercospora Leaf Spot	1.5-3.74 lbs.	22.5 lbs.	Begin applications when conditions first favor disease development and repeat at 10- to 14-day intervals if needed. Use the higher rates when conditions favor disease. Addition of a spreader/sticker is recommended.
Wheat, Barley, Oats	Fusarium Head Blight Suppression*, Helminthosporium Spot Blotch, Powdery Mildew Suppression,	1-1.5 lbs.	3.0 lbs.	Make applications for early season disease control through heading. Minimum retreatment interval is 10-days. Use higher rates when conditions favor disease. Addition of adjuvants is recommended.

	Cidillo L
*Not registered for use in C	alifornia.

	Giullie Diotoli, Otelli Must	l	1	
*Not registered for use in 0	California.	•		
			SMALL FRU	ITS
Crop	Disease	Application Rate/Acre	Maximum Annual Rate/Acre	Use Instructions
Blackberry (Aurora, Boysen, Cascade, Chehalem, Logan, Marion, Santiam, Thornless Evergreen)	Anthracnose, Cane Spot, Leaf Spot, Pseudomonas Blight, Purple Blotch, Yellow Rust	3 lbs.	28.6 lbs.	Make fall application after harvest. Apply delayed dormant spray after pruning/training in the spring. If needed, agricultural-type spray oil may be added.
	Anthracnose, Cane Spot, Leaf Spot, Purple Blotch, Yellow Rust	1.5 lbs.	28.6 lbs.	Apply when leaf buds begin to open and repeat when flower buds show white. Repeat on a 7-day interval if needed. If needed, agricultural-type spray oil may be added. NOTE: Crop injury may occur if applied to foliage under certain environmental conditions such as hot or prolonged moist periods. Discontinue applications if signs of crop injury appear.
Blueberry*	Bacterial Canker	3-6 lbs.	24 lbs.	Make first application before fall rains and a second application 4 weeks later. Use the higher rates when conditions favor disease.
	Fruit Rot, Phomopsis Twig Blight	2-4 lbs.	24 lbs.	Dormant Application: Begin applications when bloom buds begin to swell. Make additional applications at 7- to 14-day intervals if needed before blooms open.
Cranberry	Fruit Rot	6 lbs.	36 lbs.	Make first application in late bloom. Apply one or two additional applications at 7- to 14-day intervals if needed depending on disease severity.
	Rose Bloom	6 lbs.	36 lbs.	Apply three sprays on 7- to 14-day schedule if needed as soon as symptoms are observed.
	Bacterial Stem Canker	6 lbs.	36 lbs.	Apply post-harvest and again in spring at bud swell. Apply one or two additional applications at 7 to 14 day intervals if needed depending on disease severity.
	Leaf Blight, Red Leaf Spot, Stem Blight, Tip Blight (<i>Monilinia</i>)	6 lbs.	36 lbs.	Apply delayed dormant spray in the spring. Repeat at 7 to 14-day intervals it needed through pre-bloom.
Currant, Gooseberry	Anthracnose, Leaf Spot	7.5 lbs.	45.7 lbs.	Make initial application after first leaves have expanded. Continue on a 10- to 14-day schedule if needed during wet conditions in the spring. Make an additional application after harvest.
Raspberry	Anthracnose, Cane Spot, Leaf Spot, Pseudomonas Blight, Purple Blotch, Yellow Rust	3 lbs.	28.6 lbs.	Make fall application after harvest. Apply delayed dormant spray after training in the spring. If needed, agricultural-type spray oil may be added.
	Anthracnose, Cane Spot, Leaf Spot, Purple Blotch, Yellow Rust	1.5 lbs.	28.6 lbs.	Apply when leaf buds begin to open and repeat when flower buds show white. Repeat on a 7-day interval if needed. If needed, agricultural-type spray oil may be added. NOTE: Crop injury may occur if applied to foliage under certain environmental conditions such as hot or prolonged moist periods. Discontinue applications if signs of crop injury appear.
Strawberry	Angular Leaf Spot (Xanthomonas), Leaf Blight, Leaf Scorch, Leaf Spot	1.5-2.25 lbs.	23.4 lbs.	Begin application when plants are established and continue on a 7-day schedule throughout the season. Apply in at least 20 gallons of water. Use the higher rates when conditions favor disease. NOTE: Discontinue applications if signs of crop injury appear.

Crop	Disease	Application	Maximum	Use Instructions
Almond only	Destarial Diagram	Rate/Acre	Annual Rate/Acre	Abroad Only Fasherhaid bleet andrel is spiritle injected authority with
Almond only	Bacterial Blast	0.57 lb.	51.4 lbs.	Almond Only: For bacterial blast control in sprinkler irrigated orchards or whe disease is severe, apply 0.75 pound per acre post-bloom at 5- to 14-day interval if needed or just before sprinkling.
Almond, Apricot, Cherry, Plum, Prune	Bacterial Blast (<i>Pseudomonas</i>), Bacterial Canker, Coryneum Blight (Shot Hole)	6-12 lbs.	51.4 lbs.	Make first application before fall rains and a second at late dormant. Use thigher rates when conditions favor disease. Minimum retreatment interval is days. If needed, agricultural-type spray oil may be added. For Cherries: Where disease is severe, an additional application shortly aft harvest may be required. NOTE: Foliar injury may occur from post-bloom sprays on almonds, especia on NePlus varieties.
	Blossom Brown Rot, Coryneum Blight (Shot Hole)	4.3 lbs.	51.4 lbs.	Apply during early bloom. Do not apply after full bloom or injury may occur. U the higher rates when rainfall is heavy and disease pressure is high. Minimu
	Black Knot* (Plum)	3-4.3 lbs.	51.4 lbs.	retreatment interval is 5 days. Make an application at bud swell up to early bloom for early season disea suppression. Apply before full bloom. Minimum retreatment interval is 5 day Use the higher rates when rainfall is heavy and disease pressure is high. NOTE: To avoid plant injury, do not use after full bloom.
	Cherry Leaf Spot* (Sour Cherries Only)	4.3 lbs.	51.4 lbs.	Apply at petal fall as well as 1 to 2 times after petal fall. Use the lower rat where disease infection is light and use the higher rates for a dormant application or where diseases infection is moderate to heavy. Minimum retreatment inten is 5 days. Do not apply to sweet cherry or the English Morello variety as sever injury will result. The addition of 1 to 3 pounds of hydrated lime per pound Kocide 2000-O may reduce crop injury. NOTE: Moderate to severe injury such as leaf spotting and defoliation may occ from post-bloom applications.
Apple	Anthracnose, Blossom Blast, European Canker (<i>Nectria</i>),	9-12 lbs.	45.7 lbs.	Apply before fall rains. Use the higher rates when conditions favor disease. NOTE: Use on yellow varieties may cause discoloration. To avoid discoloratio
	Shoot Blast (Pseudomonas) Apple Scab*, Fire Blight	6-12 lbs.	45.7 lbs.	pick before spraying. Make application between silver-tip and green-tip. Apply as a full cover spray tearly season disease suppression. NOTE: Moderate to severe crop injury may occur from late application.
	Apple Scab*	0.75-1.75 lbs.	45.7 lbs.	discontinue use when green-tip reaches 1/2 inch. Extended spray schedule where fruit finish is not a concern: Continue
		0.5.0.75.11		applications may be made at 5- to 7-day intervals if needed between 1/2 in green-tip and first cover spray. NOTE: Moderate to severe crop injury may result from this extended spr. schedule. It is not intended for fresh market apples or for apples where fruit finis is a concern as it is likely to cause fruit russetting. The addition of 1 to 3 poun
	Fire Blight* Collar Rot, Crown Rot	0.5-0.75 lb. 3 lbs.	45.7 lbs.	of hydrated lime per pound of Kocide 2000-O may reduce crop injury. Mix in 100 gallons of water. Apply 4 gallons of suspension as a drench on the lower trunk area of each tree. Apply in early spring or in fall after harvest for be results. Do not apply to foliage or fruit. NOTE: Do not use if soil pH is below 5.5 since copper toxicity may result.
Avocado	Anthracnose, Blotch, Scab	6-9 lbs.	54 lbs.	Apply when bloom buds begin to swell and continue application at 14- to 30-di intervals for five to six applications. Use the higher rates when conditions fav disease.
Banana, Plantain	Sigatoka (Black and Yellow)	1.5 lbs.	54 lbs.	Apply at 7- to 14-day intervals if needed.
	Black Pitting	3 lbs.	54 lbs.	Mix in 100 gallons of water. Apply to the fruit stem and the basal portion of the leaf crown. Apply during the first and second weeks after fruit emergence.
Cacao	Black Pod	1.5-6.4 lbs.	45 lbs.	Begin applications at the start of the rainy season and continue while infectic conditions persist. Apply 1.5 to 3.5 lbs. at 14-to 21-day intervals if need depending on disease severity. For drier areas, make two to four application using 4.5 to 6.4 pounds per acre according to disease incidence and plantifiednsity.
Coffee	Coffee Berry Disease (Colletotrichum coffeanum)	4.5-6 lbs.	36 lbs.	Apply first spray after flowering and before onset of long rains and then at 14- 28-day intervals if needed until picking. Use the higher rates when condition favor disease.
	Bacterial Blight (Pseudomonas syringae)	4.5-6 lbs.	36 lbs.	Begin spray program before the onset of long rainy periods and continum throughout the rainy season at 14- to 21-day intervals if needed. The critical time for spraying to control this disease is just before, during and after flowering(sespecially when coinciding with wet weather. Use the higher rates when rainfing is heavy and disease pressure is high.
	Leaf Rust (Hemileia vastatrix)	1.5-3 lbs.	36 lbs.	Apply before the onset of rain and then at 14- to 21-day intervals if needed whithe rains continue. Use the higher rates when rainfall is heavy and diseast pressure is high.
	Iron Spot (Cercospora coffeicola), Pink Disease (Corticium salmonicolor)	1.5 lbs.	36 lbs.	Use concentrate or dilute spray. Begin treatment at the start of wet season ar continue at monthly intervals for three applications.
Filbert (WA, OR only)	Bacterial Blight	12-17 lbs.	68.6 lbs.	Apply as a post-harvest spray. In seasons of heavy rainfall, apply a second sprawhen three-fourths of the leaves have dropped. Wait at least 14 days betwee applications. Use the higher rates when rainfall is heavy and disease pressure high. If needed, agricultural-type spray oil may be added.
	Eastern Filbert Blight	12-17 lbs.	68.6 lbs.	Apply as a dilute spray in adequate water for thorough coverage. Mal applications starting at bud swell to bud break and continue at 14-day intervals needed until early May. Thorough coverage is essential. Use the higher rate when rainfall is heavy and disease pressure is high. If needed, agricultural-type spray oil or sticking agent may be added.
Mango	Anthracnose	6-9 lbs.	137 lbs.	Apply at 7-day intervals after fruit set until harvest. Use the higher rates whe rainfall is heavy and disease pressure is high.
Olive	Olive Knot, Peacock Spot	6-9 lbs.	51 lbs.	Make first application before winter rains begin. A second application in ear spring should be made if disease is severe. Apply the higher rates for hear disease pressure or when conditions favor disease development. Minimu retreatment interval is 30 days.
Peach, Nectarine	Bacterial Blast (<i>Pseudomonas</i>), Bacterial Canker, Bacterial Spot (<i>Xanthomonas</i>), Coryneum Blight (Shot Hole), Leaf Curl	6-12 lbs.	51.4 lbs.	Make first application before fall rains and a second at late dormant. For pear leaf curl, late dormant application must be made before leaf buds swell. Use thigher rates when rainfall is heavy and disease pressure is high. If neede agricultural-type spray oil may be added. Minimum retreatment interval is 7-day
	Blossom Brown Rot, Coryneum Blight (Shot Hole), Leaf Curl	4.3 lbs.	51.4 lbs.	Full cover spray at pink bud. Minimum retreatment interval is 5 days.
	Bacterial Spot	0.75 lb.	51.4 lbs.	Post-bloom application applied at first and second cover sprays. Minimu retreatment interval is 5 days. NOTE: Do not spray 3 weeks prior to harvest. Use only recommended rate Spotting of leaves and defoliation may occur from use in cover sprays.
Pear	Fire Blight	0.75 lb.	45.7 lbs.	Apply at 5-day intervals if needed throughout the bloom period. NOTE: Russetting may occur in copper sensitive varieties. Excessive dosage may cause fruit russet on any variety.
	Blossom Blast (Pseudomonas)	9-12 lbs.	45.7 lbs.	Apply before fall rains and again during dormancy before spring growth start Use the higher rates when disease pressure is high or when conditions fav disease development.
Pecan	Kernel Rot, Shuck Rot (Phytophthora cactorum), Zonate Leaf Spot (Cristulariella pyramidalis)	1.5-3 lbs.	24 lbs.	For suppression, apply in sufficient water to ensure complete spray coverage 14 to 28-day intervals if needed, starting at kernel growth, and continue un shucks open. Use the higher rates and shorter spray intervals if frequent rainfoccurs.
	Ball Moss*, Spanish Moss*	4.5-6 lbs.	24 lbs.	Apply in 100 gallons of water in the spring when ball moss is actively growin using 1 1/2 gallons of spray per foot of tree height. Make sure to wet ball most utils thoroughly. The addition of a non-ionic surfactant will improve control. second application may be required after 12 months.
Pistachio	Botryosphaeria Panicle and Shoot Blight, Botrytis Blight, Late Blight (<i>Alternaria</i> <i>alternata</i>), Septoria Leaf Blight	3-6 lbs.	24 lbs.	Make initial application at bud swell and repeat on a 14- to 28-day schedule needed. If disease conditions are severe, use the higher rates and shorter spraintervals.
Quince	Fire Blight	0.75 lb.	45.7 lbs.	Apply at 5-day intervals if needed throughout the bloom period. Apply in adequa water for thorough coverage.
Walnut	Walnut Blight	6-9 lbs.	91 lbs	Apply first spray at early pre-bloom prior to or when catkins are partial expanded. Make additional applications during bloom and early nutlet stage or a 7-day interval if needed when frequent rainfall or extended periods of moistum occur. Thorough coverage of catkins, leaves and nutlets is essential for effective control.

	VEGETABLES				
Crop	Disease	Application Rate/Acre	Maximum Annual Rate/Acre	Use Instructions	
Bean (Dry, Green)	Brown Spot, Common Blight, Downy Mildew*, Halo Blight	0.75-2.25 lbs.	13.5 lbs.	For protective sprays, make first application when plants are 6 inches high; repeal on a 7- to 14-day schedule if needed depending on environmental conditions. Use the higher rates for more severe disease.	
Beet (Table Beet, Beet Greens)	Cercospora Leaf Spot	1.5-3.74 lbs.	22.5 lbs.	Begin applications when conditions first favor disease development and repea at 10- to 14-day intervals if needed. Use the higher rates when conditions favor disease.	
Carrot	Alternaria Leaf Spot, Cercospora Leaf Spot	1.5-2.75 lbs.	14.3 lbs.	Begin applications when disease first threatens and repeat at 7- to 14-day intervals if needed depending on disease severity.	
Celery, Celeriac	Bacterial Blight, Cercospora Early Blight, Septoria Late Blight	1.5 lbs.	15.1 lbs.	Begin applications as soon as plants are first established in the field, repeating at 7-day intervals if needed depending on disease severity and environmental conditions.	
Crucifers (Broccoli; Brussels Sprout; Cabbage; Cabbage, Chinese; Cauliflower; Greens, Collard; Greens, Mustard; Greens, Turnip; Kale; Kohlrabi)	Black Leaf Spot (Alternaria), Black Rot (Xanthomonas), Downy Mildew	0.75-1.5 lbs.	7.57 lbs.	Begin application after transplants are set in the field, or shortly after emergence of field seeded crops or when conditions favor disease development. Apply a 7- to 10-day intervals if needed. Use the higher rates when conditions favor disease NOTE: Reddening of older leaves may occur on broccoli and a flecking of wrapper leaves may occur on cabbage.	
Cucurbits (Cantaloupe, Cucumber, Honeydew, Muskmelon, Pumpkin, Squash, Watermelon)	Alternaria Leaf Spot, Angular Leaf Spot, Anthracnose, Downy Mildew, Gummy Stem Blight, Powdery Mildew, Watermelon Bacterial Fruit Blotch (suppression)	1-2.25 lbs.	15 lbs.	Begin applications prior to disease development and continue while conditions are favorable for disease development. Repeat at 5- to 7-day intervals if needed Use the higher rates when conditions favor disease. NOTE: Crop injury may occur from application at higher rates and shorte intervals. Discontinue use if injury occurs.	
Eggplant	Alternaria Blight, Anthracnose, Phomopsis	1.5 lbs.	22.6 lbs.	Begin applications prior to development of disease symptoms. Repeat sprays at 7- to 10-day intervals if needed depending on disease severity.	
Okra*	Anthracnose, Bacterial Leaf Spot, Leaf Spots, Pod Spot, Powdery Mildew	1.5-3 lbs.	15 lbs.	Begin treatment when disease first threatens and repeat every 5- to-10 days in needed depending on disease severity. Use the higher rates and shorter spray intervals when conditions favor disease.	
Onion, Garlic, Leek	Bacterial Blight, Downy Mildew, Purple Blotch	0.75 - 1.5 lbs.	17.1 lbs.	Begin when plants are 4 to 6 inches high and repeat at 7- to 10-day intervals in needed depending on disease severity. Can cause phytotoxicity to leaves.	
Pea	Powdery Mildew	1-2.25 lbs.	11.3 lbs.	Begin applications when disease symptoms first appear and repeat at 7-day intervals if needed. Use the higher rates when conditions favor disease.	
Pepper	Anthracnose, Bacterial Spot, Cercospora Leaf Spot	1.5-2.25 lbs.	33.9 lbs.	Begin applications when conditions first favor disease development and repeal at 3- to 10-day intervals if needed depending on disease severity. Use the higher rates when conditions favor disease.	
Spinach	Anthracnose, Blue Mold, Cercospora Leaf Spot, White Rust disease	1.5-2.25 lbs.	11.3 lbs.	Begin application when disease first appears or when conditions favor disease development. Repeat at 7- to 10-day intervals if needed. Use the higher rates when conditions favor disease. NOTE: Flecking may occur on spinach leaves.	
Tomato	Anthracnose, Bacterial Speck, Bacterial Spot, Early Blight,	1.5 lbs. (processing)	49.7 lbs. (processing)	Begin applications when disease first threatens and repeat at 3- to intervals if needed depending on disease severity. Where applicable, u	
	Gray Leaf Mold, Late Blight, Septoria Leaf Spot	1.5-3 lbs. (fresh market)	22.8 lbs. (fresh market)	higher rates when conditions favor disease.	
Watercress	Cercospora Leaf Spot	1.5 lbs.	6.06 lbs.	Begin applications when plants are first established in the field, repeating at 7-to 14-day intervals if needed depending on disease severity. Do not exceed four applications per crop. Apply using ground spray equipment at no less than 50 gallons of spray solution per acre.	

VEGETABLES

NOTE: Adequate control may not be obtained when copper tolerant species or


anthomonas bacteria are present.

*Not registered for use in California

*Not registered for use in California

	VINES				
Crop	Disease	Application Rate/Acre	Maximum Annual Rate/Acre	Use Instructions	
Grape	Black Rot, Downy Mildew, Phomopsis, Powdery Mildew	1.5-3 lbs.	57.1 lbs.	Begin applications at bud break with subsequent applications throughout the season depending on disease severity. Repeat at 3-day intervals if needed. Use the higher rates when conditions favor disease. NOTE: Foliage in	
Hops	Downy Mildew	1.5 lbs.	7.57 lbs.	Make crown treatment after pruning, but before training. After training, apply at 10-day intervals if needed. NOTE: Discontinue use two weeks before harvest.	
Kiwi	Erwinia herbicola, Pseudomonas fluorescens, Pseudomonas syringae	6 lbs.	18 lbs.	Apply in 200 gallons of water per acre. Make applications on a 30-day basis. A maximum of three applications may be made.	

	MISCELLANEOUS				
Crop	Disease	Application Rate/Acre	Maximum Annual Rate/Acre	Use Instructions	
Atemoya	Anthracnose	2.25-3.5 lbs.	36 lbs.	Make initial application just before flowering and repeat on a 7-day schedule unl just before harvest. Apply in sufficient water for thorough coverage. Use th higher rates for severe disease.	
Carambola	Anthracnose	4.5-6 lbs.	30 lbs.	Make initial application just before flowering and repeat on a 7-day schedule unl just before harvest. Apply in sufficient water for thorough coverage. Use th higher rates for severe disease.	
Chives	Downy Mildew	1.5 lbs.	7.57 lbs.	Begin applications when plants are established in the field. Repeat application every 7 to 10 days if needed depending on disease conditions.	
Dill	Phoma Leaf Spot, Rhizoctonia Foliage Blight	1.5-2.25 lbs.	11.3 lbs.	Begin applications when plants are first established in the field and repeat at 7 to 10-day intervals if needed depending upon disease severity and environments conditions. Use the higher rates when conditions favor disease.	
Ginseng	Alternaria Leaf Blight, Stem Blight	2-3 lbs.	15 lbs.	Use as a tank mix with the appropriate amount of a product containing the activing redient iprodione. Use label rate of iprodione in 100 gallons of water. Use i accordance with the most restrictive of label limitations and precautions. No lab dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing. Begin Kocide 2000-C "iprodione" applications as soon as plants have emerged in spring. Application should be repeated every 7 days if needed until plants become dommant in Apply tungicides at least 8 hours before rain. Use of a spreader-sticker or stick is advised. NOTE: Alternaria Leaf and Stem Blight is most severe in humid conditions suc as those found in the dense canopies of 2- to 4-year old Ginseng. It is ver important that the stems be thoroughly covered with fungicide; therefore, use spray apparatus which distributes the fungicide throughout the canopy.	
Guava	Anthracnose, Red Algae	2.25-3.5 lbs.	14.1 lbs.	Make initial application just before flowering and repeat every 7 days until jubefore harvest. Apply in sufficient water for thorough coverage. Use the higher rates for severe disease.	
Litchi	Anthracnose	2.25-3.5 lbs.	14.1 lbs.	Make initial application just before flowering and repeat every 7 days until just before harvest. Apply in sufficient water for thorough coverage. Use the higher rates for severe disease.	



CONIFERS

For use on conifers, including Douglas Fir, Fir*, Juniper, Leyland Cypress*, Pine* and Spruce*, in Christmas tree plantings, forest stands and silviculture nurseries.

For control of foliar diseases, apply Kocide 2000-O as a thorough cover spray at rates ranging from 1.5 to 3 pounds per acre. Begin applications in the spring at the initiation of new growth and repeat at 7 to 30 day intervals if needed. Use the higher rates when disease pressure is severe or when environmental conditions favor disease development. Maximum annual rate per acre is 57.1 lbs. ended for use on the listed conifers for control of the following diseases

Crop	Scientific Name	Disease		
Douglas Fir	Pseudotsuga menziesii	Rhabdocline Needlecast		
Fir*	Abies spp.	Needlecasts		
Juniper	Juniperus spp.	Anthracnose, Phomopsis Twig Dieback*		
Leyland Cypress*	X Cupressocyparis leylandii	Cercospora Needle Blight		
Pine*	Pinus spp. Needlecasts			
Spruce*	Picea spp. Needlecasts			
Lichens*: To control lichens on any of the conifers above, apply 6 to 10 pounds of Kocide 2000-O per acre as a dormant application before new growth emerges in the spring. T addition of a non-ionic surfactant will improve control. A second application may be required after 12 months.				

NOTE: Do not buffer or combine with emulsifiable concentrate insecticides. *Not registered for use in California.

2

GREENHOUSE AND SHADEHOUSE CROPS

Notice to User: Kocide 2000-O may be used in greenhouses and shadehouses to control diseases on crops which appear on this label, and specific instructions have been developed for the crops listed. The grower should bear in mind that the sensitivity of crops grown in greenhouses and shadehouses differs greatly from crops grown under field conditions. conditions.

Neither the manufacturer nor seller has determined whether or not Kocide 2000-O can be used safely on all greenhouse and shadehouse grown crops. Consequently; injury arising from the use of Kocide 2000-O on these types of greenhouse and shadehouse crops is the responsibility of the user. The user should determine if Kocide 2000-O can be used safely prior to commercial use. In a small area, apply the recommended rates to the plants in question, i.e., foliage, fruit, etc., and observe for 7 to 10 days for symptoms of

phytotoxicity prior to commercial use.

Apply Kocide 2000-O according to specific rates given for those crops in pounds per acre. One level tablespoon of Kocide 2000-O per 1,000 square feet is equivalent to 1.5 pounds of product per acre. Kocide 2000-O should be applied in adequate water for thorough coverage of plant parts. Begin application at first sign of disease and repeat if needed; use shorter spray intervals during periods when severe disease conditions persist. For maximum annual rates per acre, refer to the crop specific directions.

NOTE: Phytotoxicity may occur on young tender flush when Kocide 2000-O is applied to citrus seedlings grown in greenhouses or shadehouses.

Crop	Disease	Rate per 1,000 Sq Ft	Use Instructions
Citrus (Non-Bearing Nursery)	Brown Rot, Citrus Canker, Greasy Spot, Melanose, Pink Pitting, Scab	3 TBSP	Begin applications when disease first threatens. Repeat at 7- to 30-day intervals if needed depending on disease severity.
Cucumber	Angular Leaf Spot, Downy Mildew	1 - 2 1/2 TBSP	Apply at 5- to 7-day intervals when plants begin to vine. Use the higher rates when conditions favor disease.
Eggplant	Alternaria Blight, Anthracnose, Phomopsis	1 1/2 TBSP	Begin applications prior to development of disease symptoms. Repeat sprays at 7- to 10-day intervals if needed depending on disease severity.
Pepper	Bacterial Spot	1 1/2 - 2 1/2 TBSP	Begin applications when conditions first favor disease development and repeat at 3- to 10-day intervals if needed depending on disease severity. Use the higher rates when conditions favor disease.
Tomato	Anthracnose, Bacterial Speck, Bacterial Spot, Early Blight, Gray Leaf Mold, Late Blight, Septoria Leaf Spot	1 1/2 - 3 TBSP	Begin applications when disease first threatens and repeat at 3- to 10-day intervals if needed depending on disease severity. Use the higher rates when conditions favor disease.

ORNAMENTALS

Use Kocide 2000-O for control of bacterial and fungal diseases of foliage, flower entals in greenhouses, shade houses, outdoor nurseries and outdoor landscape

Use Kocide 2000-O for control of bacterial and fungal diseases of foliage, flowers and stems on ornamentals in greenhouses, shade houses, outdoor nurseries and outdoor landscape plantings.

For ornamental crops in dormancy, apply as a thorough cover spray at rates ranging from 0.75 to 2 pounds per acre of Kocide 2000-O. When new growth is present, apply as a thorough cover spray at rates ranging from 0.75 to 2 pounds per acre of Kocide 2000-O. One level tablespoon of Kocide 2000-O per 1.000 square feet is equivalent to 1.5 pounds of product per acre. Begin application at first sign of disease and repeat at 7- to 14-day intervals if needed; use the higher rates and shorter spray intervals during periods of frequent rains or when severe disease conditions persist. Maximum seasonal rate per acre is 57.1 lbs.

Kocide 2000-O may be used alone or in combination with other fungicides registered for use on ornamentals as a maintenance spray. Use in accordance with the most restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

Notice to User: Plant sensitivities to Kocide 2000-O have been found to be acceptable for the specific genera and species listed on this label under the conditions tested. However, phytotoxicity may occur. Due to the large number of species and varieties of ornamental and nursery plants, and the wide range of growing conditions, it is impossible to test every one for sensitivity to Kocide 2000-O. Neither the manufacturer nor seller has determined whether or not Kocide 2000-O can be safely used on ornamental or nursery plants not listed on this label. The user should determine if Kocide 2000-O can be used safely prior to commercial use.

NOTE: This product may be reactive on masonry and metal surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.

NOTE: This product may be reactive on r	serve for 7- to 10-days for symptoms nasonry and metal surfaces such as g	alvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.
Сгор	Scientific Name	Disease
Aglaonema*	Aglaonema spp.	Bacterial Leaf Spot
Althea (Rose of Sharon)	Hibiscus syriacus	Bacterial Leaf Spot
Andromeda, Japanese*	Pieris japonica	Leaf Spots, Twig Blight
Aralia	Dizygotheca elegantissima	Alternaria, Cercospora Leaf Spot, Xanthomonas Leaf Spot
Arborvitae	Thuja spp.	Alternaria Twig Blight, Cercospora Leaf Blight
Aster*	Aster spp.	Downy Mildew, Leaf Spots
Azalea ¹	Rhododendron spp.	Botrytis Blight, Cercospora Leaf Spot, Phytophthora Dieback, Powdery Mildew
Beech*	Fagus spp.	Leaf Spots
Begonia	Begonia semperflorens	Bacterial Leaf Spot (Erwinia spp., Pseudomonas spp., Xanthomonas spp.)
Bougainvillea	Bougainvillea spectabilis	Anthracnose, Bacterial Leaf Spot
Boxwood*	Buxus spp.	Leaf Spots
Camellia	Camellia japonica, C. sasangua	Anthracnose, Bacterial Leaf Spot
Camphor Tree	Cinnamomum camphora	Pseudomonas Leaf Spot
Canna	Canna spp.	Pseudomonas Leaf Spot
Carnation ¹	Dianthus spp.	Alternaria Blight, Botrytis Blight, Pseudomonas Leaf Spot
Cedar*	Cedrus spp.	Tip Blight
Cherry, Nanking*	Prunus tomentosa	Bacterial Leaf Spot
Chinese Tallow Tree	Sapium sebiferum	
	<u>'</u>	Bacterial Leaf Spot (Pseudomonas spp., Xanthomonas spp.)
Chrysanthemum 1	Chrysanthemum morifolium	Botrytis Blight, Pseudomonas Leaf Spot, Septoria Leaf Spot
Cotoneaster	Cotoneaster spp.	Botrytis Blight
Crabapple*	Malus spp.	Fire Blight
Cypress*	Cupressus spp.	Twig Blight
Dahlia	Dahlia pinnata	Alternaria Leaf Spot, Botrytis Gray Mold, Cercospora Leaf Spot
Delphinium*	Delphinium spp.	Leaf Spots
Dianthus	Dianthus spp.	Bacterial Soft Rot, Bacterial Spot
Dogwood, Flowering	Cornus florida	Anthracnose
Dogwood, Kousa*	Comus kousa	Fungal Leaf Spots
Douglas Fir	Pseudotsuga menziesii	Rhabdocline Needlecast
Dracaena*	Dracaena marginata	Bacterial Leaf Spot
Dumb Cane*	Dieffenbachia spp.	Bacterial Leaf Spot
Dusty Miller	Senecio cineraria	Bacterial Leaf Spot (Pseudomonas cichorii)
Echinacea	Echinacea spp.	Bacterial Leaf Spot (Pseudomonas cichorii)
Elm, Chinese	Ulmus parvifolia	Xanthomonas Leaf Spot
Euonymus	Euonymus spp.	Anthracnose, Botrytis Blight
Fern Boston*	Nephrolepis exaltata	Bacterial Leaf Spot
Fern, Holly	Cyrtomium falcatum	Pseudomonas Leaf Spot
Fig, Weeping*	Ficus benjamina	Bacterial Leaf Spot
Filbert (Ornamental)*	Corylus spp.	Filbert Blight
Fir*	Abies spp.	Needlecasts
Gardenia	Gardenia jasminoides	Alternaria Leaf Spot, Botrytis Bud Rot, Cercospora Leaf Spot
Geranium	· · · · · · · · · · · · · · · · · · ·	Alternaria Leaf Spot, Botrytis Gray Mold, Cercospora Leaf Spot
	Pelargonium spp.	
Gladiola	Gladiolus spp.	Alternaria Leaf Spot, Anthracnose, Bacterial Leaf Blight, Botrytis Gray Mold
Golden Rain Tree	Koelreuteria paniculata	Bacterial Leaf Spot
Grape Ivy*	Cissus spp.	Bacterial Leaf Spot
Hawthorn*	Crataegus spp.	Fire Blight
Hibiscus ⁴	Hibiscus spp.	Bacterial Leaf Spot
Holly*	llex spp.	Bacterial Blight, Leaf Spots
Honeylocust*	Gleditsia triacanthos	Bacterial Leaf Spot
Honeysuckle, Tatarian*	Lonicera tatarica	Bacterial Leaf Spot
Impatiens	Impatiens wallerana	Bacterial Leaf Spot
Indian Hawthorn 5	Raphiolepis indica	Anthracnose, Entomosporium Leaf Spot
Iris 6*	Iris spp.	Bacterial Leaf Spot
Ivy (English, Algerian) 1	Hedera helix, H. canariensis	Xanthomonas Leaf Spot
Ixora	Ixora coccinea	Xanthomonas Leaf Spot
Juniper	Juniperus spp.	Anthracnose, Phomopsis Twig Dieback*
Lantana	Lantana camera	Bacterial Leaf Spot
Leyland Cypress*	X Cupressocyparis leylandii	Cercospora Needle Blight
Lilac	Syringa spp.	Cercospora Leaf Spot, Pseudomonas Blight*
Lily, Easter ²	Lilium longiflorum	Botrytis Blight
Linden*	Tilia spp.	Anthracnose, Leaf Blight
Loblolly Bay	Gordonia lasianthus	Anthracnose
Loquat (ornamental)	Eriobotrya japonica	Colletotrichum spp., Entomosporium maculata
Magnolia (Southern)	Magnolia grandiflora	Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot
Magnolia (Sweet Bay)	Magnolia virginiana	Anthracnose
agriolia (Ombot Day)	agriona virginiana	
Magnolia (Oriental)	Magnolia soulangiana	Bacterial Leaf Snot
Magnolia (Oriental)	Magnolia soulangiana Mandevilla son	Bacterial Leaf Spot Anthracnose
Mandevilla	Mandevilla spp.	Anthracnose
Mandevilla Maple*	Mandevilla spp. Acer spp.	Anthracnose Pseudomonas Leaf Blight
Mandevilla Maple* Marigold	Mandevilla spp. Acer spp. Tagetes spp.	Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot
Mandevilla Maple* Marigold Mountain-Ash*	Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp.	Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight
Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted*	Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis	Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot
Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted* Mulberry, Weeping	Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis Morus alba	Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot Bacterial Leaf Spot
Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted* Mulberry, Weeping Narcissus*	Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis Morus alba Narcissus spp.	Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot Bacterial Leaf Spot Leaf Blight
Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted* Mulberry, Weeping Narcissus* Nephthytis*	Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis Morus alba	Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot Bacterial Leaf Spot Leaf Blight Bacterial Leaf Spot
Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted* Mulberry, Weeping Narcissus* Nephthytis* Oak*	Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis Morus alba Narcissus spp. Syngonium podophyllum Quercus spp.	Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot Bacterial Leaf Spot Leaf Blight Bacterial Leaf Spot Leaf Spot Leaf Spots
Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted* Mulberry, Weeping Narcissus* Nephthytis*	Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis Morus alba Narcissus spp. Syngonium podophyllum	Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot Bacterial Leaf Spot Leaf Blight Bacterial Leaf Spot
Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted* Mulberry, Weeping Narcissus* Nephthytis* Oak*	Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis Morus alba Narcissus spp. Syngonium podophyllum Quercus spp.	Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot Bacterial Leaf Spot Leaf Blight Bacterial Leaf Spot Leaf Spot Leaf Spots
Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted* Mulberry, Weeping Narcissus* Nephthytis* Oak* Oak, Laurel	Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis Morus alba Narcissus spp. Syngonium podophyllum Quercus spp. Quercus laurifolia	Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot Bacterial Leaf Spot Leaf Blight Bacterial Leaf Spot Leaf Spots Algal Leaf Spot (Cephaleuros virescens)
Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted* Mulberry, Weeping Narcissus* Nephthytis* Oak* Oak, Laurel Oleander	Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis Morus alba Narcissus spp. Syngonium podophyllum Quercus spp. Quercus laurifolia Nerium oleander	Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot Bacterial Leaf Spot Leaf Blight Bacterial Leaf Spot Leaf Spots Algal Leaf Spot (Cephaleuros virescens) Bacterial Leaf Spot, Fungal Leaf Spot
Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted* Mulberry, Weeping Narcissus* Nephthytis* Oak* Oak, Laurel Oleander Oregon Grape Holly*	Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis Morus alba Narcissus spp. Syngonium podophyllum Quercus spp. Quercus laurifolia Nerium oleander Mahonia aquifolium	Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot Bacterial Leaf Spot Leaf Blight Bacterial Leaf Spot Leaf Spots Algal Leaf Spots Bacterial Leaf Spot, Fungal Leaf Spot Leaf Spots Algal Leaf Spot, Fungal Leaf Spot Leaf Spots
Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted* Mulberry, Weeping Narcissus* Nephthytis* Oak* Oak, Laurel Oleander Oregon Grape Holly* Pachysandra	Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis Morus alba Narcissus spp. Syngonium podophyllum Quercus spp. Quercus laurifolia Nerium oleander Mahonia aquifolium Pachysandra procumbens	Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot Bacterial Leaf Spot Leaf Blight Bacterial Leaf Spot Leaf Spots Algal Leaf Spot, (Cephaleuros virescens) Bacterial Leaf Spot, Fungal Leaf Spot Leaf Spots Volutella Leaf Blight
Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted* Mulberry, Weeping Narcissus* Nephthytis* Oak* Oak, Laurel Oleander Oregon Grape Holly* Pachysandra Palm, Date (ornamental)	Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis Morus alba Narcissus spp. Syngonium podophyllum Quercus spp. Quercus laurifolia Nerium oleander Mahonia aquifolium Pachysandra procumbens Phoenix canariensis	Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot Bacterial Leaf Spot Leaf Blight Bacterial Leaf Spot Leaf Spots Algal Leaf Spot (Cephaleuros virescens) Bacterial Leaf Spot, Fungal Leaf Spot Leaf Spots Volutella Leaf Blight Pestalotia Leaf Blight
Mandevilla Maple* Marigold Mountain-Ash* Mulberry, Contorted* Mulberry, Weeping Narcissus* Nephthytis* Oak* Oak, Laurel Oleander Oregon Grape Holly* Pachysandra Palm, Date (ornamental) Palm, European Fan	Mandevilla spp. Acer spp. Tagetes spp. Sorbus spp. Morus bombycis Morus alba Narcissus spp. Syngonium podophyllum Quercus spp. Quercus laurifolia Nerium oleander Mahonia aquifolium Pachysandra procumbens Phoenix canariensis Chamaerops humilis	Anthracnose Pseudomonas Leaf Blight Alternaria Leaf Spot, Botrytis Leaf Rot, Cercospora Leaf Spot, Flower Rot Fire Blight Bacterial Leaf Spot Bacterial Leaf Spot Leaf Blight Bacterial Leaf Spot Leaf Spots Algal Leaf Spot (Cephaleuros virescens) Bacterial Leaf Spot, Fungal Leaf Spot Leaf Spots Volutella Leaf Spot, Fungal Leaf Spot Pestalotia Leaf Spot Pestalotia Leaf Spot

ORNAMENTALS (continued) Jse Kocide 2000-O for control of bacterial and fungal diseases of foliage, flowers and stems on ornamentals in greenhouses, shade houses, outdoor nurseries and outdoor landscape

Use Kocide 2000-O for control of bacterial and fungal diseases of foliage, flowers and stems on ornamentals in greenhouses, shade houses, outdoor nurseries and outdoor landscape plantings.
For ornamental crops in dormancy, apply as a thorough cover spray at rates ranging from 0.75 to 2 pounds per acre of Kocide 2000-O. When new growth is present, apply as a thorough cover spray at rates ranging from 0.75 to 2 pounds per acre of Kocide 2000-O. One level tablespoon of Kocide 2000-O per 1,000 square feet is equivalent to 1.5 pounds of product per acre. Begin application at first sign of disease and repeat at 7- to 14-day intervals if needed; use the higher rates and shorter spray intervals during periods of frequent rains or when severe disease conditions persist. Maximum seasonal rate per acre is 57.1 lbs.
Kocide 2000-O may be used alone or in combination with other fungicides registered for use on ornamentals as a maintenance spray. Use in accordance with the most restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

Notice to User: Plant sensitivities to Kocide 2000-O have been found to be acceptable for the specific genera and species listed on this label under the conditions tested. However, phytotoxicity may occur. Due to the large number of species and varieties of ornamental and nursery plants, and the wide range of growing conditions, it is impossible to test every one for sensitivity to Kocide 2000-O. Neither the manufacturer nor seller has determined whether or not Kocide 2000-O can be safely used on ornamental or nursery plants not itseld on this label. The user should determine if Kocide 2000-O per present and sensitivity to commercial use. In a small area, apply the recommended rates to the plants in question, i.e., bedding plants, foliage, etc., and observe for 7- to 10-days for symptoms of phytotoxicity prior to commercial use.

NOTE: This product may be reactive

Crop	Scientific Name	Disease	
Palm, Washingtonia	Washingtonia robusta	Pestalotia Leaf Spot	
Peach (Flowering) (ornamental) 3*	Prunus spp.	Bacterial Blast, Brown Rot, Fire Blight	
Pear (Flowering) (ornamental)	Pyrus calleryana	Fire Blight, Leaf Spots	
Pentas (Egyptian Star)	Pentas spp.	Bacterial Leaf Spot (Pseudomonas spp.*, Xanthomonas spp.)	
Peony	Paeonia spp.	Botrytis Blight	
Periwinkle	Catharanthus roseus, Vinca spp.	Phomopsis Stem Blight	
Philodendron	Philodendron selloum	Bacterial Leaf Spot	
Phlox	Phlox spp.	Alternaria Leaf Spot	
Photinia (Red Tip)	Photinia x fraseri, P. glabra	Anthracnose, Entomosporium Leaf Spot	
Pine*	Pinus spp.	Needlecasts	
Pistachio (ornamental)	Pistacia chinensis	Anthracnose	
Plantain Lily ⁶	Hosta spp.	Bacterial Leaf Spot	
Plum (Flowering) (ornamental) 3*	Prunus spp.	Bacterial Blast, Brown Rot, Fire Blight	
Pothos*	Scindapsus spp.	Bacterial Leaf Spot	
Powder Puff Plant	Calliandra spp.	Bacterial Leaf Spot	
Pyracantha	Pyracantha spp.	Fire Blight, Scab	
Rhododendron	Rhododendron spp.	Alternaria Flower Spot	
Rose 1	Rosa spp.	Black Spot, Powdery Mildew	
Snapdragon	Antirrhinum majus	Anthracnose, Dieback, Downy Mildew	
Spathe Flower*	Spathiphyllum spp.	Bacterial Leaf Spot	
Spirea*	Spiraea spp.	Fire Blight	
Spruce*	Picea spp.	Needlecasts	
Sycamore	Platanus spp.	Anthracnose, Leaf Spots*	
Tulip	Tulipa spp.	Anthracnose, Botrytis Blight	
Umbrella Tree*	Schefflera spp.	Bacterial Leaf Spot	
Verbena	Verbena spp.	Xanthomonas Leaf Spot	
Viburnum	Viburnum odoratissimum, V. plicatum, V. suspensum	Anthracnose	
Viola (Pansy, Violet)	Viola spp.	Downy Mildew	
Willow	Salix spp.	Anthracnose	
Yew*	Taxus spp.	Needle Blight	
Yucca (Adam's Needle)	Yucca spp.	Cercospora Leaf Spot, Septoria Leaf Spot	
Zinnia*	Zinnia spp.	Leaf Spots	

Discoloration of foliage and/or blooms has been noted on some varieties. To prevent residues on commercial plants, do not spray immediately before selling season.

Apply Kocide 2000-0 at 2.25 to 3,75 pounds per acre. Maximum seasonal rate per acre is 214 lbs. Do not apply any additional copper pesticide to this land for 36 months. Apply Kocide 2000-O at 2.25 to 3.75 pounds per acr
 Apply dormant through bloom only.
 Hibiscus - Do not apply to plants in flower.
 For Indian Hawthom use 1.5 to 3.0 pounds per acre.
 Some cultivars may be sensitive to Kocide 2000-O.
 Not registered for use in California.

NOTE: Phytotoxicity may depend on varietal differences. If unfamiliar with the use of Kocide 2000-O, apply the recommended rate to a few plants and observe after 7 to 10 days for symptoms of phytotoxicity.

Control of Ball Moss*, Spanish Moss* and Lichens* on Ornamental and Shade Trees: Apply Kocide 2000-O in early spring when the trees are dormant. Apply 4.5 to 6 pounds of Kocide 2000-O in 100 gallons of water, using 1 1/2 gallons of spray per foot of tree height. Be sure to thoroughly wet ball moss tufts, Spanish moss or lichens. The addition of a non-ionic surfactant will improve control. A second application may be required after 12 months.

NOTE: Kocide 2000-O may be injurious to some ornamental plants growing beneath the trees. This product may be reactive on masonry and metal surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.

Cold Storage Protection for Dormant Rootstock*: To protect bars-roof unsery trees from Phytophthora Crown Rot and Botrytis, use 2 to 3 pounds of Kocide 2000-O per 100 gallons of water. Apply as a dip or spray to the roots and lower stems of dormant rootstock prior to placing in cold storage. Do not apply to rootstock less than 2 years old.

*Not registered for use in California.

GENERAL CHEMIGATION INSTRUCTIONS

Apply this product only through one or more of the following types of systems: sprinkler, including center pivot, lateral move, traveler, big gun, or plastic pipe solid set system(s). Do not apply this product through any other type of irrigation systems. In California, do not apply in systems which contain aluminum parts or components.

Crop injury, lack of effectiveness or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Shut off injection equipment after treatment and continue to operate irrigation system until Kocide 2000-O has been cleared from the last sprinkler head.

CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS
Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into the reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

NOTE: It must be determined if proper application equipment is availabl

recommended. Kocide 2000-O should be added through a traveling irrigation system continuously or at the last 30 minutes of solid set irrigation systems. Shut off injection equipment after treatment and continue to operate irrigation system until Kocide 2000-O has been cleared from the last sprinkler head.

SPRINKLER CHEMIGATION
The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The postificial principle principle principle principle are contain a functional customatic quick closing check valve to prevent the flow of fluid back toward the injection pump.

In e system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must also contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The epsticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interiock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fittled with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

NOTE: It must be determined if proper application equipment is available and if waste associated with its use can be properly handled. Agricultural chemicals are often reactive with the materials used in the construction of application equipment, such as aluminum, rubber and some synthetic materials. This factor should be taken into consideration when selecting proper application equipment. It is necessary that all application equipment be thoroughly flushed with clean water after each day's use.

When mixing, fill the nurse tank half full with water. Add Kocide 2000-O slowly to tank while hydraulic or mechanical agitation is operating and continue filling with

recommended.
Kocide 2000-O should be added through a traveling irrigation system continuously or at the last 30 minutes of solid set irrigation systems. Shut off injection equipment after treatment and continue to operate irrigation system until Kocide 2000-O has been cleared from the last sprinkler head.

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet size:

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet size:

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind speed:

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition (approximately 3 to 10 mph), and there are not sensitive areas within 250 feet downwind.

Temperature Inversions:

If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Other State and Local Requirements:

Applicators must follow all state and local pesticide drift requirements regarding application of copper compounds. Where states have stringent regulations, they must be observed.

Equipment:

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Additional requirements for aerial applications:

- The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

- Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.

- When applications are made with a crosswind, the swath must be displaced downwind. The applicator must compensate for this displacement at the up and downwind edge of the applications are made with a crosswind, the swath must be displaced downwind. The applicator must compensate for this displacement at t

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest cannot be disposed of by use according to laurel instructions, without pour other research office for guidance.

EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Completely empty paper or plastic bag, fiber sack or drum liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer for recycling if available or dispose of empty paper or plastic bag, fiber sack or fiber drum and liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact ChemTel

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States. "Curtec" is a registered trademark of Bei Incorporated "Tre-Hold" is a registered trademark of Amvac Chemical Corporation.

LIMITATION OF WARRANTY AND LIABILITY

LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read this Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of Certis. These risks can cause: ineffectiveness of the product, crop injury, or injury to non-target crops or plants. WHEN YOU BUY OR USE TIPS PRODUCT, YOU ACREE TO ACCEPT THESE RISKS.

Certis warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, CERTIS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL CERTIS OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE INSULTIVE OF CERTIS OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY, WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF CERTIS OR SELLER. THE REPLACEMENT OF THE PRODUCT.

To the extent consistent with applic

Certis LLC, 9145 Guilford Road, Suite 175, Columbia, MD 21046 All rights reserved