# **SitePro**

# Flex-4 with Dimension® Plus Quinclorac

ACTIVE INGREDIENTS:	
2,4-D	0.675%
Dicamba	0.060%
Dithiopyr	0.121%
Quinclorac	0.346%
OTHER INGREDIENTS:	<u>98.798%</u>
TOTAL	100.000%

#### THIS PRODUCT CONTAINS:

13.50 lb 2,4-dichlorophenoxyacetic acid equivalent per ton

1.20 lb 3.6-dichloro-o-anisic acid equivalent per ton

2.42 lb dithiopyr per ton

6.92 lb quinclorac per ton

Isomer Specific by AOAC Methods

FPA REG NO 2217-927-69759 EPA EST. NO. 69759-OR-002 (Lot beings with MAS)

Dimension® is a trademark of DOW AgroSciences. EPA EST. NO. 69759-OR-004 (Lot begins with BAN)

20-0-6

## GUARANTEED ANALYSIS

Total Nitrogen (N)	20.00%
Soluble Potash (K <sub>2</sub> 0)	6.00%
Sulfur (S)	2.00%

Derived from: Polymer Coated Urea, Potassium Sulfate, Urea

\*10.00% Slow Release Urea Nitrogen from Polymer Coated Urea.

# KEEP OUT OF REACH OF CHILDREN CAUTION

# STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.



Net Weight: 50lbs. (22.68 Kg) Manufactured and Guaranteed by: Marion Ag Service, Inc. 7746 St. Paul Hwy NE St. Paul. OR 97137

### PRECAUTIONARY STATEMENTS

### Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

## Personal Protective Equipment (PPE)

All loaders, applicators, and other handlers must wear:

- long-sleeved shirt and long pants, shoes plus socks, and
- waterproof gloves

### **User Safety Requirements**

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### **User Safety Recommendations**

- Users should wash hands thoroughly after handling and before eating, drinking, chewing gum using tobacco, or using the toilet
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly
- and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
  Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

# 2217-927-69759 SitePro Flex-4 with Dimension Plus Quinclorac 20-0-6 20240206 22 69759 .pdf

First Aid	
If in eyes:	Hold eye open and rinse slowly and gently with water for 15 — 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed:	Call a poison control center or doctor immediately for treatment advice.     Have person sip a glass of water if able to swallow.     Do not induce vomiting unless told to do so by a poison control center or doctor.     Do not give anything by mouth to an unconscious person.
If on skin or on clothing:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-877-800-

#### Environmental Hazards

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runof may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

Groundwater Advisory: This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into the groundwater if used in areas where soils are permeable, particularly where the water table is shallow. Application around a cistern or well may result in contamination of drinking water or groundwater

Non-Target Organism Advisory Statement: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by minimizing drift.

Surface Water Advisory: This product may impact surface water quality due to runoff of rainwater This is especially true for poorly draining soils and soils with shallow ground. This product is classified as having a high potential for reaching surface water via runoff for several weeks after application.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Do not enter or allow people (or pets) to enter the treated area until dusts have settled

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed or for research purposes.

This product is not sold, distributed, or registered within the state of New York, Do not use in the state of New York.

#### Resistance Management

For resistance management, this product contains Group 3 and Group 4 herbicides. Any weed population may contain or develop plants naturally resistant to this product and other Group 3 or 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 3 or 4 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same
- Use tank mixtures with herbicides from a different group if such use is permitted; where
  information on resistance in target weed species is available, use the less resistance-prone
  partner at a rate that will control the target weed(s) equally as well as the more resistanceprone partner. Consult your local extension service or pest control advisor if you are unsure as
- to which active ingredient is currently less prone to resistance.
   Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use, and that considers mechanical control methods, cultural (e.g., timing to favor the turf and not the weeds), biological (weed-competitive varieties) and other management practices.

  Scout area prior to application to identify the weed species present and their growth stage to
- determine if the intended application will be effective.
- determine if the intended application will be effective.
  Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: 1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; 2) a spreading patch of non-controlled plants of a particular weed species; 3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available,
- Contact your local extension specialist or pest control advisor for additional pesticide resistance ent and/or integrated weed-management recommendations for specific types of turf
- For further information or to report suspected resistance, call 877-800-5556.

### PRODUCT DESCRIPTION

This product is formulated to feed your lawn and control listed annual grasses and broadleaf weeds commonly found in home lawns. The product contains a lawn fertilizer and four different herbicides

- The fertilizer contains slow release nutrients that may extend the availability for turfgrass
- The four herbicides combine to kill listed weeds at different stages of development
- Grass weeds: Preemergent control of crabgrass, foxtails, and other annual grass weeds as listed. Postemergent control of crabgrass.
- Broadleaf weeds: Preemergent control of chickweed, henbit, knotweed, shepherd's purse, spurge, woodsorel, and other broadleaf weeds as listed. Post emergent control of over 250 broadleaf weeds including dandelion, chickweed, thistle and clover.

## WHERE TO USE:

This product may be used on the following established turf species:

Established Cool Season Turfgrass: Kentucky bluegrass, rough bluegrass (Poa trivialis), perennial ryegrass, fine fescues, tall fescue, seashore paspalum

Established Warm Season Turfgrass: Bermudagrass, buffalograss, zoysiagrass This Product may NOT be used on: annual bluegrass (Poa annua), bahiagrass, bentgrass, carpetgrass, centipedegrass, dichondra, St. Augustine. Do not use this product on turfgrass where desirable clovers are present

This Product may be used on warm-season and cool-season turfgrass in 4 use sites.

- · Residential/domestic sites are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.
- Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, industrial and institutional buildings, recreation areas, fairgrounds, areas adjacent to athletic fields and paved areas.
- Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations including, but not limited to hospitals, nursing homes, schools, museums, libraries, sport facilities, golf courses (fairways and roughs only), and office buildings
- Non-cropland sites include highway rights-of-way (principal, interstate, county, private, and unpayed roads): roadsides, roadside ditches, road shoulders, road embankments, dividers and medians; municipal, state and federal lands; airports and military installations

Guidelines to follow: The use of SitePro Flex-4 with Dimension Plus Quinclorac on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, chemical injury, disease, insects, pests, excessive temperatures, or lack of moisture may increase the chances for turfgrass injury. Allow turfgrass weakened by these stresses to recover before application of

#### WEEDS CONTROLLED hen applied as directed, this product will control and/or suppress\* the following weeds Annual Grass Weeds Goosegrass\* (crowfootgrass) Annual bluegrass (Poa Foxtail (green, yellow, Crabgrass (smooth annua) southern large) nnual ryegras Proadleaf Weeds Amaranth, purple Cupid shaving Brush Mustard, ball peedwell, commor

Mustard, black beedwell, corn maranth, slende ster, heath Aster, Daisy, English Mustard, blue peedwell, ivyleaf white heath Aster Daisy, oxeye Mustard, hedge Speedwell, persian purple Aster, white Dandelion, commor Mustard, indian. Speedwell, purslane rairie Artichoke Dandelion, false Mustard, tansy Speedwell, slender ustrian fieldcress needwell, snow Mustard, tumble Day flower Bedstraw, Catchweed Deadnettle Mustard, white beedwell, thymelea Mustard, wormseed Bedstraw, smooth Dichondra piny amaranth Bedstraw, Northerr Dock, broadleaf Nettle, stinging (burning) Spurge, spotted 3edstraw, pineywoods Spurge, prostrate Dock, curly Nettle, tall Beggarticks, hairy Dogbane Nettle, wood Spurweed etony, Florida Parsley-piert . Star of Bethlehen Dogfenne indweed, Field Dollarweed (Pennyw Strawberry, India Rindweed Hedge Dove weed Pearlwort mock Bitter Wintercress Elderberry Pennycress Tansv ragwort ttercress, hairy (frenchweed) alse sunflower Mugwort histle blessed itterweed Black-eved Susan Fiddleneck Pepperweed, perennia histle, bull Black Medic Filaree, redstem Pepperweed, virginia histle, canada Blood flower milkweed ilaree, whitester (Peppergrass) histle, flodmar Rlue lettuce leabane, daisy Pigweed, prostate prairie) Brassbuttons annual) Piaweed, redroot histle, musk Thistle, platte ristly oxtongue leabane, rough Pigweed, russian Galinsoga, hairy rigweed, smooth Thistle, plumeless Piaweed, tumble Buckhorn Galinsoga, smallflowe Thistle russian Bullnettle Goldenrod Pineapple weed Thistle, scotch Ground ivy Plains coreopsis histle, wavvlea (tickseed) Thistle, yellow turdock Groundsel urweed. lawn Gumweed Plantain, blackseed (horrible) (spurweed) Hairy fleabane Plantain, bracted Trailing crownvetch uttercup, bulbous lawkweed, meadov Plantain, broadlea Buttercup, creeping Hawkweed mouseea Plantain buckhorn Vervain blue Buttonweed, common Hawkweed, orange Plantain, hoary Vervain, hoary Hawkweed, yellow Plantain, narrowleaf Vervain, prostrat (poorjoe) -lealall Plantain, slender /ervain, tall Plantain, woolly Carolina geranium (wild) Heartleaf drymary Vetch, bird Vetch, hairy Carpetweed Poison ivv Poison oal etch, milk Catsear, spotted (false Hoary cress Pokeweed Vetch, narrowlea dandelion) Horsenettle Prairie Sunflower Vetch wild Chamber bitter Prickly lettuce (compass Horseweed Virginia creepe Chickweed, commor hickweed mouseear Johnny-Jumpup viole . Prickly sida Western salsify Chickweed, sticky Knawel Prostrate knotweed Wild aster Knotweed, prostrate Wild buckwheat Chicory Puncturevine inquefoi Kochia Purslane, commo Wild carrot Clover crimson Lambsquarters Puslev, brazil Wild four-o-clock Pusley, florida Clover, alyce Lespedeza Wild garlic Mallow, alkali Clover, hop Pusley, large flowe Wild lettuce Clover, rabbit foot Mallow, bristly Wild marigold Ragweed, bur Clover, red Mallow, commor Ragweed, commor Wild mustard Clover, strawberry Mallow, dwarf Ragweed, lancleat (vellow rocket) Mallow, venice Wild onion Clover, yellow swee Clover, white sweet Marcela Redstem filaree Wild parsnip Clover, white Matchweed Rough cinquefoil Wild radish Mexicanweed Scarlet pimpernel Wild rape Cockle . Wild strawberry Cocklebui Moneywort Shepardspurse Cocklebur, oriental Morning glory, bigroot Smartweed. Wild sweet potato Morning glory, ivyleaf Wild violet Cocklebur, spiny advsthumb ommon mullein Morning glory, mexican Smartweed, pale Woodsorrel. Smartweed. orn Spurry reeping Beggarwee Morning glory, tall Pennsylvania Woodsorrel, violet Morning glory, wolly Smooth chaff-flower Woodsorrel Creeping Jenny reeping charlie Smoothdock ellow (Oxalis) Sorrel, red (sheep Wooly croton Sowthistle, annual Wormseed Sowthistle, spiny arrow, common Spanish needles arrow, western Speedwell bi-lobed Yarrow vellow

for acceptable control of barnyardgrass and goosegrass, two applications made at a 6 to 10 week

DITHIOPYR	GROUP	3	HERBICIDE
2,4-D DICAMBA QUINCLORAC	GROUP	4	HERBICIDE

This product may be applied as a single application or as split applications in the spring, early summer, or fall,

For annual grasses such as crabgrass (smooth, southern and large) and foxtail (green,

yellow, and giant): Apply in the spring and early summer. For best results, apply this product from 2 weeks before weed germination until the weed is 4 inches tall. For dense populations, make a second application no sooner than 30 days after the first application.

For broadleaf weed control: Spring applications: for [Dandelion, common]. Fall applications: for [Chickweed, common]. For best results, apply when broadleaf weeds are young and actively growing with adequate soil moisture and favorable temperatures (50° to 80° F)

- This product needs to stick to the leaves of broadleaf weeds and should be applied when the grass is wet from dew or rain.
- Poor results will occur if applied to dry lawns. When dry conditions exist, irrigate for water the turfgrass 24 hours before application
- For best results, irrigate after the application. However do not irrigate [or water] sooner than 24 hours after application
- Avoid applications during gusty wind conditions.
- Do not spread granules onto flowers, vegetables, and ornamental shrubbery.
- Turf species listed on this label can be seeded, sodded, sprigged, or plugged into the treated areas 4 weeks after the application of this product.
- When treating newly planted areas, wait approximately 6 weeks after seedling emergence before applying this product.
- Delay mowing 2 days before and until 3 days after the application of this product. Additional stress from low mowing heights may increase the possibility of turf injury.
- Clippings from the first three mowings should be left on the treated area. Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in

#### How Much To Use

Use a drop or rotary spreader designed to apply granular fertilizers. Avoid skips or excessive overlaps as poor weed control or turfgrass injury may result.

Apply 2.4 to 4.5 pounds of product per 1,000 sq.ft. per application to established turfgrass

The maximum broadcast application rate is 4.5 pounds of product per 1.000 sq.ft, per application The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum annual (seasonal) rate is 9 pounds of product per 1,000 sq.ft.

# CALIBRATION INSTRUCTIONS

For 4.5 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 220 square feet and adjust to apply at the rate of 1 pound of product per 220 square feet.

# STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal

PESTICIDE STORAGE: Keep in original container in a cool, dry area, away from extreme heat and cold.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. To avoid wastes, use the entire contents in this bag by application according to the label instructions. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container (bag) Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by

## LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Information regarding the contents and levels of metals in this product is available on the Internet at http://www.aapfco.org/metals.html

Rev 10/04/2023 BL/SCS

Blend #: 31439