

STORAGE AND DISPOSAL: STORAGE: Do not contaminate water, food, or feed by storage or disposal. Do not open or remove from package before ready to install. Store in a cool, dry place. Foil Pouch: Do not open or remove from package before ready to use the before ready to use the property of the property o FIRST AID: IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person. IF INHALED: Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison gently with water for 13-20 minutes. Nemove contact tenses, if present, arter the first 5 minutes, then continue missing eye. Call a poison control center or doctor for treatment advice. IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. NOTE TO PHYSTICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. PolSoNING SYMPTOMS: CONTAINS AN ORGANOPHOSPHATE THAT INHIBITS CHOLINESTERASE.

Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps. <u>TREATMENT</u>. Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10 minute intervals in doses of 1 to 2 milligrams until complete atropinization has occurred. Morphine is contraindicated. 2-PAM is also antidotal and may be administered in conjunction with atropine. Clear chest by postural drainage. Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment. FOR THE FOLLOWING EMERGENCIES, PHONE 24 HOURS A DAY: For Medical Emergencies phone 1-800-222-1222. For Transportation Emergencies, including spill, leak or fire, phone 1-800-424-9300. For Product Use Information phone: 1-800-800-1819. PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING: May be fatal if swallowed. Do not get in mouth. Harmful if inhaled. Causes moderate eye irritation. Avoid breathing vapors. Avoid warkinus: way do eratal ir swarlowed. Oo not get in mouth, rahmful in inhaled, causes moderate eye irritation, avoid creaming Vapors, avoid contact with eyes or clothing. After prolonged storage, a small amount of liquid may form on strip. Do not get liquid in eyes. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Do not use in hospitals or clinic rooms, such as patient rooms, wards, nurseries, operating or emergency areas. Do not use in any area of the home where people will be present for extended periods of time. Do not use in kitchens. Do not use in the food/feed area of restaurants or any area where food is prepared or served. Not to be taken internally by humans or animals. **NOTICE:** Buyer assumes all responsibility for safety and use not in accordance with directions.