



TRIPLE ACTION  
LAWN CARE



# TRIPLE ACTION LAWN CARE

Prevents Crabgrass, Kills Existing Weeds And Feeds Your Lawn In One Step

Feeds up to **3** Months



18-0-1


Net 20 lbs  
Wt (9.07 kg)

Covers up to  
**8,300** Square Feet

Covers up to  
**8,300** Square Feet

Net 20 lbs  
Wt (9.07 kg)

18-0-1



Prevents And Kills Existing Weeds, And Feeds Your Lawn

KEEP OUT OF REACH OF CHILDREN  
**CAUTION**  
See back panel for additional Precautionary Statements and First Aid

Covers up to **8,300** Square Feet

Net Wt 20 lbs  
(9.07 kg)

18-0-1

ACTIVE INGREDIENTS:  
2,4-D ..... 0.675%  
Dicamba ..... 0.060%  
Dithiopyr ..... 0.121%  
Quinclorac ..... 0.346%  
OTHER INGREDIENTS: ..... 98.798%  
TOTAL ..... 100.000%  
  
THIS PRODUCT CONTAINS:  
13.50 lb 2,4-dichlorophenoxyacetic acid equivalent per ton (CAS 1928-43-4)  
1.20 lb 3,6-dichloro-o-anisic acid equivalent per ton (CAS 1918-00-9)  
2.42 lb dithiopyr per ton (CAS 97886-45-8)  
6.92 lb quinclorac per ton (CAS 84087-01-4)  
Isomer Specific by AOC Methods.

TRIPLE ACTION  
LAWN CARE



TRIPLE ACTION  
LAWN CARE

Feeds up to **3** Months



# TRIPLE ACTION LAWN CARE



**Prevents Crabgrass, Kills Existing Weeds And Feeds Your Lawn In One Step.**

## GUARANTEED FERTILIZER ANALYSIS

F825

Total Nitrogen (N) .....	18.00%
18.00% Urea Nitrogen	
Soluble Potash (K <sub>2</sub> O) .....	1.00%

**Derived from:** Polymer Coated Urea, Urea and Muriate of Potash.  
\*4.50% slow-release urea nitrogen from polymer coated urea.

**STOP** READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

## PRECAUTIONARY STATEMENTS

### Hazards to Humans and Domestic Animals

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wear long-sleeved shirt and long pants, shoes plus socks and gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

## FIRST AID

<b>If in Eyes</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If Swallowed</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on Skin or Clothing</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.

## ENVIRONMENTAL HAZARDS:

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

These chemicals (2,4-D and dicamba) have properties and characteristics associated with chemicals detected in groundwater. The use of these chemicals in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

### General Precautions and Restrictions:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed or for research purposes.
- This product is not sold, distributed, or registered within the state of New York. Do not use in the state of New York.

## Product Description

This product is formulated to feed your lawn and control most annual grasses and broadleaf weeds commonly found in home lawns. The product contains a lawn fertilizer and four different herbicides.

- The fertilizer contains slow release nutrients that may extend the availability for turfgrass uptake.
- The four herbicides combine to kill many different kinds of weeds at different stages of development.
- Grass weeds: Preemergent control of crabgrass, foxtails, and other annual grass weeds. Postemergent control of crabgrass.
- Broadleaf weeds: Preemergent control of chickweed, henbit, knotweed, shepherds-purse, spurge, woodsorrel, and other broadleaf weeds. Postemergent control of broadleaf weeds including dandelion, chickweed, thistle and clover.

## Where To Use

This product may be used on the following established turf species:

**Established Cool Season Turfgrass:** Kentucky bluegrass, rough bluegrass (Poa trivialis), perennial ryegrass, fine fescues, tall fescue, seashore paspalum

**Established Warm Season Turfgrass:** Bermudagrass, buffalograss, zoysiagrass

This product may NOT be used on: annual bluegrass (Poa annua), bahiagrass, bentgrass, carpetgrass, centipedegrass, dichondra, St. Augustine. Do not use this product on turfgrass where desirable clovers are present.

**Guidelines to follow:** The use of this product on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, chemical injury, disease, insects, pests, excessive temperatures, or lack of moisture may increase the chances for turfgrass injury. Allow turfgrasses weakened by these stresses to recover before application of this product.

## WEEDS CONTROLLED

When applied as directed, this product will control and/or suppress\* the following weeds:

### Annual Grass Weeds:

Annual bluegrass (Poa annua)	Barnyardgrass*	Foxtail (green, yellow, giant)
Annual ryegrass	Crabgrass (smooth, southern, large)	Goosegrass* (crowfootgrass)

\* for acceptable control of barnyardgrass and goosegrass, two applications made at a 6 to 10 week interval are needed

### Broadleaf Weeds:

Amaranth, purple	Burdock	Common mullein	Goldenrod	Gallinsoga, smallflower	Mallow, common
Amaranth, slender	Burweed, lawn (spurweed)	Corn Spurry	Creeping Jenny (creeping charlie)	Goldenrod	Mallow, dwarf
Amaranth, spiny	Buttercup, bulbous	Creeping Beggarweed	Cudweed	Ground ivy	Mallow, venice
Artichoke	Buttercup, creeping	Cupid-shaving Brush	Daisy, English	Gumweed	Marcela
Aster, heath	Buttonweed, common (poorjoe)	Carolina geranium (wild)	Daisy, oxeye	Hairy fleabane	Matchweed
Aster, purple	Buttercup, creeping	Chickweed, common	Dandelion	Hawkweed, meadow	Mexicanweed
Aster, white heath	Buttercup, creeping	Chickweed, mouseear	Dandelion, false (spotted catsear)	Hawkweed, mousear	Moneywort
Aster, white prairie	Catnip	Chickweed, sticky	Day flower	Hawkweed, orange	Morningglory, bigroot
Austrian fieldress	Chickweed, common	Chickweed, mouseear	Deadnettle	Hawkweed, yellow	Morningglory, ivyleaf
Bedstraw, catchweed	Chickweed, mouseear	Chickweed, sticky	Dichondra	Heartleaf drymary	Morningglory, mexican
Bedstraw, Northern	Chickweed, mouseear	Chickweed, sticky	Dock, broadleaf	Hemp	Morningglory, red
Bedstraw, pineywoods	Chickweed, mouseear	Chickweed, sticky	Dock, curly	Henbit	Morningglory, tall
Bedstraw, smooth	Chickweed, mouseear	Chickweed, sticky	Dock, smooth	Hemp	Morningglory, woolly
Beggaricks	Chickweed, mouseear	Chickweed, sticky	Dogbane	Hoary cress	Mustard, ball
Beggaricks, hairy	Chickweed, mouseear	Chickweed, sticky	Dogbane	Horsenettle	Mustard, black
Betony, Florida	Chickweed, mouseear	Chickweed, sticky	Dollarweed (Pennywort)	Horsenettle	Mustard, blue
Bindweed, Field	Chickweed, mouseear	Chickweed, sticky	Dove weed	Horsenettle	Mustard, hedge
Bindweed, Hedge	Chickweed, mouseear	Chickweed, sticky	Elderberry	Horsenettle	Mustard, indian
Bitter Wintercress	Chickweed, mouseear	Chickweed, sticky	False sunflower	Horsenettle	Mustard, tansy
Bitterweed	Chickweed, mouseear	Chickweed, sticky	False sunflower	Horsenettle	Mustard, tumble
Black medic	Chickweed, mouseear	Chickweed, sticky	Falsiflax	Horsenettle	Mustard, white
Black-eyed Susan	Chickweed, mouseear	Chickweed, sticky	Fiddleneck	Horsenettle	Mustard, wormseed
Blood flower milkweed	Chickweed, mouseear	Chickweed, sticky	Filaree, redstem	Horsenettle	Nettle, stinging (burning)
Blue lettuce	Chickweed, mouseear	Chickweed, sticky	Filaree, whitestem	Horsenettle	Nettle, tall
Brass buttons	Chickweed, mouseear	Chickweed, sticky	Fleabane, daisy (annual)	Horsenettle	Nettle, wood
Bristly oxtongue	Chickweed, mouseear	Chickweed, sticky	Fleabane, rough	Horsenettle	Parsley-piert
Broomweed	Chickweed, mouseear	Chickweed, sticky	Gallinsoga, hairy	Horsenettle	Parsnip
Bullnettle	Chickweed, mouseear	Chickweed, sticky		Horsenettle	Pearwort

## WEEDS CONTROLLED - continued

Pennycress (frenchweed)	Prairie Sunflower	Spanish needles	Thistle, plumeless	Wild garlic
Pepperweed	Prickly lettuce (compass plant)	Speedwell, common	Thistle, russian	Wild lettuce
Pepperweed, perennial	Prickly sida	Speedwell, corn	Thistle, scotch	Wild mustard (yellow rocket)
Pepperweed, virginia (peppergrass)	Puncturevine	Speedwell, ivyleaf	Thistle, yellow (horrible)	Wild onion
Pigweed, prostrate	Purslane, common	Speedwell, persian	Trailing crownvetch	Wild parsnip
Pigweed, redroot	Pusley, brazil	Speedwell, purslane	Velvetleaf	Wild radish
Pigweed, russian	Pusley, florida	Speedwell, slender	Vervain, blue	Wild rape
Pigweed, smooth	Pusley, large flower	Speedwell, snow	Vervain, hoary	Wild strawberry
Pigweed, tumble	Ragweed, bur	Speedwell, thymeleaf	Vervain, prostrate	Wild sweet potato
Pineapple weed	Ragweed, common	Spurge, prostrate	Vervain, tall	Wild violet
Plains coreopsis (tickseed)	Ragweed, landleaf	Spurge, spotted	Vetch, bird	Woodsorrel, creeping
Plantain, blackseed	Ragweed, western	Spurweed	Vetch, hairy	Woodsorrel, violet
Plantain, bracted	Rough cinquefoil	Star of Bethlehem	Vetch, milk	Woodsorrel, yellow (Oxalis)
Plantain, broadleaf	Scarlet pimpernel	Strawberry, India mock	Vetch, narrowleaf	Woolly croton
Plantain, buckhorn	Shepherds-purse	Tansy ragwort	Vetch, wild	Wormseed
Plantain, hoary	Smartweed, ladysthumb	Tanweed	Virginia creeper	Yarrow, common
Plantain, narrowleaf	Smartweed, pale	Thistle, blessed	Western clematis	Yarrow, western
Plantain, slender	Smartweed, pennsylvania	Thistle, bull	Western salsify	Yarrow, yellow
Plantain, woolly	Smooth chaff-flower	Thistle, canada	Wild aster	
Poison ivy	Sorel, red (sheep)	Thistle, floodman (prairie)	Wild buckweed	
Poison oak	Southistle, annual	Thistle, musk	Wild carrot	
Pokeweed	Southistle, spiny	Thistle, platte	Wild four-o'clock	

## When To Apply

This product may be applied as a single application or as split applications in the spring, early summer, or fall.

For annual grasses such as crabgrass (smooth, southern and large) and foxtail (green, yellow, and giant): Apply in the spring and early summer. For best results, apply this product from 2 weeks before weed germination until the weed is 4 inches tall. For dense populations, make a second application no sooner than 30 days after the first application.

For broadleaf weed control: For best results, apply when broadleaf weeds are young and actively growing with adequate soil moisture and favorable temperatures (50° to 80°F).

## How To Apply

- This product needs to stick to the leaves of broadleaf weeds and should be applied when the grass is wet from dew or rain.
- Poor results will occur if applied to dry lawns. When dry conditions exist, irrigate (or water) the turfgrass 24 hours before application.
- For best results, irrigate after the application. However do not irrigate sooner than 24 hours after application.
- Apply this product when the wind is calm.
- Do not spread granules onto flowers, vegetables, and ornamental shrubbery.
- Turf species listed on this label can be seeded, sodded, sprigged, or plugged into the treated areas 4 weeks after the application of this product.
- When treating newly planted areas, wait approximately 6 weeks after seedling emergence before applying this product.
- Delay mowing 2 days before and until 3 days after the application of this product. Additional stress from low mowing heights may increase the possibility of turf injury.
- Clippings from the first three mowings should be left on the treated area.
- Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

## How Much To Use

Use a drop or rotary spreader designed to apply granular fertilizers. Avoid skips or excessive overlaps as poor weed control or turfgrass injury may result.

Apply 2.4 pounds of product per 1,000 sq. ft. per application to established turfgrass.

The maximum broadcast application rate is 2.4 pounds of product per 1,000 sq. ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum annual (seasonal) rate is 4.8 pounds of product per 1,000 sq. ft.

## CALIBRATION INSTRUCTIONS:

For 2.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 420 square feet and adjust to apply at the rate of 1 pound of product per 420 square feet.

## STORAGE and DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Keep in original container in a cool, dry area, away from extreme heat and cold.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container (bag). Completely empty bag into application equipment. If empty: Place bag in trash or offer empty bag for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.html>

EPA Reg. No. 2217-926-59144

EPA Est. No. 59144-GA-1 (A), 32802-MO-1 (B), 37351-CO-1 (C), 8378-IN-1 (D), 9404-FL-1 (E), 961-PA-01 (F), 961-IL-01 (G), 35497-OR-1 (H) 49991-TX-2 (Q), 69759-OR-002 (R), 7401-TX-01 (S)

Letter in parenthesis is first letter of lot number.

## LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Expert Gardener  
Triple Action Lawn Care 20lbs



Accugreen, EdgeGuard and Scotts are registered trademarks of OMS Investments, Inc. SpeedyGreen is a registered trademark of Everris International B.V. EV-N-SPRED is a registered trademark of Earthway Products, Inc.

Distributed and Guaranteed by:  
Gro Tec, Inc., P.O. Box 290, Madison, GA 30650

Expert Gardener is a registered trademark of  
Walmart Inc. Bentonville, AR 72716

**Net Wt 20 lbs (9.07 kg)**