

LAWN WEED CONTROL

Kills Dandelions, Clovers & Other Major Lawn Weeds

Kills the Weeds, Not the Lawn

Money Back Guarantee


See back panel for details

Covers up to 5,000 Square

LAWN WEED

^{Net} 14 lbs (6.35 kg)





KEEP OUT OF REACH OF CHILDREN
CAUTION

See back panel for complete Precautionary
Statements and Directions for Use, including
First Aid, and Storage and Disposal

Covers up to Square Feet

Net 14 lbs (6.35 kg)

ACTIVE INGREDIENTS:
2, 4-D, 2-ethylhexyl ester
1.1083
Mecoprop-p acid
0.1679
Dicamba acid
0.0719
OTHER INGREDIENTS
98.6549
TOTAL
100.0009
THIS PRODUCT CONTAINS:
14.70 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.735%
CAS 1928-43-4.
3.34 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.167%.
CAS 16484-77-8

Isomer Specific by AOAC Methods



(6.35 kg)

Covers up to 5,000 Square

LAWN WEED CONTROL



LAWN WEED CONTROL







LAWN WEED CONTROL

Kills Dandelions, Clovers & Other Major Lawn Weeds



Multiply the length by the width of your yard and then subtract the area of your house to find the square footage you need to cover.



Apply when broadleaf weeds are actively growing in Spring or early Fall. Apply to a wet lawn.



Always apply with a rotary or drop spreader to distribute granules evenly (see chart below). Never apply by hand.





No rain or water for 1 to 2 days after application. After 2 days, water thoroughly.



STOP READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

If in Eyes

Hold eye open and rinse slowly and gently with water for 15-20 minutes.

 Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye Call a poison control center or doctor for treatment advice

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this product, contact the National Pesticide Information Center, 1-800-858-7378, Monday - Friday, 8:00 am to 12:00 pm PST You may also contact the National Poison Control Center, 1-800-222-1222, day or night, for emergency medical treatment information. FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is a combination of three proven weed killers-2,4-D, mecoprop-p, and dicamba. This herbicide controls a wide range of eeds, including dandelion, chickweed, knotweed, plantain, henbit, and spurge.

- · Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- · Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes. Do not use on dichondra, carpetgrass, 'Floratam' St. Augustinegrass, creeping bentgrass, or turfgrass where desirable clove
- Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees. Do not exceed specified dosage for any area and do not apply this product within the dripline of, or over surface roots of trees, shrubs, and other
- · Do not use this product on or near desirable plants and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

USE PRECAUTIONS:

 Do not apply this product to newly seeded grasses until well-established or after the second or third mowing. • Reseeding or overseeding areas may occur three (3) weeks after the application of this product.

For use on Kentucky Bluegrass, Fescue spp., Perennial Ryegrass, common Bermudagrass, Bahiagrass, Zoysiagrass, St. Augustinegrass, Fescue spp., Perennial Ryegrass, Common Bermudagrass, Bahiagrass, Zoysiagrass, St. Augustinegrass, Fescue spp., Perennial Ryegrass, Common Bermudagrass, Bahiagrass, Zoysiagrass, St. Augustinegrass, Fescue spp., Perennial Ryegrass, Common Bermudagrass, Bahiagrass, Zoysiagrass, St. Augustinegrass, Fescue spp., Perennial Ryegrass, Common Bermudagrass, Bahiagrass, Zoysiagrass, St. Augustinegrass, Fescue spp., Perennial Ryegrass, Common Bermudagrass, Bahiagrass, Zoysiagrass, St. Augustinegrass, Fescue spp., Perennial Ryegrass, Common Bermudagrass, Bahiagrass, Zoysiagrass, St. Augustinegrass, Fescue spp., Perennial Ryegrass, Common Bermudagrass, Bahiagrass, Zoysiagrass, St. Augustinegrass, Fescue spp., Fescue spp.,Centipedegrass, hybrid Bermudagrass, and Colonial Bentgrass. Apply 2.8 pounds of product per 1,000 square feet of turfgrass. The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application. The maximum number of broadcast applications is 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per

WHEN TO USE

Apply when Broadleaf weeds are actively growing in Spring or early Fall.

Apply when the air is calm to ensure uniform coverage and avoid spreading granules onto flowers, vegetables and ornamental shrubbery. Possible retreatment may be necessary if lawn is heavily overrun by weeds, if adverse low moisture conditions prevail or if weeds are in a state of poor growth.

APPLICATION INSTRUCTIONS

- 1. Mow lawn to normal height 1 to 2 days before application. 2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).
- 3. Apply when Broadleaf weeds are young and actively growing, preferably in the morning when dew is on the grass. At the time of application, moisture on the weed leaves from dew, rainfall or irrigation/watering aids in control. If grass is not moist at time of application, sprinkle lightly with water before applying product.
- 4. Avoid applying when temperatures are consistently above 90°F 5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

SPREADER SETTINGS **BROADCAST SPREADER** xpert Gardener® Broadcast xpert Gardener® Deluxe Broadcast cotts® SpeedyGreen 3.7 Scotts® Basic cotts® Lawn Pro 3.75 cotts® EdgeGuard® Deluxe 3.7 cotts® EdgeGuard® Mini 3.75 **DROP SPREADER** Scotts® Accuareen®

HOW TO APPLY - SPREADERS & CALIBRATION

Always apply with a rotary or drop spreader to distribute the granules evenly. Never apply by hand. **COVÉRAGE**: To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One pag covers up to 5,000 square feet.

IMPORTANT: It is important to follow directions as indicated on this quality lawn care product as overapplication may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult Owner's

Manual. Never apply by hand. For rotary spreaders: Operate at intervals corresponding to the swath width.

For drop spreaders: Overlap the wheel tracks to ensure complete coverage. For best results, apply this product uniformly and avoid streaks, skips or overlaps. Walk at a steady pace to ensure even coverage. Close the spreader opening when turning to change directions or when stopping

Expert Gardener Lawn Weed Control II 0-0-1 F525

GUARANTEED ANALYSIS Soluble Potash (K₃O). 1.0% Derived from: Muriate of Potash.

CALIBRATION INSTRUCTIONS AND SPREADER SETTINGS

Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended rate under actual application conditions.

To provide proper distribution, calibrate spreader before application. BROADCAST SPREADERS: For best results, apply in a circular path starting on the outside and working toward the center

overlapping slightly to ensure complete coverage **DROP SPREADERS:** Apply 2 strips along each end. Then apply along length of lawn, overlapping slightly and shutting off the

Horseweed

Lespedeza

Mallow, comm

Johnny-Jumpup violet Knawel

Knotweed, prostrate

WEEDS CONTROLLED

Amaranth, purple Amaranth, slende Clover, white Cockle Aster, heath Aster, white heath Cocklebur Aster, purple Aster, white prairie Artichoke Austrian fieldcress Cocklebur, oriental Cocklebur, spiny Common mulleir Bedstraw, Catchweed Bedstraw, smooth Corn Spurry Creeping Beggarweed Bedstraw, Nonthern Creeping Jenny (creeping charlie) Bedstraw, pineywoods Beggartricks, hairy Betony, Florida Cupid shaving Brush Daisy, English Bindweed, Field Bindweed, Hedge Dandelion, false Bitter Wintercress Bittercress, hairy Day flowe Black-eyed Susan Dichondra Dock, broadleaf Dock, curly Black Medic Blood flower milkweed Blue lettuce Dogbane Dogfennel Dollarweed (Pennywort) Bristly oxtongue Dove weed Elderberry Falseflax Buckhorn False sunflower Burclover Fiddleneck Filaree, redstem Burdock Burweed, lawn (spurweed) Buttercup, bulbous Filaree, whitestem Fleabane, daisy (annual) Fleabane, rough Buttercup, creeping Buttonweed, comm Buttonweed, Virginia Galinsoga, hairy Carolina geranium (wild) Ground ivy Catsear, spotted (false dandelion) Groundsel Hairy fleabane Chickweed, commor Chickweed, mouseear Hawkweed, meadow Hawkweed, mouseea Chickweed, sticky Hawkweed, orange Cinquefoil Clover, crimsor Hawkweed, yellow

Marcela Matchweed Moneywort orningglory, i Morningglory, Mexican Morningglory, red Morningglory, tall Morningglory, woolly Mustard, black Mustard, blue Mustard, Indian Mustard, tansy Mustard, tumble Mustard, white Mustard, wormseed Nettle, stinging (burning) Nettle, tall Parsley-piert Parsnip Pennycress (frenchweed Pepperweed, perennial Pepperweed, virginia (Pepp Pigweed, prostrate Pigweed, redroot Pigweed, russian Heartleaf drymary Pigweed, tumble Pineapple weed Plains coreopsis (tickseed) Hoary cress Plantain, blackseed Plantain, bracted

Plantain, buckhorn Plantain, hoary Plantain, narrowlea Plantain, slender Plantain, woolly Poison ivy Poison oak Prickly lettuce (compass plant) Prostrate knotweed Puncturevine Purslane, commo Pusley, florida Ragweed, commo Ragweed, lancleaf Ragweed, western Redstem filaree Rough cinquefoil Scarlet pimpernel Shepherd's purse Smartweed, ladysthumb Smartweed, pale Smartweed, pennsylvania Smooth chaff-flower Sorrel, red (sheep) Sowthistle, annua Sowthistle, spiny Spanish needles peedwell, bi-lobed Speedwell, corn Speedwell, ivyleaf Speedwell, persian Speedwell, purslane Speedwell, slender Speedwell, thymeleaf Spiny amaranth Spurge, spotted Spurge, prostrate Star of Bethlehem

Plantain, broadleaf

Thistle, blessed Thistle, bull Thistle, canada Thistle, flodman (prairie) Thistle, musk Thistle, platte Thistle, plumeless Thistle, russian Thistle, scotch Thistle, wavyleaf Thistle, yellow (horrible) Trailing crownvetch Velvetleaf Vervain, blue Vervain, hoary Vervain, prostrate Vetch, hairy Vetch, milk Vetch, narrowlead Vetch, wild Virginia creeper Western clemat Western salsify Wild aster Wild carrot Wild four-o-clock Wild lettuce Wild marigold Wild mustard (yellow rocket) Wild onion Wild parsnip Wild radish Wild rape Wild strawberry Wild violet Woodsorrel, creeping Woodsorrel, yellow (Oxalis) Woolly croton Yarrow, commor Yarrow, yellow Strawberry, India mock

Clover, strawberry STORAGE AND DISPOSAL

Clover, alyce

Clover, red

Do not contaminate water, food or feed by storage and disposal. **PESTICIDE STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container: Do not reuse or refill this container. If empty: Place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place any unused product down any indoor or outdoor drain

 $Keep\ this\ product\ off\ sidewalks,\ patios,\ driveways,\ etc.\ This\ fertilizer\ may\ stain\ concrete\ and\ painted\ surfaces.\ If\ it\ comes\ in\ contact$ with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders, which scatter fertilizer beyond lawn edge and onto concrete. EPA Reg. No. 2217-1047-59144

EPA Est. No, 59144-GA-1 (A), 32802-MO-1 (B), 37351-CO-1(C), 8378-IN-1 (D), 9404-FL-1 (E), 961-PA-01 (F), 961-IL-01(G), 35497-OR-1 (H), 49991-TX-2 (Q), 69759-OR-002 (R), 7401-TX-01 (S)

Letter in parenthesis is first letter of lot number. When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials. Information regarding the contents and levels of metals in this product is available on the Internet at

http://www.aapfco.org/metals.html **LIMITED WARRANTY AND DISCLAIMER**

FOR USE ONLY AS DIRECTED. THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUEN-TIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS **PRODUCT.** If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Great Quality. Great Price. Guaranteed.

* Our guarantee is our promise that you'll be fully satisfied with the quality of every Expert Gardener $^{ ext{@}}$ product. If for any reason you aren't happy, we'll replace it or return your money – whichever you prefer. All you need to do is return the empty packaging. It's that simple. Guaranteed. Guaranteed within 1 year of the date of purchase.

Questions or comments? Call 855-481-2977.

Expert Gardener Lawn Weed Control II 14lbs



Accugreen, EdgeGuard and Scotts are registered trademarks of OMS Investments, Inc SpeedyGreen is a registered trademark of

Distributed by: Gro Tec, Inc. P.O. Box 290, Madison, GA 30650 Expert Gardener is a registered trademark of




LAWN WEED CONTROL.

Kills Dandelions, Clovers & Other Major Lawn Weeds

Kills the Weeds, Not the Lawn

Money Back Guarantee

See back panel for details



KEEP OUT OF REACH OF CHILDREN CAUTION See back panel for complete Precautionary Statements and Directions for Use, including First Aid, and Storage and Disposal

AWN WEED

NTROL

Covers up to 5,000 Square Feet

Net 14 lbs (6.35 kg)

2, 4-D, 2-ethylhexyl este 3.34 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.167%. 1.42 lb 3, 6-dichloro-o-anisic acid equivalent per ton or 0.071% **Isomer Specific by AOAC Methods**



LAWN WEED



LAWN WEED CONT

Expert Gardener is a registered trademark of Walmart Inc. Bentonville, AR 72716 Distributed by: Gro Tec, Inc. P.O. Box 290, Madison, GA 30650

SpeedyGreen is a registered trademark of trademarks of OMS Investments, Inc. Accugreen, EdgeGuard and Scotts are registered



Questions or comments? Call 855-481-2977.

Guaranteed within I year of the date of purchase. the empty packaging. It's that simple. Guaranteed. money – whichever you prefer. All you need to do is return reason you aren't happy, we'll replace it or return your vith the quality of every Expert Gardener $^{\scriptscriptstyle \odot}$ product. It for any * Our guarantee is our promise that you'll be fully satisfied Great Quality. Great Price. Guaranteed. agreements at the point of sale or elsewhere.

Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or the point of purchase, and the purchase price will be refunded in full. The terms of this PRODUCT. If these terms are not acceptable, return this product unopened immediately to TIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUEN-EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF FOR USE ONLY AS DIRECTED. THE MANUFACTURER NEITHER MAKES NOR INTENDS

LIMITED WARRANTY AND DISCLAIMER http://www.aapfco.org/metals.html

Officials. Information regarding the contents and levels of metals in this product is available on the Internet at When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Letter in parenthesis is first letter of lot number.

49991-TX-2 (Q), 69759-OR-002 (R), 7401-TX-01 (S) Eby Eşt. No, 59144-GA-1 (A), 32802-MO-1 (B), 37351-CO-1(C), 8378-IN-1 (D), 9404-FL-1 (E), 961-PA-01 (F), 961-IL-01(G), 32497-OR-1 (H), EPA Reg. No. 2217-1047-59144 impossible to prevent. Be careful when using rotary spreaders, which scatter fertilizer beyond lawn edge and onto concrete.

with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be Keep this product off sidewalks, patios, driveways, etc. This fertilizer may stain concrete and painted surfaces. If it comes in contact down any indoor or outdoor drain.

recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place any unused product AND CONTAINER HANDLING: Nonrefillable container: Do not reuse or refill this container. If empty: Place in trash or offer for bags containing unused product in a locked storage area inaccessible to children and domestic animals. **PESTICIDE DISPOSAL** Do not contaminate water, food or feed by storage and disposal. PESTICIDE STORAGE: Store in original container and keep any **STORAGE AND DISPOSAL** Strawberry, India mock

Clover red	Hoary cross	haasyseld nietnel@	Strawherry India mock	
Clover, rabbit foot	Jidn9H	Plains coreopsis (tickseed)	Star of Bethlehem	Yarrow, yellow
Clover, hop	Hemp	Pineapple weed	zbnxweed	Yarrow, western
Clover, alyce	Heartleaf drymary	Pigweed, tumble	Spurge, prostrate	Хаггом, соттоп
Clover, crimson	Healall	Pigweed, smooth	Spurge, spotted	Wormseed
Cinquefoil	Hawkweed, yellow	Pigweed, russian	дыпу этагалт	Woolly croton
Chicory	намкмеед, огапде	Pigweed, redroot	Speedwell, thymelear	Woodsorrel, yellow (Oxalis)
Chickweed, sticky	Намкмеед, тоизееаг	Pigweed, prostrate	ypeedwell, snow	Woodsorrel, violet
Chickweed, mouseear	нэмкмеед, теадом	Pepperweed, virginia (Peppergrass)	Speedwell, slender	Woodsorrel, creeping
Сыскweed, соттоп	Hairy fleabane	Pepperweed, perennial	Speedwell, purslane	Wild violet
Chamber bitter	remweed	Pennycress (frenchweed)	Speedwell, persian	Wild sweet potato
Catsear, spotted (false dandelion)	expunosej	Pearlwort	Speedwell, ivyleat	Wild strawberry
dinje	conuq ivy	Parsnip	Speedwell, corn	Wild rape
Carpetweed	Goldenrod	Parsley-piert	Speedwell, common	Asiber bliW
Carolina geranium (wild)	Galinsoga, smallflower	Nettle, wood	Speedwell, bi-lobed	Wild parsnip
Buttonweed, Virginia	valinsoga, hairy	Nettle, tall	səlbəən akineq	noino bliW
Buttonweed, common (poorjoe)	Heabane, rough	Nettle, stinging (burning)	Sowthistle, spiny	Wild mustard (yellow rocket)
Buffercup, creeping	Heabane, daisy (annual)	Mustard, wormseed	Sowthistle, annual	Wild marigold (wellow rocket)
Buttercup, bulbous	Filaree, whitestem	Mustard, white	Sorrel, red (sheep)	Wild lettuce
Buttercup hulbous	Filaree, redstem	Mustard, tumble	Smoothdock	Wild garlic
Burdock Burdock	Fiddleneck	Mustard, tansy	Smooth chaff-flower	Wild four-o-clock
Burdock	False sunflower	Mustard, Indian	Smartweed, pennsylvania	Wild Carrot
Bullnettle	X6lf92l67 Falca cultura a sle3	Mustard, hedge	Smartweed, pale	Wild buckwheat
	Elderberry yestealay		Smartweed, ladysthumb	
вискроги Вгоотмееа		Mustard, blue	Shepherd's purse	Wild aster
Bronweed Bronweed	Dollarweed (Pennywort) Dove weed	Mustard, black	Scarlet pimpernel	Western salsify
Brassbuttons Bristly oxtonnie		Mustard, ball		Vestern clematis
	Dogfennel	Mugwort	Rough cinquefoil	Virginia creeper
Blue lettuce	Dogbane	Morningglory, woolly	Redstem filaree	Vetch, wild
Blood flower milkweed	Dock, curly	Meringglory, tall	Ragweed, western	Vetch, narrowleaf
Black Medic	Dock, broadleaf	Morningglory, red	Ragweed, lancleaf	Vetch, milk
Black-eyed Susan	Dichondra	Morningglory, Mexican	вадмеед, соттоп	Vetch, hairy
Bitterweed	Deadnettle	Momingglory, ivyleaf	Ragweed, bur	Vetch, bird
Bittercress, hairy	рэу Помег	Morningglory, bigroot	Pusley, large flower	Vervain, tall
Bitter Wintercress	əslət ,fooiləbneQ	Moneywort	Pusley, florida	Vervain, prostrate
Bindweed, Hedge	Dandelion, common	Mexicanweed	Pusley, brazil	Vervain, hoary
Bindweed, Field	Daisy, oxeye	beewdətsM	Purslane, common	Vervain, blue
Betony, Florida	Daisy, English	Marcela	Puncturevine	łseltevleV
Beggartricks, hairy	pəəwpn	Mallow, venice	Prostrate knotweed	Trailing crownvetch
Bedstraw, pineywoods	dsura Bruish biqu.	Mallow, dwarf	Prickly sida	Thistle, yellow (horrible)
Bedstraw, Nonthern	Creeping Jenny (creeping charlie)	Mallow, common	Prickly lettuce (compass plant)	Thistle, wavyleaf
Bedstraw, smooth	Creeping Beggarweed	Mallow, bristly	Prairie Sunflower	Thistle, scotch
Bedstraw, Catchweed	сот Spurry	Mallow, alkali	Pokeweed	Thistle, russian
szərəbləfi neirtzuA	niəllum nommo)	Геsbedeza	Poison oak	Thistle, plumeless
Artichoke	Cocklebur, spiny	Lambsquarters	Yvi nosio9	Thistle, platte
Aster, white prairie	Cocklebur, oriental	Kochia	Plantain, woolly	Thistle, musk
Aster, purple	Cocklebur	Knotweed, prostrate	Plantain, slender	Thistle, flodman (prairie)
Aster, white heath	Cockle	Knawel	Plantain, narrowleaf	Thistle, canada
Aster, heath	Clover, white	təloiv quqmul-ynndol	Plantain, hoary	Thistle, bull
Amaranth, slender	Clover, white sweet	bəəwnosmil	Plantain, buckhorn	Thistle, blessed
Amaranth, purple	Clover, yellow sweet	Horseweed	Plantain, broadleat	lanweed

WEEDS CONTROLLED

overlapping slightly to ensure complete coverage. **DROP SPREADERS:** Apply 2 strips along each end. Then apply along length of lawn, overlapping slightly and shutting off the spreader before each turn at edge of border strips. recommended rate under actual application conditions.

To provide proper distribution, calibrate spreader before application.

To provide proper distribution, calibrate spreader before application.

To provide proper distribution, calibrate spreader before application. Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the CALIBRATION INSTRUCTIONS AND SPREADER SETTINGS

Derived from: Muriate of Potash. Soluble Potash (KرO,). %0.1 **GUARANTEED ANALYSIS** Expert Gardener Lawn Weed Control | 0-0-1 F525

directions or when stopping. steady pace to ensure even coverage. Close the spreader opening when turning to change For drop spreaders: Overlap the wheel tracks to ensure complete coverage. Walk at a For rotary spreaders: Operate at intervals corresponding to the swath width Manual. Never apply by hand. treated. Be sure to check spreader before beginning, adjusting if necessary. Consult Owner's Differences may occur due to condition of spreader, speed at which applied and roughness of area overapplication may cause injury to your lawn. Spreader settings are approximate.

bag covers up to 5,000 square feet. IMPORTANT: It is important to follow directions as indicated on this quality lawn care product as Always apply with a rotary or drop spreader to distribute the granules evenly. Never apply by hand. COVERAGE: To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One HOW TO APPLY - SPREADERS & CALIBRATION

DROP SPREADER cotts" EdgeGuard" Mini cotts® EdgeGuard® Deluxe Scotts® Lawn Pro cotts® Basic cotts SpeedyGreen Expert Gardener® Deluxe Broadcast xpert Gardener Broadcast **BROADCAST SPREADER SPREADER SETTINGS**

″n991gu⊃⊃A ®stfo⊃∂

5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made. 4. Avoid applying when temperatures are consistently above 90°F. application, moisture on the weed leaves from dew, rainfall or irrigation/watering aids in control. If grass is not moist at time of application, sprinkle lightly with water before applying product. 3. Apply when Broadleaf weeds are young and actively growing, preferably in the morning when dew is on the grass. At the time of 1. Mow lawn to normal height 1 to 2 days before application to sustain moisture until the next watering (see step 5). 2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).

SNOITOURTSNI NOITADIJ99A

Possible retreatment may be necessary if lawn is heavily overrun by weeds, if adverse low moisture conditions prevail or if weeds are in a Apply when Broadleaf weeds are actively growing in Spring or early Fall.

Apply when the air is calm to ensure uniform coverage and avoid spreading granules onto flowers, vegetables and ornamental shrubbery. WHEN TO USE

applications is 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per Centipedegrass, hybrid Bermudagrass, and Colonial Bentgrass. Apply 2.8 pounds of product per 1,000 square feet of furfgrass. The maximum number of product per 1,000 sq.ft. per application. The maximum number of product set For use on Kentucky Bluegrass, Fescue spp., Perennial Ryegrass, common Bermudagrass, Bahiagrass, Zoysiagrass, Augustinegrass, YJ99A OT HOUM WOH • Reseeding or overseeding areas may occur three (3) weeks after the application of this product. • Do not apply this product to newly seeded grasses until well-established or after the second or third mowing.

USE PRECAUTIONS: over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not Bo not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lesterams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your jawn, and

(rhizomes) within the drip line of the roots of desirable trees and shrubs, since injury may result. • Do not use this product on or near desirable plants and avoid spreading granules onto exposed root systems or adventitious shoots Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees. Do not exceed specified dosage for any area and do not apply this product within the dripline of, or over surface roots of trees, shrubs, and other • Do not use on dichondra, carpetgrass, Floratam' St. Augustinegrass, creeping bentgrass, or turfgrass where desirable clovers

area during application. Do not allow people or pets to enter the treated area until dusts have settled. • Do not apply this product in a way that will contact any person or pet, either directly or through drift, Keep people and pets out of the This product is a combination of three proven weed killers—2,4-D, mecoprop-p, and dicamba. This herbicide controls a wide range of

• Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, and spurge. PRODUCT DESCRIPTION:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. **DIRECTIONS FOR USE** the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm **ENVIRONMENTAL HAZARDS** nformation. FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300. You may also contact the National Poison Control Center, 1-800-222-1222, day or night, for emergency medical treatment tion on this product, contact the National Pesticide Information Center, 1-800-858-7378, Monday - Friday, 8:00 am to 12:00 pm PST.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where

Call a poison control center or doctor for treatment advice. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

Hold eye open and rinse slowly and gently with water for 15-20 minutes. If in Eyes **GIA T2811**

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Hazards to Humans and Domestic Animals PRECAUTIONARY STATEMENTS


STOP READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

2 days, water thoroughly. days after application. After No rain or water for 1 to 2



below). Mever apply by hand. dranies evenly (see chart drop spreader to distribute Always apply with a rotary or






Apply to a wet lawn.

in Spring or early Fall.

weeds are actively growing

Apply when broadleaf



footage you need to cover. house to find the square subtract the area of your width of your yard and then Multiply the length by the





& Other Major Lawn Weeds

Kills Dandelions, Clovers LAWN WEED CONTROL.










ESKO:


PressMatch: 13












50%.... **77.5**%



30%.... 56.8%

50%.... 75.7%

Process only

