



FIFRA Section 24(c)
SPECIAL LOCAL NEED LABEL

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON

Ziram 76DF Fungicide

GROUP M3 FUNGICIDE

EPA Reg. No. 70506-173

EPA SLN No. OR-180001

For control of Eastern Filbert Blight on Filberts (Hazelnuts)

This label is valid until December 31, 2028 or until otherwise amended, withdrawn, cancelled, or suspended.

ACTIVE INGREDIENT:

Ziram (Zinc dimethyldithiocarbamate) 76.0%

OTHER INGREDIENTS: 24.0%

TOTAL..... **100.0%**

Contains 0.76 lbs ai of ziram per pound

**KEEP OUT OF REACH OF CHILDREN
PELIGRO DANGER**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

ENVIRONMENTAL HAZARDS

Follow all applicable directions, restrictions, and precautions in the Environmental Hazards section of the label affixed to the main container of Ziram 76DF Fungicide (EPA Reg. No. 70506-173).

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This SLN label and the label affixed to the main container of Ziram 76DF Fungicide must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the label affixed to the main container of Ziram 76 DF Fungicide.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

USE INSTRUCTIONS

Crop	Lbs. Per Acre	Disease	Application Instructions
Filberts	3 – 8	Eastern Filbert Blight (EFB)	Apply at bud break and continue at 2 week intervals until early May. Thorough coverage of all branches is essential. Use the higher rate if disease pressure is high and/or if crop canopy is dense. Do not apply more than 5 applications per crop cycle. Do not apply more than 30 lbs (22.8 lbs a.i.) per crop cycle. A crop cycle is defined as pre-bloom through postharvest. For ground applications, apply in a minimum of 20 gallons of spray per acre. For aerial applications, apply in a minimum of 10 gallons of spray per acre. For both ground and aerial applications, always use sufficient water to provide thorough coverage.

RESTRICTIONS

Do not apply this product through any type of irrigation system.
Do not graze or feed cover crops from treated orchards.

Rev. 02/09/2024