

**FIFRA Section 24(c) Special  
Local Need Label**

**EPA Reg. No. 279-3220  
EPA SLN No.: OR-250005**

**FOR DISTRIBUTION AND USE  
ONLY IN THE STATE OF  
OREGON**



**FOR CONTROL OF HAIRCAP MOSS AND SUSCEPTIBLE WEEDS IN  
CRANBERRIES**

<p>This label for Spartan 4F Herbicide is valid until December 31, 2027 or until otherwise amended, withdrawn, cancelled, or suspended.</p>
<p><b>IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.</b></p>
<p><b>THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.</b></p>

<b>Active Ingredient:</b>	
Sulfentrazone: .....	39.6%
<b>Other Ingredients:</b> .....	<u>60.4%</u>
<b>Total:</b>	100.0%

Contains 4 lbs of active ingredient per gallon.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

**Environmental Hazards**

Follow all applicable directions, restrictions and precautions in the Environmental Hazards section of the label affixed to the main container of Spartan 4F Herbicide.

**DIRECTIONS FOR USE**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and the label affixed to the main container of Spartan 4F Herbicide must be in the possession of the user at the time of pesticide application.
- Read this entire SLN label and the entire label affixed to the main container of Spartan 4F Herbicide.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements and precautions on this SLN label and the label affixed to the main container of Spartan 4F Herbicide (EPA Reg No. 279-3220).

## **Application Information**

**Proper handling instructions:** Spartan 4F Herbicide may not be mixed or loaded within 50 feet of any wells (including abandoned wells and drainage wells), sinkholes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pads or properly diked mixing/loading areas. Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or washwater, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specific minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment. Product must be used in a manner that will prevent back siphoning in wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

### **Application Timing: Dormant Spring Application**

Spartan 4F Herbicide can be applied after removal of winter flood and prior to cranberry plants breaking dormancy for control of haircap moss and other susceptible weed species. For preemergence control of susceptible weeds, apply Spartan 4F Herbicide at 8 to 12 fl oz/A (0.25 - 0.375 lb ai/A) before cranberry bud scales begin to loosen, or stunting may occur.

### **Application Methods**

**Chemigation Application:** Spartan 4F Herbicide may be applied via injection through sprinkler chemigation systems. **Spartan 4F Herbicide is active in the soil.** Best results are obtained when the soil is moist at the time of application. Additional irrigation may be applied after injection if needed. Follow all applicable directions, precautions, and restrictions in the Chemigation Application section of the label affixed to the main container of Spartan 4F Herbicide.

**Broadcast Boom Application:** Spartan 4F Herbicide may be applied by traditional boom application at broadcast rates listed above using a minimum of 15 GPA carrier volume. After application, run sprinkler irrigation and apply 0.1 – 0.25" of water volume per acre to remove Spartan 4F Herbicide from the foliage and provide soil incorporation.

The directions on this SLN label are based upon the interactive effects of Spartan 4F Herbicide and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under Rotational Crop Restrictions, Conditions of Sale and Limitations of Warranty and Liability, and any other section on the label affixed to the main container of Spartan 4F Herbicide pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been evaluated under treatment with Spartan 4F Herbicide. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on Spartan 4F Herbicide under specific local conditions.

### **Precautions**

- Application to cranberry plants at the cabbage head growth stage has shown limited phytotoxicity, usually temporary, in the form of stunting.

## **Restrictions**

- To minimize risk of injury, applications should be made when the buds are dormant (bud scales tight). Risk of injury increases as bud development progresses. Do not apply any later than cabbage head stage.
- Do not apply more than 12.0 fl oz/A of Spartan 4F Herbicide per year.
- Do not apply more than 0.375 lb sulfentrazone/A per year.
- Do not apply more than once per year.
- Pre-harvest Interval (PHI): 3 days.
- Do not apply additional sulfentrazone containing products to cranberry if Spartan 4F Herbicide has been previously applied within the same twelve-month period.
- The 12-month period is considered to begin upon the initial Spartan 4F Herbicide application.
- Do not apply to frozen soils or to existing snow cover to prevent Spartan 4F Herbicide runoff from rain or snow melt that may occur following application.

## **MANUFACTURER'S CONDITIONS AND RISK OF USE**

**NOTICE:** This 24(c) label will only be utilized by growers located in the state of Oregon. **SUCH GROWER USERS AGREE FMC CORPORATION WILL NOT BE HELD RESPONSIBLE FOR FAILURE TO PERFORM AND/OR CROP RESPONSE AND OR YIELD LOSSES RESULTING FROM THE USE OF SPARTAN 4F HERBICIDE IN CRANBERRY.** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability on the label affixed to the main container of Spartan 4F Herbicide and this SLN label before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product shall be followed carefully.

**USE OF SPARTAN 4F HERBICIDE ON CRANBERRY IN ACCORDANCE WITH THIS LABELING MAY RESULT IN FAILURE TO PERFORM, CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS. READ AND UNDERSTAND THESE CONDITIONS AND RISKS OF USE FOR SECTION 24(C) REGISTRATION BEFORE USING THIS PRODUCT ON THE CROP. FMC Corporation recommends that the user and/or grower test product in order to determine its suitability for such intended use.**

FMC Corporation makes the Product available for use in the manner described in this Special Local Need (SLN) on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of failure to perform, Crop injury, Crop yield reduction and/or Crop loss. The decision to use the Product in the manner described in this SLN Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the potential risk of failure to perform, Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls and (iv) any other relevant factors.

**By purchasing the Product for use, or using the Product, in the manner described in this SLN Labeling, you acknowledge and accept that, to the extent consistent with applicable law:**

- (1) You assume all risk of failure to perform, Crop injury, Crop yield reduction and Crop loss;
- (2) FMC Corporation does not make, and does not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Labeling;
- (3) FMC Corporation does not make, and does not authorize any agent or representative to make, any warranties, expressed or implied, with respect to the use of the Product on the Crop and disclaim all warranties, expressed or implied, including any implied warranty of fitness for a particular purpose or merchantability;
- (4) FMC Corporation disclaims all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to failure to perform, Crop injury, Crop yield reductions and/or Crop loss, including without limitation incidental, consequential or special damages;
- (5) These Conditions and Risks of Use registration supersede any contrary representations or

recommendations by FMC Corporation, its agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

**If these Conditions and Risk of Use are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container. These Conditions and Risks of Uses are required by FMC Corporation and not specified by the U.S. EPA or the State of Oregon.**

**Section 24(c) Registrant:**



**FMC Corporation**

2929 Walnut Street

Philadelphia Pennsylvania 19104

FMC, the FMC logo and Spartan are trademarks of FMC Corporation or an affiliate.

©2025 FMC Corporation All Rights Reserved

Label code DR-4759 030525