

FIFRA Section 24(c) Special Local Need Label
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON

TIAFENACIL	GROUP	14	HERBICIDE
------------	-------	----	-----------

TIAFENACIL 70WG

EPA Reg. No. 71512-36

SLN No. OR-230010

For Postemergence Weed Control in Non-bearing Hazelnuts

This SLN label is valid until December 31, 2028 or until otherwise amended, withdrawn, cancelled, or suspended

Active Ingredient

Tiafenacil*	70.0%
Other Ingredients	30.0%
Total	100.0%

*methyl N-[2-[[2-chloro-5-[3,6-dihydro-3-methyl-2,6-dioxo-4-(trifluoromethyl)-1(2H)-pyrimidinyl]-4-fluorophenyl]thio]-1-oxopropyl]-β-alaninate

TIAFENACIL 70WG is formulated as a water dispersible granule (WG) and contains 0.70 pounds of active ingredient per pound of formulated product.

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

SEE LABEL BOOKLET FOR FIRST AID, ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE INCLUDING STORAGE AND DISPOSAL

READ ENTIRE LABEL CAREFULLY AND USE ONLY AS DIRECTED.

ENVIRONMENTAL HAZARDS

Follow all applicable directions, restrictions, and precautions in the Environmental Hazards section of the label affixed to the main container of TIAFENACIL 70WG (EPA Reg. No. 71512-36).

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

This SLN label and the label affixed to the main container of TIAFENACIL 70WG must be in the possession of the user at the time of application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the label affixed to the main container of TIAFENACIL 70WG. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

PRODUCT USES & APPLICATION INSTRUCTIONS:

USE FOR:
Post-emergence weed and sucker control in non-bearing hazelnuts (only). Weeds controlled include Italian ryegrass, annual bluegrass, prostrate knotweed, and wild carrot.
SPRAY MIXTURE PREPARATION:
For sucker control, apply the labeled rate of TIAFENACIL 70WG in a minimum of 10 gallons of spray solution per acre. For weed control, apply the labeled rate of TIAFENACIL 70WG in 15-20 gallons of spray solution per acre.. Follow the instructions in the Sprayer Mixing: Mixing and Loading section of the label affixed to the main container of TIAFENACIL 70WG.

POSTEMERGENCE SUCKER AND WEED CONTROL IN NON-BEARING HAZELNUTS:		
Application Timing	Rate Range (oz/A)	Additional Information & Restrictions
Postemergence Burndown (Directed)	0.5 to 1.5 (0.022-0.067 lbs tiafenacil/acre)	<ul style="list-style-type: none"> Apply as a spray directed to the weeds or suckers, using conventional low-pressure ground spray equipment. Follow manufacture's recommendations for spraying pressure. Follow all applicable directions, restrictions and precautions related to spray drift management on the label affixed to the main container of TIAFENACIL 70WG. Do not allow spray solution to contact green stems (except suckers) or foliage. Do not apply to hazelnut established less than 1 year. DO NOT make the second application within 30 days of the first application. Do not make the third application within 14 days of the second application. Use higher rate for dense and/or mature weed infestations. Always apply product with an effective tank mixture partner, if electing to use the low rate. Refer to the label affixed to the main container of TIAFENACIL 70WG for tank mixing directions, restrictions, and precautions. To avoid crop injury, do not allow the spray solution to drift into the canopy.

Always use a methylated seed oil/crop oil concentrate (MSO/COC) or nonionic surfactant (NIS) at the recommended rate when applying Tiafenacil 70WG or reduced performance will occur. For adjuvant requirements, refer to the Adjuvants section of the label affixed to the main container of TIAFENACIL 70WG.

RESTRICTIONS:

- **DO NOT** apply more than 1.5 oz of TIAFENACIL 70WG (0.067 lbs of tiafenacil) per acre per application.
- **DO NOT** apply more than 4.5 oz. of TIAFENACIL 70WG (0.2 lbs of tiafenacil) per acre per year.
- **DO NOT** make more than 3 applications per year.
- This SLN label allows application of this product to non-bearing hazelnuts **ONLY**. **DO NOT** harvest hazelnuts for at least 1 year (365 days) after the last date this product was applied.
- **DO NOT** apply this product by aerial application.
- **DO NOT** apply this product through any type of irrigation system.
- Apply by ground application equipment only.

NOTICE

Read the "WARRANTY DISCLAIMER" and "LIMITATION OF LIABILITY" in the label booklet for this product before using this product. Those terms apply to this SLN label and if those terms are not acceptable, return the product unopened at once.

(RV050823)

[Section 24\(c\) Registrant:](#)
ISK Biosciences Corporation
7470 Auburn Road, Suite A
Concord, OH 44077